

| | Anthem BCBS Medicaid | CoventryCares of Kentucky | | | Humana CareSource | | | Kentucky Spirit Health Plan | | Passport Health Plan | | | WellCare of Kentucky | | | Kentucky Averages | | |
|--|----------------------|---------------------------|--------|--------|-------------------|--------|------|-----------------------------|--------|----------------------|--------|--------|----------------------|--------|--------|-------------------|--------|--------|
| | 2014 | 2014 | 2013 | 2012 | 2014 | 2013 | 2012 | 2013 | 2012 | 2014 | 2013 | 2012 | 2014 | 2013 | 2012 | 2014 | 2013 | 2012 |
| Effectiveness of Care | | | | | | | | | | | | | | | | | | |
| Prevention and Screening | | | | | | | | | | | | | | | | | | |
| Adult BMI Assessment (ABA) | | | | | | | | | | | | | | | | | | |
| ABA: Rate | N/A | 86.13% | 60.88% | N/A | 66.91% | N/A | | | N/A | 89.35% | 85.43% | 76.38% | 91.32% | 85.65% | N/A | | 77.09% | 76.38% |
| Weight Assessment & Counseling for Nutrition & Physical Activity for Children/Adolescents (WCC) | | | | | | | | | | | | | | | | | | |
| WCC: BMI Percentile Ages 3 - 11 years | N/A | 47.27% | 25.25% | 15.16% | 56.88% | 31.44% | | | 23.05% | 85.22% | 76.41% | 57.98% | 42.49% | 32.62% | 23.51% | | 38.84% | 29.90% |
| WCC: BMI Percentile Ages 12 - 17 years | N/A | 42.58% | 24.44% | 23.87% | 56.34% | 41.07% | | | 28.00% | 88.27% | 79.61% | 65.75% | 46.38% | 34.64% | 27.89% | | 40.11% | 36.24% |
| WCC: BMI Percentile Total | N/A | 45.50% | 25.00% | 18.29% | 56.69% | 34.06% | | | 24.77% | 86.31% | 77.48% | 60.49% | 43.80% | 33.33% | 25.00% | | 39.27% | 32.08% |
| WCC: Counseling for Nutrition Ages 3 - 11 years | N/A | 55.08% | 25.59% | 30.32% | 51.30% | 39.80% | | | 33.69% | 71.48% | 65.78% | 64.50% | 43.22% | 43.01% | 30.88% | | 41.13% | 40.09% |
| WCC: Counseling for Nutrition Ages 12 - 17 years | N/A | 45.16% | 20.00% | 29.68% | 46.48% | 47.32% | | | 30.67% | 75.31% | 68.42% | 63.01% | 37.68% | 43.79% | 31.29% | | 40.11% | 38.70% |
| WCC: Counseling for Nutrition Total | N/A | 51.34% | 23.84% | 30.09% | 49.64% | 41.85% | | | 32.64% | 72.85% | 66.67% | 64.02% | 41.36% | 43.29% | 31.02% | | 40.83% | 39.61% |
| WCC: Counseling for Physical Activity Ages 3 - 11 years | N/A | 40.63% | 21.21% | 21.66% | 47.58% | 31.77% | | | 21.63% | 60.82% | 46.51% | 34.85% | 39.56% | 36.92% | 23.86% | | 32.92% | 25.65% |
| WCC: Counseling for Physical Activity Ages 12 - 17 years | N/A | 41.29% | 23.70% | 29.03% | 46.48% | 49.11% | | | 27.33% | 68.52% | 65.79% | 64.38% | 42.03% | 49.67% | 40.14% | | 43.38% | 40.35% |
| WCC: Counseling for Physical Activity Ages Total | N/A | 40.88% | 21.99% | 24.31% | 47.20% | 36.50% | | | 23.61% | 63.58% | 52.98% | 44.37% | 40.39% | 41.44% | 29.40% | | 36.47% | 30.60% |
| Childhood Immunization Status (CIS) | | | | | | | | | | | | | | | | | | |
| CIS: DtaP/DT | N/A | 78.83% | 78.19% | 84.49% | 64.23% | 17.02% | | | 76.40% | 87.47% | 85.62% | 88.50% | 72.99% | 78.70% | 74.94% | | 80.01% | 84.66% |
| CIS: IPV | N/A | 89.29% | 91.18% | 92.82% | 77.86% | 31.91% | | | 87.85% | 94.43% | 95.80% | 97.79% | 86.86% | 89.35% | 87.47% | | 91.46% | 94.48% |
| CIS: MMR | N/A | 87.83% | 90.26% | 90.28% | 86.62% | 70.21% | | | 86.21% | 94.66% | 95.58% | 95.35% | 87.35% | 88.66% | 87.01% | | 90.91% | 92.37% |
| CIS: Hib | N/A | 87.35% | 93.04% | 92.59% | 82.00% | 53.19% | | | 86.68% | 96.29% | 95.35% | 97.79% | 85.89% | 90.74% | 87.70% | | 92.64% | 94.32% |
| CIS: Hepatitis B | N/A | 90.27% | 87.70% | 85.65% | 69.83% | 57.45% | | | 76.17% | 93.74% | 96.02% | 97.57% | 79.32% | 84.72% | 75.17% | | 88.56% | 90.22% |
| CIS: VZV | N/A | 89.54% | 92.11% | 90.74% | 86.37% | 70.21% | | | 88.32% | 95.36% | 95.35% | 95.58% | 90.02% | 90.74% | 89.10% | | 92.34% | 93.10% |
| CIS: Pneumococcal Conjugate | N/A | 80.29% | 79.12% | 78.01% | 64.48% | 12.77% | | | 72.66% | 86.77% | 86.50% | 87.83% | 75.67% | 80.79% | 73.32% | | 81.34% | 82.58% |
| CIS: Hepatitis A | N/A | 74.45% | 72.39% | 71.30% | 73.24% | 59.57% | | | 74.77% | 84.45% | 83.63% | 80.53% | 68.13% | 69.91% | 67.98% | | 74.20% | 76.80% |
| CIS: Rotavirus | N/A | 62.53% | 60.79% | 64.81% | 55.72% | 44.68% | | | 60.98% | 71.00% | 75.22% | 74.12% | 54.26% | 61.81% | 54.29% | | 64.65% | 68.54% |
| CIS: Influenza | N/A | 36.98% | 35.50% | 31.25% | 35.28% | 14.89% | | | 30.37% | 56.38% | 51.33% | 51.55% | 29.93% | 34.03% | 20.42% | | 38.78% | 41.83% |
| CIS: Combination 2 | N/A | 75.18% | 71.69% | 75.23% | 54.26% | 8.51% | | | 65.65% | 83.53% | 83.19% | 87.17% | 64.96% | 71.99% | 63.11% | | 74.43% | 79.60% |
| CIS: Combination 3 | N/A | 71.05% | 67.52% | 68.75% | 51.09% | 8.51% | | | 63.08% | 80.05% | 80.75% | 82.74% | 62.04% | 70.37% | 59.16% | | 71.63% | 75.15% |
| CIS: Combination 4 | N/A | 61.56% | 54.52% | 55.79% | 44.04% | 4.26% | | | 54.44% | 71.69% | 73.45% | 73.67% | 49.64% | 56.48% | 46.64% | | 59.73% | 65.09% |
| CIS: Combination 5 | N/A | 52.55% | 48.96% | 53.24% | 37.96% | 4.26% | | | 46.03% | 63.57% | 68.14% | 65.93% | 41.61% | 51.39% | 40.84% | | 54.42% | 58.31% |
| CIS: Combination 6 | N/A | 31.63% | 26.91% | 25.93% | 23.84% | 2.13% | | | 24.07% | 50.35% | 46.68% | 47.79% | 25.06% | 27.55% | 16.47% | | 31.92% | 37.49% |
| CIS: Combination 7 | N/A | 48.18% | 42.23% | 46.53% | 33.82% | 4.26% | | | 41.36% | 58.83% | 62.83% | 60.40% | 36.01% | 45.60% | 35.96% | | 48.40% | 52.79% |
| CIS: Combination 8 | N/A | 30.41% | 23.90% | 23.61% | 21.90% | 2.13% | | | 23.13% | 47.80% | 44.47% | 45.13% | 21.90% | 25.93% | 14.15% | | 29.62% | 35.14% |
| CIS: Combination 9 | N/A | 25.79% | 22.04% | 23.15% | 19.71% | 2.13% | | | 19.16% | 43.39% | 42.26% | 40.71% | 18.98% | 21.99% | 14.85% | | 26.92% | 32.03% |
| CIS: Combination 10 | N/A | 25.06% | 20.19% | 21.30% | 18.49% | 2.13% | | | 18.22% | 41.53% | 40.27% | 38.94% | 17.52% | 21.30% | 12.76% | | 25.47% | 30.31% |
| Immunizations for Adolescents (IMA) | | | | | | | | | | | | | | | | | | |
| IMA: Meningococcal | N/A | 83.07% | 76.62% | 74.31% | 69.83% | 71.15% | | | 54.44% | 88.89% | 86.75% | 75.00% | 80.77% | 81.99% | 79.86% | | 80.99% | 72.94% |
| IMA: Tdap/Td | N/A | 90.10% | 86.11% | 78.70% | 71.78% | 71.15% | | | 55.79% | 93.52% | 91.61% | 86.95% | 88.08% | 91.32% | 80.56% | | 89.24% | 80.93% |
| IMA: Combination #1 | N/A | 82.11% | 75.23% | 71.99% | 67.15% | 71.15% | | | 50.97% | 88.43% | 85.21% | 73.45% | 79.23% | 81.03% | 77.08% | | 79.72% | 70.87% |
| Lead Screening in Children (LSC) | | | | | | | | | | | | | | | | | | |
| LSC: Rate | N/A | 60.58% | 66.13% | 65.51% | 65.45% | 53.19% | | | 61.92% | 77.26% | 81.42% | 82.30% | 61.11% | 68.18% | 59.63% | | 70.59% | 74.29% |
| Breast Cancer Screening (BCS) | | | | | | | | | | | | | | | | | | |
| BCS: Rate | N/A | 51.58% | N/A | N/A | N/A | N/A | | | 51.67% | 56.95% | 59.25% | N/A | 51.47% | N/A | N/A | | 59.25% | 51.67% |
| Cervical Cancer Screening (CCS) | | | | | | | | | | | | | | | | | | |
| CCS: Rate | 25.29% | 49.15% | 51.41% | 47.89% | 35.52% | 37.96% | | | 38.67% | 55.53% | 72.36% | 64.11% | 48.18% | 48.93% | 46.28% | | 52.96% | 49.61% |
| Chlamydia Screening in Women (CHL) | | | | | | | | | | | | | | | | | | |
| CHL: Ages 16 - 20 years | 44.62% | 46.04% | 44.13% | 46.93% | 55.93% | 65.96% | | | 46.65% | 62.43% | 63.51% | 63.14% | 44.49% | 41.84% | 45.57% | | 47.82% | 50.92% |
| CHL: Ages 21 - 24 years | 53.00% | 56.98% | 58.71% | 53.98% | 54.51% | 69.12% | | | 63.99% | 65.16% | 72.14% | 69.62% | 54.68% | 54.46% | 53.15% | | 59.43% | 59.94% |
| CHL: Total Rate | 51.27% | 50.43% | 48.27% | 48.98% | 55.02% | 66.99% | | | 51.92% | 63.49% | 65.58% | 65.00% | 48.69% | 45.56% | 47.85% | | 51.08% | 53.58% |
| Human Papillomavirus Vaccine for Female Adolescents (HPV) | | | | | | | | | | | | | | | | | | |
| HPV: Rate | N/A | 15.82% | 16.32% | 13.92% | 13.25% | N/A | | | 3.39% | 30.63% | 28.95% | 29.40% | 20.44% | 15.28% | 11.81% | | 19.03% | 21.30% |
| Respiratory Conditions | | | | | | | | | | | | | | | | | | |
| Appropriate Testing for Children with Pharyngitis (CWP) | | | | | | | | | | | | | | | | | | |
| CWP: Rate | N/A | 65.23% | 62.72% | 63.95% | 78.39% | N/A | | | 60.19% | 84.46% | 80.54% | 73.57% | 67.27% | 64.47% | 64.74% | | 67.01% | 66.45% |

| | Anthem BCBS Medicaid | CoventryCares of Kentucky | Humana CareSource | Kentucky Spirit Health Plan | Passport Health Plan | WellCare of Kentucky | Kentucky Averages |
|---|----------------------|---------------------------|-------------------|-----------------------------|----------------------|----------------------|--|
| Appropriate Treatment for Children with Upper Respiratory Infection (URI) | | | | | | | |
| URI: Rate | N/A | 61.64% | 58.33% | 56.40% | 80.35% | 75.47% | 58.06% 81.16% 81.34% 77.74% 60.24% 59.17% 61.81% 62.75% 63.60% |
| Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis (AAB) | | | | | | | |
| AAB: Rate | 27.56 | 19.38% | 16.60% | 11.53% | 22.49% | N/A | 31.30% 19.38% 13.37% 31.99% 17.59% 30.45% 30.81% 22.81% 27.05% |
| Use of Spirometry Testing in the Assessment and Diagnosis of COPD (SPR) | | | | | | | |
| SPR: Rate | N/A | 23.09% | N/A | N/A | N/A | N/A | N/A 33.81% 38.15% 33.57% 23.27% N/A N/A 38.15% 33.57% |
| Pharmacotherapy Management of COPD Exacerbation (PCE) | | | | | | | |
| PCE: Systemic Corticosteroid | 63.73% | 64.20% | 65.10% | 66.74% | 57.78% | 43.42% | 37.64% 72.04% 76.78% 38.20% 65.07% 37.13% 35.37% 46.83% 44.25% |
| PCE: Bronchodilator | 71.48% | 76.11% | 80.66% | 84.36% | 69.05% | 59.21% | 45.29% 86.02% 91.63% 56.01% 79.61% 47.05% 45.36% 58.48% 57.47% |
| Use of Appropriate Medications for People with Asthma (ASM) | | | | | | | |
| ASM: Ages 5 - 11 years | N/A | 93.83% | 94.38% | N/A | 92.98% | N/A | N/A 94.36% 94.89% 92.95% 95.12% 94.79% N/A 94.64% 92.95% |
| ASM: Ages 12 - 18 years | N/A | 89.96% | 88.38% | N/A | 92.54% | N/A | N/A 88.96% 91.77% 87.98% 89.60% 87.72% N/A 89.16% 87.98% |
| ASM: Ages 19 - 50 years | N/A | 63.83% | 57.89% | N/A | 82.35% | N/A | N/A 73.11% 76.70% 67.74% 66.88% 66.67% N/A 64.21% 67.74% |
| ASM: Ages 51 - 64 years | N/A | 53.46% | 50.00% | N/A | N/A | N/A | N/A 65.05% 65.31% 58.02% 59.71% 64.74% N/A 58.31% 58.02% |
| ASM: Total Rate | N/A | 85.82% | 84.31% | N/A | 90.35% | N/A | N/A 88.48% 91.31% 87.88% 85.67% 85.64% N/A 86.53% 87.88% |
| Cardiovascular Conditions | | | | | | | |
| Cholesterol Management for Patients with Cardiovascular Conditions (CMC) | | | | | | | |
| CMC: LDL-C Screening | | | 79.63% | N/A | | N/A | N/A 79.47% 79.91% 80.09% N/A 79.89% 79.91% |
| CMC: LDL-C Level <100 mg/dL | | | 29.63% | N/A | | N/A | N/A 41.50% 44.59% 44.21% N/A 39.94% 44.59% |
| Controlling High Blood Pressure (CBP) | | | | | | | |
| CBP: Total Rate | 52.79% | 57.70% | 54.22% | 49.11% | 50.61% | 42.34% | 43.14% 51.66% 61.95% 62.97% 55.99% 54.98% 58.68% 55.34% 54.61% |
| Persistence of Beta Blocker Treatment After a Heart Attack (PBH) | | | | | | | |
| PBH: Rate | 71.43% | 83.13% | 89.87% | 87.50% | 79.73% | N/A | 59.65% 86.00% 94.44% 73.42% 82.95% 71.13% 72.62% 77.83% 75.11% |
| Diabetes | | | | | | | |
| Comprehensive Diabetes Care (CDC) | | | | | | | |
| CDC: HbA1c Testing | 90.89% | 87.04% | 83.08% | 80.37% | 88.87% | 83.92% | 80.79% 90.78% 86.59% 84.08% 88.78% 88.25% 86.64% 86.69% 83.38% |
| CDC: HbA1c Poor Control (>9.0%) | 51.14% | 39.78% | 45.73% | 52.82% | 66.97% | 69.48% | 62.03% 38.43% 36.28% 35.57% 40.61% 39.29% 44.54% 40.97% 47.42% |
| CDC: HbA1c Control (<8.0%) | 38.70% | 50.18% | 45.43% | 40.18% | 32.85% | 28.34% | 32.45% 50.61% 54.12% 55.97% 50.49% 51.04% 45.32% 49.69% 44.51% |
| CDC: HbA1c Control (<7.0%) | 26.20% | 36.08% | 36.38% | 32.08% | 21.67% | 21.18% | N/A 34.72% 40.68% 41.85% 39.47% 39.83% 32.58% 38.65% 35.00% |
| CDC: Eye Exam | 84.06% | 40.51% | 38.11% | 37.14% | 39.96% | 37.87% | 42.83% 40.70% 57.93% 52.74% 43.66% 35.25% 35.52% 39.07% 41.91% |
| CDC: LDL-C Screening | | | 78.66% | 71.39% | | 70.57% | 71.30% 77.13% 76.99% 84.22% 79.18% 81.67% 75.27% |
| CDC: LDL-C Level (<100 mg/dL) | | | 28.81% | 26.48% | | 22.07% | 23.18% 39.18% 42.54% 36.75% 35.08% 34.90% 32.80% |
| CDC: Medical Attention for Nephropathy | 84.06% | 80.84% | 79.57% | 75.34% | 84.49% | 75.20% | 67.99% 81.74% 80.64% 79.48% 82.68% 79.72% 80.62% 79.73% 76.67% |
| CDC: Blood Pressure Control (<140/80 mm Hg) | | | 32.16% | 34.25% | | 28.34% | 27.81% 37.96% 42.04% 39.63% 39.64% 37.39% 36.76% |
| CDC: Blood Pressure Control (<140/90 mm Hg) | 58.14% | 57.85% | 52.13% | 54.19% | 49.09% | 44.41% | 46.36% 66.43% 63.41% 64.68% 60.00% 58.76% 58.02% 57.53% 56.67% |
| Musculoskeletal | | | | | | | |
| Disease Modifying Anti-Rheumatic Drug Therapy in Rheumatoid Arthritis (ART) | | | | | | | |
| ART: Rate | 63.64% | 69.19% | 65.84% | 63.01% | 60.91% | N/A | 37.13% 61.14% 72.29% 44.21% 61.25% 42.88% 44.62% 50.84% 47.34% |
| Use of Imaging Studies for Low Back Pain (LBP) | | | | | | | |
| LBP: Rate | 55.18% | 64.34% | 69.18% | 68.84% | 56.97% | N/A | 68.30% 60.20% 62.69% 68.09% 61.48% 64.20% 66.38% 65.95% 67.96% |
| Behavioral Health | | | | | | | |
| Follow-up Care for Children Prescribed ADHD Medication (ADD) | | | | | | | |
| ADD: Initiation Phase | NR | 44.59% | 49.85% | N/A | 42.28% | N/A | N/A 44.84% 31.81% 29.12% 61.49% 64.86% N/A 47.65% 29.12% |
| ADD: Continuation and Maintenance Phase | NR | 52.35% | 52.68% | N/A | 54.55% | N/A | N/A 58.26% 41.67% 28.83% 69.62% 74.11% N/A 56.49% 28.83% |
| Diabetes Screening for People With Schizophrenia or Bipolar Disorder Who Are Using Antipsychotic Medications (SSD) | | | | | | | |
| SSD: Rate | 82.93% | 79.35% | 75.32% | 77.39% | 80.79% | 77.17% | 75.33% 89.30% 89.07% 75.96% 82.24% 78.15% 80.84% 78.11% 77.59% |
| Diabetes Monitoring for People With Diabetes And Schizophrenia (SMD) | | | | | | | |
| SMD: Rate | N/A | 68.89% | 71.21% | N/A | 58.70% | N/A | N/A 58.33% 54.41% N/A 70.08% 71.95% N/A 69.94% 66.27% |
| Cardiovascular Monitoring For People With Cardiovascular Disease and Schizophrenia (SMC) | | | | | | | |
| SMC: Rate | N/A | N/A | N/A | N/A | N/A | N/A | N/A N/A N/A N/A 82.14% 82.35% N/A 82.35% 80.00% |
| Adherence to Antipsychotic Medications for Individuals With Schizophrenia (SAA) | | | | | | | |
| SAA: Rate | 57.14% | 63.50% | 61.79% | N/A | 45.91% | N/A | N/A 56.09% 78.42% N/A 66.58% 64.70% N/A 65.09% 64.70% |
| Medication Management | | | | | | | |
| Annual Monitoring for Patients on Persistent Medications (MPM) | | | | | | | |

| | Anthem BCBS Medicaid | CoventryCares of Kentucky | | | Humana CareSource | | | Kentucky Spirit Health Plan | Passport Health Plan | | | WellCare of Kentucky | | | Kentucky Averages | | |
|---|----------------------|---------------------------|--------|--------|-------------------|--------|--|-----------------------------|----------------------|--------|--------|----------------------|--------|--------|-------------------|--------|--------|
| MPM: ACE inhibitors or ARBs | 90.15% | 89.34% | 88.97% | 89.51% | 90.02% | N/A | | 87.29% | 92.00% | 91.78% | 91.01% | 91.21% | 91.28% | 90.03% | | 90.65% | 89.61% |
| MPM: Digoxin | N/A | 47.93% | 90.24% | 90.72% | 50.00% | N/A | | 84.78% | 61.46% | 93.75% | 91.45% | 53.66% | 91.02% | 90.24% | | 91.21% | 89.77% |
| MPM: Diuretics | 90.62% | 89.54% | 89.80% | 89.01% | 90.36% | N/A | | 89.20% | 92.68% | 92.95% | 91.02% | 92.09% | 91.98% | 90.93% | | 91.51% | 90.06% |
| MPM: Anticonvulsants | | | 64.23% | 63.74% | | N/A | | 66.02% | | 61.93% | 59.94% | | 69.76% | 67.13% | | 66.36% | 64.06% |
| MPM: Total Rate | 90.10% | 89.07% | 87.07% | 86.82% | 89.93% | N/A | | 85.66% | 92.04% | 89.37% | 87.59% | 91.24% | 89.81% | 88.27% | | 88.88% | 87.21% |
| Medication Management for People With Asthma (MMA) | | | | | | | | | | | | | | | | | |
| MMA: 50% Ages 5 - 11 years | N/A | 73.71% | 67.98% | N/A | 68.87% | N/A | | N/A | 64.47% | 72.38% | 70.54% | 61.92% | 59.45% | N/A | | 66.71% | 70.54% |
| MMA: 50% Ages 12 - 18 years | N/A | 64.78% | 65.91% | N/A | 64.52% | N/A | | N/A | 61.13% | 65.53% | 63.53% | 56.72% | 57.07% | N/A | | 63.16% | 63.53% |
| MMA: 50% Ages 19 - 50 years | N/A | 69.44% | 66.77% | N/A | N/A | N/A | | N/A | 65.29% | 70.89% | 71.20% | 67.12% | 69.18% | N/A | | 68.49% | 71.20% |
| MMA: 50% Ages 51 - 64 years | N/A | 78.82% | 86.17% | N/A | N/A | N/A | | N/A | 68.66% | 93.55% | 91.30% | 77.71% | 80.49% | N/A | | 84.27% | 91.30% |
| MMA: 50% Total | N/A | 70.39% | 67.72% | N/A | 67.48% | N/A | | N/A | 63.51% | 70.03% | 68.37% | 61.65% | 61.08% | N/A | | 66.28% | 68.37% |
| MMA: 75% Ages 5 - 11 years | N/A | 50.52% | 42.92% | N/A | 43.40% | N/A | | N/A | 38.08% | 47.16% | 48.49% | 34.16% | 35.14% | N/A | | 41.83% | 48.49% |
| MMA: 75% Ages 12 - 18 years | N/A | 42.28% | 43.28% | N/A | 30.65% | N/A | | N/A | 32.35% | 39.10% | 42.29% | 29.35% | 32.44% | N/A | | 38.81% | 42.29% |
| MMA: 75% Ages 19 - 50 years | N/A | 46.29% | 46.11% | N/A | N/A | N/A | | N/A | 39.67% | 49.37% | 56.00% | 43.13% | 43.84% | N/A | | 45.92% | 56.00% |
| MMA: 75% Ages 51 - 64 years | N/A | 49.41% | 63.83% | N/A | N/A | N/A | | N/A | 52.24% | 67.74% | 71.74% | 59.64% | 65.04% | N/A | | 64.92% | 71.74% |
| MMA: 75% Total | N/A | 47.25% | 44.04% | N/A | 38.35% | N/A | | N/A | 36.67% | 44.65% | 47.17% | 35.03% | 37.01% | N/A | | 42.02% | 47.17% |
| Asthma Medication Ratio (AMR) | | | | | | | | | | | | | | | | | |
| AMR: 5-11 years | N/A | 85.85% | 85.71% | N/A | 75.45% | N/A | | N/A | 81.19% | 85.49% | 75.14% | 85.26% | 86.91% | N/A | | 86.00% | 75.14% |
| AMR: 12-18 years | N/A | 66.94% | 71.74% | N/A | 56.72% | N/A | | N/A | 68.19% | 75.66% | 66.79% | 72.87% | 74.26% | N/A | | 73.63% | 66.79% |
| AMR: 19-50 years | N/A | 39.43% | 38.81% | N/A | 57.58% | N/A | | N/A | 54.24% | 57.56% | 43.32% | 48.12% | 51.03% | N/A | | 46.37% | 43.32% |
| AMR: 51-64 years | N/A | 38.36% | 38.83% | N/A | N/A | N/A | | N/A | 49.51% | 55.10% | 40.74% | 45.68% | 50.53% | N/A | | 45.90% | 40.74% |
| AMR: Total Rate | N/A | 70.43% | 71.31% | N/A | 66.37% | N/A | | N/A | 71.77% | 78.43% | 68.23% | 71.61% | 74.22% | N/A | | 74.05% | 68.23% |
| Access/Availability of Care | | | | | | | | | | | | | | | | | |
| Adults' Access to Preventive/Ambulatory Health Services (AAP) | | | | | | | | | | | | | | | | | |
| AAP: Ages 20 - 44 years | 74.61% | 81.16% | 86.43% | 88.38% | 72.68% | 73.35% | | 82.03% | 82.58% | 90.45% | 85.12% | 84.36% | 88.90% | 88.15% | | 87.86% | 86.22% |
| AAP: Ages 45 - 64 years | 84.17% | 87.23% | 92.16% | 93.70% | 83.52% | 80.11% | | 86.64% | 89.39% | 94.78% | 90.68% | 91.05% | 94.48% | 93.26% | | 93.56% | 91.32% |
| AAP: Ages 65 years and older | N/A | 82.47% | 86.47% | 88.79% | 78.36% | 83.72% | | 86.78% | 94.00% | 93.21% | 92.07% | 88.94% | 94.51% | 93.68% | | 93.74% | 91.31% |
| AAP: Total Rate | 78.49% | 83.21% | 88.62% | 90.45% | 77.16% | 77.37% | | 84.42% | 85.17% | 92.39% | 88.22% | 86.91% | 91.98% | 90.97% | | 90.69% | 88.75% |
| Children and Adolescents' Access to Primary Care Practitioners (CAP) | | | | | | | | | | | | | | | | | |
| CAP: Ages 12 - 24 months | N/A | 97.20% | 95.46% | 97.94% | 92.39 | 93.40% | | 96.89% | 98.35% | 98.55% | 97.85% | 97.49% | 98.07% | 97.72% | | 97.20% | 97.65% |
| CAP: Ages 25 months - 6 years | N/A | 90.63% | 92.42% | 93.93% | 82.52 | 79.98% | | 91.34% | 90.25% | 92.00% | 89.37% | 92.02% | 93.02% | 93.61% | | 92.15% | 92.07% |
| CAP: Ages 7 - 11 years | N/A | 96.32% | 97.57% | N/A | 88.32% | N/A | | N/A | 94.19% | 94.70% | 91.95% | 96.30% | 97.47% | N/A | | 96.81% | 91.95% |
| CAP: Ages 12 - 19 years | N/A | 95.05% | 96.39% | N/A | 85.11% | N/A | | N/A | 92.92% | 93.95% | 91.64% | 95.22% | 96.45% | N/A | | 95.80% | 91.64% |
| Annual Dental Visits (ADV) | | | | | | | | | | | | | | | | | |
| ADV: Ages 2 - 3 years | N/A | 37.97% | 37.77% | 39.70% | 38.65% | 37.84% | | 29.45% | 48.63% | 47.01% | 42.27% | 42.32% | 43.98% | 39.89% | | 42.25% | 38.50% |
| ADV: Ages 4 - 6 years | N/A | 67.74% | 68.16% | 71.17% | 59.80% | 57.74% | | 57.94% | 72.94% | 72.68% | 67.93% | 69.99% | 72.85% | 70.86% | | 70.57% | 67.68% |
| ADV: Ages 7 - 10 years | N/A | 66.48% | 67.10% | 69.06% | 54.91% | 56.86% | | 56.70% | 70.46% | 71.57% | 67.87% | 69.13% | 72.96% | 70.14% | | 70.02% | 66.63% |
| ADV: Ages 11 - 14 years | N/A | 59.61% | 61.13% | 62.79% | 51.39% | 50.79% | | 49.64% | 66.23% | 67.76% | 63.59% | 63.74% | 67.55% | 63.96% | | 64.76% | 60.78% |
| ADV: Ages 15 - 18 years | N/A | 51.55% | 53.24% | 55.25% | 42.99% | 43.75% | | 45.13% | 58.80% | 61.57% | 57.11% | 55.38% | 60.33% | 56.97% | | 57.51% | 54.17% |
| ADV: Ages 19 - 21 years | 21.68% | 34.57% | 38.62% | 42.65% | 24.71% | 22.78% | | 30.50% | 40.66% | 45.67% | 39.64% | 38.20% | 42.72% | 38.60% | | 41.01% | 38.19% |
| ADV: Total Rate | 21.49% | 57.34% | 59.22% | 61.07% | 46.40% | 50.26% | | 48.87% | 63.64% | 65.48% | 60.95% | 60.36% | 65.00% | 61.79% | | 62.53% | 58.86% |
| Prenatal and Postpartum Care (PPC) | | | | | | | | | | | | | | | | | |
| PPC: Timeliness of Prenatal Care | 64.29% | 84.91% | 82.41% | N/A | 69.34% | 82.26% | | N/A | 86.89% | 91.61% | N/A | 87.29% | 86.54% | N/A | | 86.24% | 88.35% |
| PPC: Postpartum Care | 48.70% | 59.12% | 50.93% | N/A | 51.58% | 61.02% | | N/A | 68.67% | 67.33% | N/A | 51.41% | 52.44% | N/A | | 55.47% | 62.65% |
| Call Answer Timeliness (CAT) | | | | | | | | | | | | | | | | | |
| CAT: Rate | 87.75% | 78.20% | 85.31% | N/A | 90.82% | 93.01% | | N/A | 77.62% | 86.69% | N/A | 80.43% | 86.80% | N/A | | 86.43% | 78.43% |
| Use of Services | | | | | | | | | | | | | | | | | |
| Frequency of Ongoing Prenatal Care (FPC) | | | | | | | | | | | | | | | | | |
| FPC: <21 percent | 17.53% | 5.84% | 10.56% | N/A | 13.63% | 8.06% | | N/A | 5.81% | 3.75% | N/A | 4.52% | 6.03% | N/A | | 7.06% | 3.59% |
| FPC: 21 - 40 percent | 3.90% | 2.68% | 5.59% | N/A | 8.03% | 6.99% | | N/A | 2.33% | 1.99% | N/A | 1.98% | 4.41% | N/A | | 4.30% | 2.59% |
| FPC: 41 - 60 percent | 4.55% | 6.57% | 9.72% | N/A | 9.98% | 8.33% | | N/A | 4.65% | 4.86% | N/A | 5.93% | 4.41% | N/A | | 6.35% | 5.07% |
| FPC: 61 - 80 percent | 14.29% | 12.90% | 20.40% | N/A | 20.68% | 14.78% | | N/A | 12.09% | 11.04% | N/A | 15.25% | 15.55% | N/A | | 16.13% | 11.36% |
| FPC: >= 81 percent | 59.74% | 72.02% | 53.73% | N/A | 47.45% | 61.83% | | N/A | 75.12% | 78.37% | N/A | 72.32% | 69.61% | N/A | | 66.17% | 77.39% |
| Well-Child Visits in the First 15 Months of Life (W15) | | | | | | | | | | | | | | | | | |

| | Anthem BCBS Medicaid | CoventryCares of Kentucky | | | Humana CareSource | | | Kentucky Spirit Health Plan | Passport Health Plan | | | WellCare of Kentucky | | | Kentucky Averages | |
|--|----------------------|---------------------------|-----------|-----------|-------------------|---------|--|-----------------------------|----------------------|-----------|-----------|----------------------|-----------|-----------|-------------------|-----------|
| W15: 0 Visits | N/A | 2.43% | 2.78% | 1.16% | 2.72% | N/A | | 2.88% | 0.76% | 1.05% | 1.25% | 3.32% | 1.62% | 3.47% | 1.91% | 1.51% |
| W15: 1 Visit | N/A | 3.13% | 2.78% | 2.31% | 2.42% | N/A | | 4.40% | 1.27% | 0.62% | 1.46% | 5.61% | 0.69% | 3.47% | 1.45% | 1.87% |
| W15: 2 Visits | N/A | 4.91% | 3.71% | 3.47% | 5.44% | N/A | | 4.23% | 1.69% | 1.37% | 2.62% | 5.36% | 3.94% | 6.02% | 3.22% | 3.04% |
| W15: 3 Visits | N/A | 7.14% | 6.73% | 6.71% | 7.25% | N/A | | 8.80% | 4.13% | 3.68% | 4.06% | 6.89% | 6.25% | 9.03% | 5.79% | 4.94% |
| W15: 4 Visits | N/A | 12.06% | 10.44% | 7.87% | 13.29% | N/A | | 14.55% | 8.30% | 8.17% | 7.96% | 8.67% | 8.80% | 16.67% | 9.25% | 9.03% |
| W15: 5 Visits | N/A | 20.09% | 15.78% | 15.74% | 22.36% | N/A | | 23.18% | 17.62% | 14.76% | 14.66% | 22.96% | 16.90% | 18.75% | 15.96% | 15.63% |
| W15: >= 6 Visits | N/A | 50.24% | 57.77% | 62.73% | 46.53% | N/A | | 41.96% | 66.24% | 70.34% | 67.98% | 47.19% | 61.81% | 42.59% | 62.43% | 63.97% |
| Well-Child Visits in the Third, Fourth, Fifth and Sixth Years of Life (W34) | | | | | | | | | | | | | | | | |
| W34: Rate | N/A | 57.73% | 62.73% | 55.79% | 62.29% | 56.69% | | 58.14% | 74.77% | 75.47% | 70.68% | 59.75% | 56.31% | 61.81% | 62.75% | 61.82% |
| Adolescent Well-Care Visits (AWC) | | | | | | | | | | | | | | | | |
| AWC: Rate | 24.19% | 43.07% | 46.30% | 45.83% | 32.60% | 39.42% | | 35.28% | 51.37% | 57.99% | 52.46% | 41.85% | 43.75% | 38.89% | 47.48% | 43.98% |
| Frequency of Selected Procedures (FSP)* | | | | | | | | | | | | | | | | |
| FSP: Bariatric Weight Loss Surgery F Ages 0-19 Procs/1000 MM | | 0 | 0.00 | 0.00 | 0.00 | 0.00 | | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| FSP: Bariatric Weight Loss Surgery F Ages 20-44 Procs/1000 MM | | 0.05 | 0.09 | 0.15 | 0.01 | 0.04 | | 0.05 | 0.12 | 0.24 | 0.27 | 0.10 | 0.11 | 0.15 | 0.12 | 0.15 |
| FSP: Bariatric Weight Loss Surgery F Ages 45-64 Procs/1000 MM | | 0.03 | 0.11 | 0.13 | 0.02 | 0.22 | | 0.08 | 0.14 | 0.25 | 0.26 | 0.11 | 0.12 | 0.09 | 0.17 | 0.14 |
| FSP: Bariatric Weight Loss Surgery M Ages 0-19 Procs/1000 MM | | 0 | 0.00 | 0.00 | 0.00 | 0.00 | | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| FSP: Bariatric Weight Loss Surgery M Ages 20-44 Procs/1000 MM | | 0.01 | 0.05 | 0.03 | 0.01 | 0.00 | | 0.01 | 0.02 | 0.12 | 0.01 | 0.02 | 0.04 | 0.02 | 0.05 | 0.02 |
| FSP: Bariatric Weight Loss Surgery M Ages 45-64 Procs/1000 MM | | 0.02 | 0.00 | 0.02 | 0.01 | 0.00 | | 0.01 | 0.01 | 0.00 | 0.03 | 0.00 | 0.05 | 0.04 | 0.01 | 0.03 |
| FSP: Tonsillectomy MF Ages 0-9 Procs/1000 MM | | 0.85 | 1.07 | 1.46 | 0.56 | 0.57 | | 1.14 | 0.78 | 1.01 | 1.00 | 1.00 | 1.18 | 1.22 | 0.96 | 1.20 |
| FSP: Tonsillectomy MF Ages 10-19 Procs/1000 MM | | 0.36 | 0.41 | 0.70 | 0.24 | 0.33 | | 0.49 | 0.30 | 0.44 | 0.48 | 0.41 | 0.54 | 0.70 | 0.43 | 0.59 |
| FSP: Hysterectomy Abdominal F Ages 15-44 Procs/1000 MM | | 0.24 | 0.26 | 0.29 | 0.25 | 0.26 | | 0.20 | 0.26 | 0.19 | 0.26 | 0.24 | 0.25 | 0.31 | 0.24 | 0.26 |
| FSP: Hysterectomy Abdominal F Ages 45-64 Procs/1000 MM | | 0.4 | 0.35 | 0.28 | 0.43 | 0.43 | | 0.23 | 0.41 | 0.21 | 0.34 | 0.35 | 0.32 | 0.31 | 0.33 | 0.29 |
| FSP: Hysterectomy Vaginal F Ages 15-44 Procs/1000 MM | | 0.21 | 0.20 | 0.29 | 0.23 | 0.05 | | 0.20 | 0.39 | 0.22 | 0.24 | 0.31 | 0.34 | 0.33 | 0.20 | 0.26 |
| FSP: Hysterectomy Vaginal F Ages 45-64 Procs/1000 MM | | 0.2 | 0.08 | 0.19 | 0.29 | 0.00 | | 0.11 | 0.43 | 0.09 | 0.12 | 0.32 | 0.27 | 0.22 | 0.11 | 0.16 |
| FSP: Cholecystectomy, Open M Ages 30-64 Procs/1000 MM | | 0.05 | 0.02 | 0.03 | 0.04 | 0.00 | | 0.06 | 0.08 | 0.09 | 0.11 | 0.03 | 0.02 | 0.04 | 0.03 | 0.06 |
| FSP: Cholecystectomy, Open F Ages 15-44 Procs/1000 MM | | 0.02 | 0.02 | 0.02 | 0.04 | 0.05 | | 0.00 | 0.03 | 0.02 | 0.03 | 0.01 | 0.02 | 0.01 | 0.03 | 0.02 |
| FSP: Cholecystectomy Open F Ages 45-64 Procs/1000 MM | | 0.06 | 0.02 | 0.08 | 0.05 | 0.11 | | 0.05 | 0.07 | 0.02 | 0.15 | 0.05 | 0.02 | 0.05 | 0.04 | 0.08 |
| FSP: Cholecystectomy Closed M Ages 30-64 Procs/1000 MM | | 0.62 | 0.63 | 0.75 | 0.60 | 0.08 | | 0.64 | 0.64 | 0.34 | 0.55 | 0.73 | 0.67 | 0.72 | 0.43 | 0.67 |
| FSP: Cholecystectomy Closed F Ages 15-44 Procs/1000 MM | | 1.18 | 1.08 | 1.57 | 1.21 | 0.67 | | 1.31 | 1.06 | 0.89 | 0.87 | 1.47 | 1.39 | 1.60 | 1.01 | 1.34 |
| FSP: Cholecystectomy Closed F Ages 45-64 Procs/1000 MM | | 1.13 | 0.88 | 1.29 | 1.19 | 0.54 | | 0.97 | 1.05 | 0.80 | 0.76 | 1.20 | 1.14 | 1.20 | 0.84 | 1.06 |
| FSP: Back Surgery M Ages 20-44 Procs/1000 MM | | 0.35 | 0.30 | 0.41 | 0.49 | 0.32 | | 0.43 | 0.46 | 0.54 | 0.61 | 0.33 | 0.44 | 0.47 | 0.40 | 0.48 |
| FSP: Back Surgery F Ages 20-44 Procs/1000 MM | | 0.22 | 0.21 | 0.27 | 0.23 | 0.47 | | 0.29 | 0.36 | 0.47 | 0.40 | 0.27 | 0.33 | 0.31 | 0.37 | 0.32 |
| FSP: Back Surgery M Ages 45-64 Procs/1000 MM | | 0.58 | 0.47 | 0.59 | 0.76 | 1.18 | | 0.75 | 0.85 | 1.15 | 0.96 | 0.58 | 0.67 | 0.64 | 0.87 | 0.73 |
| FSP: Back Surgery F Ages 45-64 Procs/1000 MM | | 0.44 | 0.47 | 0.56 | 0.57 | 1.19 | | 0.57 | 0.84 | 0.93 | 1.16 | 0.56 | 0.65 | 0.78 | 0.81 | 0.77 |
| FSP: Mastectomy F Ages 15-44 Procs/1000 MM | | 0.02 | 0.01 | 0.02 | 0.02 | 0.00 | | 0.04 | 0.04 | 0.02 | 0.03 | 0.06 | 0.02 | 0.05 | 0.01 | 0.03 |
| FSP: Mastectomy F Ages 45-64 Procs/1000 MM | | 0.12 | 0.13 | 0.13 | 0.10 | 0.32 | | 0.16 | 0.23 | 0.09 | 0.10 | 0.19 | 0.14 | 0.13 | 0.17 | 0.13 |
| FSP: Lumpectomy F Ages 15-44 Procs/1000 MM | | 0.14 | 0.15 | 0.16 | 0.17 | 0.05 | | 0.13 | 0.18 | 0.20 | 0.21 | 0.16 | 0.15 | 0.18 | 0.14 | 0.17 |
| FSP: Lumpectomy F Ages 45-64 Procs/1000 MM | | 0.25 | 0.38 | 0.36 | 0.41 | 0.22 | | 0.46 | 0.47 | 0.43 | 0.31 | 0.37 | 0.33 | 0.40 | 0.34 | 0.38 |
| Inpatient Utilization - General Hospital/Acute Care: Total (IPUA)* | | | | | | | | | | | | | | | | |
| IPUA: Total Discharges/1000 MM | | 9.14 | 9.85 | 11.16 | 9.57 | 116.27 | | 11.30 | 10.23 | 10.21 | 12.19 | 10.31 | 12.87 | 16.43 | 37.30 | 12.77 |
| IPUA: Medicine Discharges/1000 MM | | 4.08 | 4.08 | 5.54 | 5.72 | 99.29 | | 4.40 | 4.24 | 3.86 | 6.05 | 4.84 | 6.40 | 8.49 | 28.41 | 6.12 |
| IPUA: Surgery Discharges/1000 MM | | 1.39 | 1.22 | 1.69 | 2.27 | 8.80 | | 2.46 | 2.99 | 2.05 | 2.68 | 2.56 | 2.83 | 3.14 | 3.72 | 2.49 |
| IPUA: Maternity Discharges/1000 MM | | 5.6 | 8.08 | 6.81 | 0.08 | 13.32 | | 7.68 | 4.37 | 8.12 | 6.46 | 4.21 | 6.30 | 8.35 | 8.95 | 7.32 |
| Antibiotic Utilization: Total (ABXA)* | | | | | | | | | | | | | | | | |
| ABXA: Total # of Antibiotic Prescriptions M&F | | 581,787 | 494,491 | 555,955 | 103,836 | 25,910 | | 228,615 | 246,773 | 159,967 | 272,204 | 619,303 | 438,288 | 406,686 | 279,664 | 365,865 |
| ABXA: Average # of Antibiotic Prescriptions PMPY M&F | | 2 | 2.31 | 2.71 | 1.36 | 1.40 | | 1.54 | 1.34 | 1.43 | 1.58 | 1.64 | 1.68 | 2.45 | 1.70 | 2.07 |
| ABXA: Total Days Supplied for all Antibiotic Prescriptions M&F | | 5,377,518 | 4,560,637 | 5,093,854 | 928,053 | 238,654 | | 2,020,874 | 2,253,360 | 1,500,889 | 2,507,341 | 5,674,046 | 4,067,483 | 2,956,948 | 2,591,915 | 3,144,754 |
| ABXA: Average # Days Supplied per Antibiotic Prescription M&F | | 9.24 | 9.22 | 9.16 | 8.94 | 9.21 | | 8.84 | 9.13 | 9.38 | 9.21 | 9.16 | 9.28 | 7.27 | 9.27 | 8.62 |
| ABXA: Total # of Prescriptions for Antibiotics of Concern M&F | | 277,069 | 236,136 | 277,842 | 49,264 | 11,134 | | 108,771 | 101,897 | 65,028 | 115,066 | 305,832 | 210,527 | 199,808 | 130,706 | 175,372 |
| ABXA: Average # of Prescriptions for Antibiotics of Concern M&F | | 0.95 | 1.10 | 1.35 | 0.65 | 0.60 | | 0.73 | 0.55 | 0.58 | 0.67 | 0.81 | 0.81 | 1.20 | 0.77 | 0.99 |
| ABXA: Percent Antibiotics of Concern of all Antibiotic Prescriptions | | 47.62% | 47.75% | 49.98% | 47.44 | 42.97% | | 47.58% | 41.29% | 40.65% | 42.27% | 49.38% | 48.03% | 49.13% | 44.85% | 47.24% |
| Health Plan Descriptive Information | | | | | | | | | | | | | | | | |
| Board Certification (BCR)* | | | | | | | | | | | | | | | | |
| BCR: % of Family Medicine Board Certified | NR | 67.70% | 63.91% | 49.18% | 41.18% | 51.52% | | N/A | 24.86% | 76.70% | 78.39% | 45.11% | 46.52% | 38.49% | 59.66% | 55.35% |

| | Anthem BCBS Medicaid | CoventryCares of Kentucky | | | Humana CareSource | | | Kentucky Spirit Health Plan | Passport Health Plan | | | WellCare of Kentucky | | | Kentucky Averages | |
|--|----------------------|---------------------------|---------|---------|-------------------|---------|--|-----------------------------|----------------------|---------|---------|----------------------|---------|---------|-------------------|---------|
| BCR: % of Internal Medicine Board Certified | NR | 77.23% | 73.73% | 73.03% | 65.28% | 65.17% | | N/A | 39.86% | 77.93% | 77.91% | 47.87% | 50.54% | 41.89% | 66.84% | 64.28% |
| BCR: % of OB/GYNs Board Certified | NR | 67.95% | 70.85% | 66.89% | 72.22% | 52.96% | | N/A | 30.38% | 70.89% | 71.04% | 45.77% | 48.94% | 43.52% | 60.91% | 60.48% |
| BCR: % of Pediatricians Board Certified | NR | 73.57% | 78.75% | 78.81% | 78.55% | 73.79% | | N/A | 43.29% | 78.03% | 79.69% | 36.06% | 38.79% | 39.00% | 67.34% | 65.83% |
| BCR: % of Geriatricians Board Certified | NR | 58.06% | 57.14% | 58.62% | 20.00% | 16.67% | | N/A | 47.37% | 100.00% | 100.00% | 63.16% | 58.54% | 63.33% | 58.09% | 73.98% |
| BCR: % of Other Physician Specialists Board Certified | NR | 81.84% | 73.48% | 70.66% | 68.44% | 59.23% | | N/A | 16.69% | 68.19% | 67.19% | 38.37% | 39.53% | 38.30% | 60.11% | 58.72% |
| Follow-Up after Hospitalization (FUH) | | | | | | | | | | | | | | | | |
| FUH Follow-up Hosp Men Illness-7 Day | 20.52% | 31.98% | 38.98% | 41.81% | 20.99 | 15.50% | | 30.32% | 25.00% | 1.56% | N/A | 33.82% | 36.07% | 35.92% | 35.30% | 36.60% |
| FUH Follow-up Hosp Men Illness-30 Day | 37.02% | 55.82% | 63.37% | 69.15% | 39.35 | 28.68% | | 54.56% | 49.55% | 9.38% | N/A | 57.18% | 61.79% | 61.74% | 59.43% | 62.55% |
| Mental Health Utilization (MPT) | | | | | | | | | | | | | | | | |
| MPT:MH Svs Any Tot F Pct | | 7.76% | 8.76% | 15.20% | 10.91% | 10.65% | | 13.91% | 1.94% | 2.64% | N/A | 9.35% | 10.94% | 17.70% | 8.64% | 15.63% |
| MPT:MH Svs Any Tot M Pct | | 8.33% | 10.17% | 15.65% | 9.99% | 12.41% | | 15.04% | 4.15% | 6.14% | N/A | 9.47% | 11.30% | 17.79% | 9.97% | 16.15% |
| MPT:MH Svs Any Tot Pct | | 8.02% | 9.41% | 15.40% | 10.47% | 11.47% | | 14.41% | 2.95% | 4.24% | N/A | 9.41% | 11.10% | 17.74% | 9.25% | 15.87% |
| MPT:MH Svs Intensive Tot F Pct | | 0.20% | 0.26% | 0.32% | 6.77% | 3.01% | | 0.29% | 0.00% | 0.00% | N/A | 0.57% | 0.74% | 0.76% | 0.51% | 0.45% |
| MPT:MH Svs Intensive Tot M Pct | | 0.23% | 0.28% | 0.37% | 6.37% | 3.26% | | 0.49% | 0.00% | 0.00% | N/A | 0.64% | 0.80% | 0.86% | 0.54% | 0.56% |
| MPT:MH Svs Intensive Tot Pct | | 0.21% | 0.27% | 0.35% | 6.58% | 3.12% | | 0.38% | 0.00% | 0.00% | N/A | 0.60% | 0.77% | 0.81% | 0.52% | 0.50% |
| MPT:MH Svs Inpat Tot F Pct | | 0.87% | 1.00% | 1.09% | 1.09% | 2.71% | | 1.10% | 0.12% | 0.04% | N/A | 1.23% | 1.31% | 1.33% | 1.01% | 1.17% |
| MPT:MH Svs Inpat Tot M Pct | | 0.90% | 1.02% | 1.12% | 1.48% | 3.21% | | 1.11% | 0.15% | 0.04% | N/A | 1.27% | 1.27% | 1.39% | 1.01% | 1.20% |
| MPT:MH Svs Inpat Tot Pct | | 0.88% | 1.01% | 1.11% | 1.28% | 2.94% | | 1.10% | 0.13% | 0.04% | N/A | 1.25% | 1.29% | 1.36% | 1.01% | 1.19% |
| MPT:MH Svs Outpat Tot F Pct | | 7.38% | 8.31% | 14.93% | 10.42% | 7.58% | | 13.61% | 1.83% | 2.61% | N/A | 8.74% | 10.28% | 17.28% | 8.10% | 15.31% |
| MPT:MH Svs Outpat Tot M Pct | | 7.98% | 9.73% | 15.32% | 9.23% | 9.25% | | 14.73% | 4.01% | 6.11% | N/A | 8.77% | 10.64% | 17.33% | 9.43% | 15.79% |
| MPT:MH Svs Outpat Tot Pct | | 7.65% | 8.97% | 15.11% | 9.85% | 8.35% | | 14.11% | 2.82% | 4.21% | N/A | 8.76% | 10.44% | 17.30% | 8.70% | 15.53% |
| Weeks of Pregnancy (WOP) | | | | | | | | | | | | | | | | |
| WOP:Wks of Preg <0 week Pct | | 24.67% | 21.83% | 10.66% | 20.78% | 5.45% | | N/A | 19.43% | 12.09% | 19.92% | 25.79% | 17.88% | 10.37% | 18.50% | 15.28% |
| WOP:Wks of Preg 1-12 weeks Pct | | 43.65% | 32.69% | 25.97% | 30.75% | 12.19% | | N/A | 32.55% | 30.40% | 41.88% | 41.74% | 35.45% | 26.20% | 33.15% | 33.43% |
| WOP:Wks of Preg 13-27 weeks Pct | | 18.14% | 22.74% | 35.97% | 22.04% | 20.96% | | N/A | 29.05% | 30.58% | 21.01% | 18.99% | 25.26% | 34.07% | 25.80% | 31.64% |
| WOP:Wks of Preg 28+ weeks Pct | | 7.36% | 16.83% | 19.28% | 20.63% | 56.68% | | N/A | 11.55% | 19.85% | 10.17% | 7.43% | 15.82% | 22.15% | 21.63% | 18.53% |
| WOP:Wks of Preg Tot all Pregs Pct | | 100.00% | 100.00% | 100.00% | 100.00% | 100.00% | | N/A | 100.00% | 100.00% | 100.00% | 100.00% | 100.00% | 100.00% | 100.00% | 100.00% |
| WOP:Wks of Preg Unknown Pct | | 6.19% | 5.91% | 8.13% | 5.80% | 4.71% | | N/A | 7.43% | 7.08% | 7.01% | 6.06% | 5.60% | 7.22% | 6.06% | 7.49% |
| Antidepressant Medication Management (AMM) | | | | | | | | | | | | | | | | |
| AMM:Rate - Effect.Acute Phase Treatment | 69.55% | 62.35% | 59.33% | 64.54% | 68.35% | N/A | | 45.75% | 62.67% | 58.82% | 60.50% | 52.29% | 49.86% | 57.70% | 55.23% | 58.36% |
| AMM:Rate - Effect.Continuation Phase Treat. | 62.05% | 46.78% | 42.59% | 44.44% | 59.71% | N/A | | 29.74% | 46.83% | 42.45% | 46.60% | 37.61% | 33.53% | 46.23% | 38.71% | 42.98% |
| Initiation and Engagement of Alcohol and other Drug (IET) | | | | | | | | | | | | | | | | |
| IET:Rate - Engagement - 13-17 Years | N/A | 12.69% | 9.17% | 15.56% | N/A | N/A | | 22.38% | 2.56% | 1.76% | N/A | 16.76% | 19.60% | 16.61% | 11.93% | 17.89% |
| IET:Rate - Engagement - 18+ Years | 9.89% | 7.09% | 5.27% | 6.17% | N/A | N/A | | 6.52% | 2.91% | 1.76% | N/A | 5.73% | 4.46% | 6.05% | 4.36% | 6.24% |
| IET:Rate - Engagement - Total | 9.87% | 7.43% | 5.63% | 6.94% | N/A | N/A | | 7.64% | 2.89% | 1.76% | N/A | 6.16% | 5.35% | 6.76% | 4.95% | 7.10% |
| IET:Rate - Initiation - 13-17 Years | N/A | 37.86% | 31.66% | 39.80% | N/A | N/A | | 43.71% | 13.92% | 6.47% | N/A | 41.26% | 38.04% | 43.25% | 29.24% | 41.99% |
| IET:Rate - Initiation - 18+ Years | 34.29% | 32.68% | 31.47% | 34.39% | N/A | N/A | | 42.64% | 25.92% | 33.19% | N/A | 30.36% | 24.62% | 37.98% | 28.01% | 38.13% |
| IET:Rate - Initiation - Total | 34.32% | 32.99% | 31.48% | 34.84% | N/A | N/A | | 42.72% | 25.18% | 30.22% | N/A | 30.79% | 25.42% | 38.33% | 28.10% | 38.42% |
| Identification of Alcohol and other Drug Services (IAD) | | | | | | | | | | | | | | | | |
| IAD:ID Svs Outpat Tot Pct | | 3.52% | 2.84% | 3.41% | 6.91% | 0.17% | | 3.75% | 3.89% | 1.73% | N/A | 5.24% | 3.54% | 4.49% | 2.88% | 3.85% |
| IAD:D Svs Any Tot Pct | | 3.94% | 3.25% | 3.74% | 8.36% | 0.56% | | 4.43% | 4.61% | 2.17% | N/A | 6.16% | 4.08% | 5.13% | 3.35% | 4.38% |
| IAD:ID Svs Intensive Tot Pct | | 0.07% | 0.03% | 0.04% | 1.34% | 0.02% | | 0.04% | 0.00% | 0.00% | N/A | 0.10% | 0.07% | 0.07% | 0.04% | 0.05% |
| IAD:ID Svs Inpat Tot Pct | | 0.82% | 0.69% | 0.68% | 2.47% | 0.42% | | 1.31% | 1.30% | 0.70% | N/A | 1.73% | 0.87% | 1.31% | 0.77% | 1.06% |
| Ambulatory Care:(AMBA) | | | | | | | | | | | | | | | | |
| AMB ER <1 Visit/1000 | | 108.57 | 114.16 | 122.28 | 130.38 | 6.80 | | 116.01 | 125.49 | 125.67 | 134.76 | 108.65 | 112.49 | 121.71 | 113.21 | 123.91 |
| AMB ER 1-9 Visit/1000 | | 59.72 | 63.02 | 66.60 | 64.99 | 2.81 | | 64.40 | 61.89 | 62.39 | 66.97 | 62.40 | 65.06 | 68.29 | 61.73 | 66.63 |
| AMB ER 10-19 Visit/1000 | | 54.3 | 54.68 | 59.59 | 61.17 | 3.60 | | 58.33 | 54.94 | 51.64 | 52.15 | 56.39 | 56.66 | 60.20 | 53.23 | 57.58 |
| AMBA: TOTAL Outpatient Visits/1000 MM | | 571.16 | 660.84 | 818.21 | 415.84 | 336.28 | | 473.11 | 367.69 | 398.89 | 469.38 | 546.89 | 594.18 | 769.57 | 573.72 | 645.76 |
| AMBA: TOTAL Emergency Department Visits/1000 MM | | 79.32 | 78.72 | 85.99 | 92.71 | 5.42 | | 83.29 | 84.33 | 76.23 | 81.30 | 85.15 | 83.58 | 86.85 | 78.11 | 84.45 |
| Non Recommended Cervical Cancer Screening (NCS) | | | | | | | | | | | | | | | | |
| NCS: Rate | 13.13% | 7.86% | 9.62% | N/A | 6.37% | 0.33% | | N/A | 6.00% | 6.98% | N/A | 7.57% | 9.84% | N/A | 8.88% | |