CDC

KENTUCKY

Children Aged <5 Years

2006

Pediatric Nutrition
Surveillance System

Graphics

Comparing Contributor and National Data

<u>2005</u> (prior year) national PedNSS data are presented.

Contributors included 41 states, the District of Columbia, 6 tribal governments, and 1 U.S. territory.

Source of data

among children aged <5 years



* *Special Supplemental Nutrition Program for Women, Infants and Children.*
** *Early Periodic Screening, Diagnosis, and Treatment Program.*
\+ *Title V Maternal and Child Health Program.*
++ *Includes Head Start.*

2006 KY PedNSS Table 1C

Racial and ethnic distribution

among children aged <5 years



2006 KY PedNSS Table 1C

Age distribution

among children aged <5 years



2006 KY PedNSS Table 1C

Income distribution*

among children aged <5 years



* *Percent of poverty level.*

** *Adjunctive eligibility: eligible for Medicaid, Food Stamps, or other entitlement programs.*

2006 KY PedNSS Table 1C

Program participation*

among children aged <5 years



* *Represents program participation among children enrolled in the PH program contributing PedNSS data.*

** *Special Supplemental Nutrition Program for Women, Infants and Children.*

*** *Temporary Assistance for Needy Families.*

2006 KY PedNSS Table 1C

Migrant status*

among children aged <5 years



* *Family whose principal employment is agricultural on a seasonal basis; adapted from Federal Register 50744, 9/99.*

2006 KY PedNSS Table 1C

Prevalence of low birthweight*

by race and ethnicity



* *< 2500 grams, among infants born during the reporting period.*

** *Reduce low birthweight to 5% of live births.*

2006 KY PedNSS Table 8C

Trends in prevalence of low birthweight*

by race and ethnicity



* *< 2500 grams, among infants born during the reporting period.*
Year 2010 target: reduce low birthweight to 5% of live births.

2006 KY PedNSS Table 18C

Prevalence of high birthweight*

by race and ethnicity



* *> 4000 grams, among infants born during the reporting period.*

2006 KY PedNSS Table 8C

Trends in prevalence of high birthweight*

by race and ethnicity



* *> 4000 grams, among infants born during the reporting period.*

2006 KY PedNSS Table 18C

Prevalence of short stature, underweight, and overweight*

among children aged <5 years



* *Short stature: < 5th percentile length or height-for-age; underweight: < 5th percentile weight-for-length or BMI-for-age; overweight: ≥ 95th percentile weight-for-length or BMI-for-age, CDC Growth Charts, 2000.*

2006 KY PedNSS Table 2C

Prevalence of short stature*

among children aged <5 years, by race and ethnicity



* *< 5th percentile length or height-for-age, CDC Growth Charts, 2000.*

** *Reduce growth retardation (short stature) among low-income children under age five years to 5%.*

2006 KY PedNSS Table 8C

Prevalence of short stature*

among children aged <5 years, by age



* *< 5th percentile length or height-for-age, CDC Growth Charts, 2000.*

** *Reduce growth retardation (short stature) among low-income children under age five years to 5%.*

2006 KY PedNSS Table 8C

Trends in prevalence of short stature*

among children aged <5 years, by race and ethnicity



* *< 5th percentile length or height-for-age, CDC Growth Charts, 2000.*
Year 2010 target: reduce growth retardation (short stature) among low-income children under age five years to 5%.

2006 KY PedNSS Table 18C

Prevalence of underweight*

among children aged <5 years, by race and ethnicity



* *< 5th percentile weight-for-length or BMI-for-age, CDC Growth Charts, 2000. 5% of children are expected to fall below the 5th percentile.*

2006 KY PedNSS Table 8C

Prevalence of underweight*

among children aged <5 years, by age



* *< 5th percentile weight-for-length or BMI-for-age, CDC Growth Charts, 2000.*
5% of children are expected to fall below the 5th percentile.

2006 KY PedNSS Table 8C

Trends in prevalence of underweight*

among children aged <5 years, by race and ethnicity



* *< 5th percentile weight-for-length or BMI-for-age, CDC Growth Charts, 2000.*
5% of children are expected to fall below the 5th percentile.

2006 KY PedNSS Table 18C

Prevalence of overweight*

among children aged <5 years, by race and ethnicity



* *≥ 95th percentile weight-for-length or BMI-for-age, CDC Growth Charts, 2000.*
5% of children are expected to fall above the 95th percentile.

2006 KY PedNSS Table 8C

Prevalence of overweight*

among children aged <5 years, by age



* *≥ 95th percentile weight-for-length or BMI-for-age, CDC Growth Charts, 2000.*
5% of children are expected to fall above the 95th percentile.

2006 KY PedNSS Table 8C

Trends in prevalence of overweight*

among children aged <5 years, by race and ethnicity



* *≥ 95th percentile weight-for-length or BMI-for-age, CDC Growth Charts, 2000.*
5% of children are expected to fall above the 95th percentile.

2006 KY PedNSS Table 18C

Trends in prevalence of overweight*

among children aged <5 years, by age



* *≥ 95th percentile weight-for-length or BMI-for-age, CDC Growth Charts, 2000.*
5% of children are expected to fall above the 95th percentile.

2006 KY PedNSS Table 20C

Prevalence of overweight and risk of overweight*

among children aged 2 to <5 years, by race and ethnicity



* *Overweight: ≥ 95th percentile BMI-for-age; at risk of overweight: ≥ 85th-<95th percentile BMI-for-age, CDC Growth Charts, 2000. 15% of children are expected to fall above the 85th percentile (5% above the 95th percentile and 10% between the 85th and 95th percentiles).*

2006 KY PedNSS Table 8C

Prevalence of overweight and risk of overweight*

among children aged 2 to <5 years, by age



* *Overweight: ≥ 95th percentile BMI-for-age; at risk of overweight: ≥ 85th-<95th percentile BMI-for-age, CDC Growth Charts, 2000. 15% of children are expected to fall above the 85th percentile (5% above the 95th percentile and 10% between the 85th and 95th percentiles).*

2006 KY PedNSS Table 8C

Trends in prevalence of at risk of overweight

among children aged 2 to <5 years, by race and ethnicity



* *≥ 85th-<95th percentile BMI-for-age, CDC Growth Charts, 2000.*
10% of children are expected to fall between these percentiles.

2006 KY PedNSS Table 18C

Trends in prevalence of overweight

among children aged 2 to <5 years, by race and ethnicity



* *≥ 95th percentile BMI-for-age, CDC Growth Charts, 2000.*
5% of children are expected to fall above the 95th percentile.

2006 KY PedNSS Table 18C

Prevalence of anemia*

among children aged <5 years, by race and ethnicity



* *Hb or Hct < 5th percentile, CDC MMWR vol. 47 (No. RR-3), 1998.*

2006 KY PedNSS Table 8C

Prevalence of anemia*

among children aged <5 years, by age



* *Hb or Hct < 5th percentile, CDC MMWR vol. 47 (No. RR-3), 1998.*

2006 KY PedNSS Table 21C

Trends in prevalence of anemia*

among children aged <5 years, by race and ethnicity



* *Hb or Hct < 5th percentile, CDC MMWR vol. 47 (No. RR-3), 1998.*

2006 KY PedNSS Table 18C

Trends in prevalence of anemia*

among children aged <5 years, by age



* *Hb or Hct < 5th percentile, CDC MMWR vol. 47 (No. RR-3), 1998.*

2006 KY PedNSS Table 21C

Percentage of infants ever breastfed*

by race and ethnicity



* *Among infants born during the reporting period.*

** *Increase the proportion of mothers who breastfeed their babies in the early postpartum period to 75%.*

2006 KY PedNSS Table 9C

Trends in the percentage of infants ever breastfed*

by race and ethnicity



* *Among infants born during the reporting period.*
Year 2010 target: increase the proportion of mothers who breastfeed their babies in the early postpartum period to 75%.

2006 KY PedNSS Table 19C

Percentage of infants breastfed at least 6 months*

by race and ethnicity



* *Among infants who turned six months of age during the reporting period.*

** *Year 2010 target: increase the proportion of mothers who breastfeed their babies at six months to 50%.*

2006 KY PedNSS Table 9C

Trends in the percentage of infants breastfed at least 6 months*

by race and ethnicity



* *Among infants who turned six months of age during the reporting period.*
Year 2010 target: increase the proportion of mothers who breastfeed their babies at six months to 50%.

2006 KY PedNSS Table 19C

Percentage of infants breastfed at least 12 months*

by race and ethnicity



* *Among infants who turned twelve months of age during the reporting period.*

** *Increase the proportion of mothers who breastfeed their babies at one year to 25%.*

2006 KY PedNSS Table 9C

Trends in the percentage of infants breastfed at least 12 months*

by race and ethnicity



* *Among infants who turned twelve months of age during the reporting period.*
Year 2010 target: increase the proportion of mothers who breastfeed their babies at one year to 25%.

2006 KY PedNSS Table 19C

Trends in the percentage of infants ever breastfed, and breastfed at least 6 and 12 months



Year 2010 targets: increase the proportion of mothers who breastfeed their babies a) in the early postpartum period to 75%, b) at six months to 50%, and c) at one year to 25%.

2006 KY PedNSS Table 13C

Percentage of infants exclusively breastfed at least 3 and 6 months*

Insufficient Data

** Among infants who turned three and six months of age respectively during the reporting period.*

2006 KY PedNSS Table 3C

Percentage of children aged 2 to <5 years who view TV ≤2 hours/day

by race and ethnicity



* *Increase the proportion of children and adolescents who view TV two or fewer hours per day to 75%.*

2006 KY PedNSS Table 9C

Percentage of children aged 2 to <5 years who view TV ≤2 hours/day

by age



* *Increase the proportion of children and adolescents who view TV two or fewer hours per day to 75%.*

2006 KY PedNSS Table 9C

Percentage of children aged <5 years who live in households with smokers*

by race and ethnicity



* *Defined as smokers who smoke inside the home.*

** *Reduce the proportion of children who are regularly exposed to tobacco smoke at home to 10%.*

2006 KY PedNSS Table 9C

Percentage of children aged <5 years who live in households with smokers*

by age



* *Defined as smokers who smoke inside the home.*

** *Reduce the proportion of children who are regularly exposed to tobacco smoke at home to 10%.*

2006 KY PedNSS Table 9C

Graphics

Comparing Contributor and Local Data

Racial and ethnic distribution

among children aged <5 years, by region*



* *Data for some regions not presented if <100 records are available for analysis after exclusions.*

2006 KY PedNSS Table 4E

Age distribution

among children aged <5 years, by region*



* *Data for some regions not presented if <100 records are available for analysis after exclusions.*

2006 KY PedNSS Table 5E

Prevalence of low birthweight*

by region**



* *< 2500 grams, among infants born during the reporting period.*

** *Data for some regions not presented if <100 records are available for analysis after exclusions.*

*** *Reduce low birthweight to 5% of live births.*

2006 KY PedNSS Table 6E

Prevalence of high birthweight*

by region**



* *> 4000 grams, among infants born during the reporting period.*

** *Data for some regions not presented if <100 records are available for analysis after exclusions.*

2006 KY PedNSS Table 6E

Prevalence of short stature*

among children aged <5 years, by region**



* *< 5th percentile length or height-for-age, CDC Growth Charts, 2000.*

** *Data for some regions not presented if <100 records are available for analysis after exclusions.*

*** *Reduce growth retardation (short stature) among low-income children under age five years to 5%.*

2006 KY PedNSS Table 6E

Prevalence of underweight*

among children aged <5 years, by region**



* *< 5th percentile weight-for-length or BMI-for-age, CDC Growth Charts, 2000.*
5% of children are expected to fall below the 5th percentile.

** *Data for some regions not presented if <100 records are available for analysis after exclusions.*

2006 KY PedNSS Table 6E

Prevalence of overweight*

among children aged <5 years, by region**



* *≥ 95th percentile weight-for-length or BMI-for-age, CDC Growth Charts, 2000.*
5% of children are expected to fall above the 95th percentile.

** *Data for some regions not presented if <100 records are available for analysis after exclusions.*

2006 KY PedNSS Table 6E

Prevalence of overweight and risk of overweight*

among children aged 2 to <5 years, by region**



* *At risk of overweight: ≥ 85th-<95th percentile BMI-for-age. Overweight: ≥ 95th percentile BMI-for-age, CDC Growth Charts, 2000. 5% of children are expected to fall above the 95th percentile; 15% of children are expected to fall above the 85th percentile.*

** *Data for some regions not presented if <100 records are available for analysis after exclusions.*

2006 KY PedNSS Table 6E

Prevalence of anemia*

among children aged <5 years, by region**



* *Hb or Hct < 5th percentile, CDC MMWR vol. 47 (No. RR-3), 1998.*

** *Data for some regions not presented if <100 records are available for analysis after exclusions.*

2006 KY PedNSS Table 6E

Percentage of infants ever breastfed*

by region**



* *Among infants born during the reporting period.*

** *Data for some regions not presented if <100 records are available for analysis after exclusions.*

*** *Increase the proportion of mothers who breastfeed their babies in the early postpartum period to 75%.*

2006 KY PedNSS Table 7E

Percentage of infants breastfed at least 6 months*

by region**



* *Among infants who turned six months of age during the reporting period.*

** *Data for some regions not presented if <100 records are available for analysis after exclusions.*

*** *Year 2010 target: increase the proportion of mothers who breastfeed their babies at six months to 50%.*

2006 KY PedNSS Table 7E

Percentage of infants breastfed at least 12 months*

by region**



* *Among infants who turned twelve months of age during the reporting period.*

** *Data for some regions not presented if <100 records are available for analysis after exclusions.*

*** *Increase the proportion of mothers who breastfeed their babies at one year to 25%.*

2006 KY PedNSS Table 7E

Percentage of infants exclusively breastfed at least 3 months*

by region**

Insufficient Data

* *Among infants who turned three months of age during the reporting period.*

** *Data for some regions not presented if <100 records are available for analysis after exclusions.*

2006 KY PedNSS Table 7E

Percentage of children aged 2 to <5 years who view TV ≤2 hours/day

by region*



* *Data for some regions not presented if <100 records are available for analysis after exclusions.*

** *Increase the proportion of children and adolescents who view TV two or fewer hours per day to 75%.*

2006 KY PedNSS Table 7E

Percentage of children aged <5 years who live in households with smokers*

by region**



* *Defined as smokers who smoke inside the home.*

** *Data for some regions not presented if <100 records are available for analysis after exclusions.*

*** *Reduce the proportion of children who are regularly exposed to tobacco smoke at home to 10%.*

2006 KY PedNSS Table 7E

Maps

State Maps of County Data

Prevalence of low birthweight*

by county



** < 2500 grams, among infants born during the reporting period.*
Year 2010 target: reduce low birthweight to 5% of live births.

2006 KY PedNSS Table 6B

Prevalence of high birthweight*

by county



** > 4000 grams, among infants born during the reporting period.*

2006 KY PedNSS Table 6B

Prevalence of short stature*

among children aged <5 years, by county



** < 5th percentile length or height-for-age, CDC Growth Charts, 2000.*
Year 2010 target: reduce growth retardation (short stature) among low-income children under age five years to 5%.

2006 KY PedNSS Table 6B

Prevalence of underweight*

among children aged <5 years, by county



** < 5th percentile weight-for-length or BMI-for-age, CDC Growth Charts, 2000.*
5% of children are expected to fall below the 5th percentile.

2006 KY PedNSS Table 6B

Prevalence of overweight*

among children aged 2 to <5 years, by county



* ≥ 95th percentile BMI-for-age, CDC Growth Charts, 2000.
5% of children are expected to fall above the 95th percentile.
Year 2010 target: reduce overweight among children aged six to nineteen years to 5%.

2006 KY PedNSS Table 6B

Prevalence of anemia*

among children aged <5 years, by county



** Hb or Hct < 5th percentile, CDC MMWR vol. 47 (No. RR-3), 1998.*

2006 KY PedNSS Table 6B

Percentage of infants ever breastfed*

by county

Insufficient Data

* *Among infants born during the reporting period.*
Year 2010 target: increase the proportion of mothers who breastfeed their babies in the early postpartum period to 75%.

2006 KY PedNSS Table 7B

Percentage of infants breastfed at least 6 months*, by county

Insufficient Data

** Among infants who turned six months of age during the reporting period.*
Year 2010 target: increase the proportion of mothers who breastfeed their babies at six months to 50%.

2006 KY PedNSS Table 7B

Percentage of infants breastfed at least 12 months*, by county

Insufficient Data

** Among infants who turned twelve months of age during the reporting period.*
Year 2010 target: increase the proportion of mothers who breastfeed their babies at one year to 25%.

2006 KY PedNSS Table 7B

Percentage of infants exclusively breastfed at least 3 months*, by county

Insufficient Data

** Among infants who turned three months during the reporting period.*

2006 KY PedNSS Table 7B

Percentage of children aged 2 to <5 years who view TV ≤2 hours/day, by county



Year 2010 target: increase the proportion of children and adolescents who view TV two or fewer hours per day to 75%.

2006 KY PedNSS Table 7B

Percentage of children aged <5 years who live in households with smokers*, by county



** Defined as smokers who smoke inside the home.*
Year 2010 target: reduce the proportion of children who are regularly exposed to tobacco smoke at home to 10%.

2006 KY PedNSS Table 7B

CDC