



**Commonwealth of Kentucky
Cabinet for Health and Family Services**

FOR IMMEDIATE RELEASE

Contact: Barbara Fox
502-564-6786, ext. 3102

Beth Fisher
502-564-6786, ext.3101

Don't Let the Flu Catch Up With You

FRANKFORT, Ky. (Oct. 26, 2018) - The Department for Public Health (DPH), within the Cabinet for Health and Family Services, is reminding Kentuckians to get vaccinated against the flu.

"Getting the flu can be debilitating and sometimes life-threatening," said Jeffrey Howard, M.D., commissioner of DPH. "Vaccination is the best tool we have to prevent the flu and it is also extremely important to take simple preventive steps to avoid the flu and other illnesses that tend to circulate at this time of year – wash your hands frequently, cover your mouth when you cough or sneeze and stay home when you're sick."

DPH officials report weekly influenza activity to the CDC as part of statewide flu surveillance efforts. The weekly report is located online at <https://healthalerts.ky.gov/Pages/FluActivity.aspx> will be updated by noon each Friday. Kentucky currently is reporting 28 laboratory-confirmed cases of flu and one influenza-related death. The state flu activity level is currently classified as "sporadic". The report consists of laboratory-confirmed cases of influenza that are defined by molecular virus testing and positive virus culture test results. Rapid positive influenza tests are not

included in this report, but are used as an indicator of flu-like illness circulating across the state.

The Centers for Disease Control and Prevention (CDC) Advisory Committee on Immunization Practices (ACIP) recommends flu vaccine for all individuals six months of age and older. People who are strongly encouraged to receive the flu vaccine because they may be at higher risk for complications or negative consequences include:

- Children age six months through 59 months; • Women who are or will be pregnant during the influenza season; • Persons 50 years of age or older;
- Persons with extreme obesity (Body Mass Index of 40 or greater); • Persons aged six months and older with chronic health problems; • Residents of nursing homes and other long-term care facilities; • Household contacts (including children) and caregivers of children aged ≤ 59 months (i.e., aged <five years, particularly contacts of children aged <six months) and adults aged ≥ 50 years;
- Household contacts and caregivers or people who live with a person at high-risk for complications from the flu; and
- Health care workers, including physicians, nurses, and other workers in inpatient and outpatient-care settings, medical emergency-response workers (e.g., paramedics and emergency medical technicians), employees of nursing home and long-term care facilities who have contact with patients or residents, and students in these professions who will have contact with patients.

Adequate supplies of flu vaccine are expected to be available for this year's flu season. Vaccination can be given any time during the flu season, but providers are encouraged to administer the vaccine as soon as possible.

During the 2016-17 and 2017-18 influenza seasons, ACIP recommended that Flu Mist not be used because of concerns about low effectiveness against influenza A (H1N1). However, ACIP recommends that FluMist once again be an option for vaccination during the 2018-2019 influenza season for persons for whom it is age-appropriate and who

have no medical contraindications. The change in recommendation is a result of the influenza A (H1N1) strain being changed in the current FluMist formulation.

Due to the timing of the decision by the ACIP regarding use of FluMist, the Kentucky Vaccines for Children Program (VFC) will not be offering FluMist for VFC or KCHIP patients during the 2018-2019 influenza season. However, some private providers will offer FluMist this season for their patients.

The flu vaccine is especially important in light of the severe season experienced last year across the U.S. The CDC reports that more than 900,000 flu-related hospitalizations occurred and more than 80,000 people died as a result of flu. In Kentucky there were 333 flu-related deaths, 5 of which were pediatric. Of the pediatric flu deaths reported in last year's season, more than one-third of these occurred among healthy children between the ages of 6 months and 4 years.

It is especially important for pregnant women to be vaccinated before or during pregnancy. Antibodies cross the placenta and provide immunity to infants up to 6 months of age, when the infant is eligible to receive their dose of flu vaccine.

DPH relies on sites such as doctors' offices, hospitals and health departments to help track the level of influenza activity in the state and to identify which strains of the flu are circulating in Kentucky. These voluntary sites collect data and report influenza-like illness (ILI) cases according to age groups each week. This sampling represents only a small percentage of influenza cases for the state, but contributes to the ongoing assessment of flu activity in the Commonwealth and helps determine the weekly level of flu activity.

Infection with the flu virus can cause fever, headache, cough, sore throat, runny nose, sneezing and body aches. Flu can be very contagious. For more information on influenza or the availability of flu vaccine, Kentuckians should contact their primary care medical provider or local health department. Influenza information is also available online at <https://www.cdc.gov/flu/index.htm>.

The Cabinet for Health and Family Services is home to most of the state's human services and healthcare programs, including the Department for Medicaid Services, the Department for Community Based Services the Department for Public Health, the Department for Aging and Independent Living and the Department for Behavioral Health, Developmental and Intellectual Disabilities. CHFS is one of the largest agencies in state government, with nearly 8,000 full- and part-time employees located across the Commonwealth focused on improving the lives and health of Kentuckians.