



# HIV/AIDS Surveillance Report 2024

Kentucky Cabinet for Health and Family Services  
Department for Public Health  
HIV/AIDS Section

(Data complete through 2022, preliminary for 2023)

Dear Reader:

Enclosed please find Kentucky's HIV/AIDS Annual Surveillance Report 2024, which contains data on HIV infections among Kentuckians reported to the Department for Public Health. This annual edition is a Centers for Disease Control and Prevention (CDC) grant deliverable and is produced to fulfill the requirements of Funding Opportunity Announcement (FOA): PS24-0047.

Confidential AIDS reporting started in 1982, whereas legislation requiring confidential HIV name-based reporting was not enacted until July 2004. Prior to July 2004, HIV infections were reported with a unique code.

Section I (and throughout the report) profiles HIV infections diagnosed among Kentuckians, regardless of progression to AIDS. A total of 12,418 cumulative HIV infections were diagnosed and reported as of December 31, 2023. Of these HIV infections, 59% have progressed to AIDS as of the report date.

Section II profiles new HIV infections diagnosed among Kentuckians. In calendar year 2022, there were 405 new HIV infections diagnosed among Kentucky residents, a diagnosis rate of 9.0 per 100,000. This is an increase from the rate of 8.7 per 100,000 population for 2021. Trends among people with newly diagnosed infections are presented in this section, and disparities by race/ethnicity, age at diagnosis, sex and mode of transmission are highlighted. As per CDC guidance, the data for the last two years (2023 and 2024) are considered preliminary and not included in the trends analysis.

Section III profiles Kentuckians with HIV infection who were diagnosed with AIDS within 30 days of initial HIV diagnosis, also referred to as concurrent diagnoses. Analyses focus on the most recent 10-year period: January 1, 2014, through December 31, 2023. Twenty-two percent (22%) of the 3,568 individuals with new HIV disease diagnoses within that period were diagnosed with AIDS within 30 days of the initial HIV diagnosis.

Please read the data sources and technical notes on pages 3-5 for further information concerning interpretation of the data. The data presented in this report are available at <https://chfs.ky.gov/agencies/dph/dehp/hab/Documents/AnnualReport2024.pdf>.

Sincerely,

Manny Singh, MBBS, MPH  
Senior Epidemiologist  
HIV/AIDS Section

Release Date: 01/31/2025



**HIV/AIDS Surveillance Report**  
**Kentucky HIV/AIDS Section**  
**Division of Epidemiology and Health Planning**  
**Department for Public Health**  
**Cabinet for Health and Family Services**

**Address:**

Kentucky Department for Public Health  
HIV/AIDS Section  
275 East Main Street, HS2E-C  
Frankfort, KY 40621

**Phone:** (502) 564-6539 or (800) 420-7231

**Case Reporting Only:** (866) 510-0008

**Kentucky AIDS Drug Assistance Program (KADAP):** (866) 510-0005

**Website:** <https://chfs.ky.gov/agencies/dph/dehp/hab/Pages/default.aspx>

**Suggested citation:** Kentucky Cabinet for Health and Family Services. Department for Public Health. HIV/AIDS Surveillance Report, 2024; vol. 24. <https://chfs.ky.gov/agencies/dph/dehp/hab/Pages/default.aspx>. Published January 31, 2025. Accessed [date].

**For all media inquiries, please call the Office of Communications at (502) 564-6786 for assistance.**

**Kentucky Department for Public Health HIV/AIDS useful links:**

**HIV Reporting and Statistics**

**Fillable Adult HIV Confidential Case Report Form:**

[https://chfs.ky.gov/agencies/dph/dehp/hab/Documents/ACRF\\_Fillable.pdf](https://chfs.ky.gov/agencies/dph/dehp/hab/Documents/ACRF_Fillable.pdf)

**Fillable Pediatric HIV Confidential Case Report Form:**

[https://chfs.ky.gov/agencies/dph/dehp/hab/Documents/PCRF\\_Fillable.pdf](https://chfs.ky.gov/agencies/dph/dehp/hab/Documents/PCRF_Fillable.pdf)

**HIV Prevention**

**Syringe Services Programs:**

<https://chfs.ky.gov/agencies/dph/dehp/hab/Pages/kyseps.aspx>

**HIV Test Sites in Kentucky:**

<https://chfs.ky.gov/agencies/dph/dehp/hab/Documents/KYHIVTestSites.pdf>

**HIV Services**

**HIV Care Coordinator Regions and Contact Information:**

<https://chfs.ky.gov/agencies/dph/dehp/hab/Documents/KYHIVCCRs.pdf>

**Ryan White Services Eligibility Application:**

<https://chfs.ky.gov/agencies/dph/dehp/hab/Documents/RWEligApp.pdf>



## Data Sources

The HIV/AIDS Annual Report presents data regarding HIV disease cases diagnosed among Kentuckians and reported to the Kentucky Department for Public Health's HIV/AIDS Surveillance Program through December 31, 2023. In this annual edition, HIV disease cases diagnosed among Kentuckians are presented, regardless of disease progression. The data only include those persons who have been confidentially tested and reported to the HIV/AIDS Surveillance Program. No adjustments are made to the data presented to account for undiagnosed, anonymously tested or unreported cases.

Kentucky population estimates used in the calculation of rates were obtained from the Kentucky State Data Center. Source: Population Division, U.S. Census Bureau, 2020, available at <http://www.ksdc.louisville.edu/>. Accessed September 13, 2024.

## HIV/AIDS Reporting Requirements

According to state regulation 902 KAR 2:020 Reportable Disease Surveillance, Section 16, health professionals licensed under KRS Chapters 311 through 314, health facilities licensed under KRS 216B.015(13), and medical laboratories licensed under KRS Chapter 333, are required to report HIV and AIDS cases to the Kentucky Department for Public Health within five business days of diagnosis.

Cases of confirmed HIV and AIDS are reported to the Kentucky Department for Public Health's HIV/AIDS Surveillance Program at 866-510-0008 on the Confidential Adult HIV Case Report form for patients  $\geq 13$  years of age and on Pediatric Case Report form for patients less than 13 years of age at the time of diagnosis. Data from the case report forms are compiled to produce this report. Additional case reporting information can be found on the Kentucky HIV/AIDS Section Website: <https://chfs.ky.gov/agencies/dph/dehp/hab/Pages/reportsstats.aspx>.

## Key Terminology

The terminology used in this report is in a format consistent with CDC's technical guidelines for HIV surveillance grantees in the United States, and also consistent with the National HIV Surveillance Report, available online at: <https://www.cdc.gov/hiv-data/index.html>.

**Current Age:** An individual's age or age group as of December 31, 2023.

**Age at Diagnosis:** An individual's age or age group at the time of initial HIV disease diagnosis.

**Adults and Adolescents:** An individual aged 13 years and older.

**Pediatric:** An individual aged less than 13 years.

**Acquired Immunodeficiency Syndrome (AIDS):** Advanced stage of HIV infection characterized by severe immune deficiency and diagnosed by the presence of at least one of 26 opportunistic illnesses or a CD4 T-lymphocyte count of less than 200 cells/ml of blood. The CD4 T-lymphocyte count takes precedence over the CD4 T-lymphocyte percentage, and a percentage of less than 14% is considered only if the count is missing.

**Concurrent Diagnosis:** Both HIV and AIDS are diagnosed within a 30-day period.

**Date of Diagnosis:** The date of an individual's initial HIV disease diagnosis.

**Human Immunodeficiency Virus (HIV):** A retrovirus that infects the helper T cells of the immune system resulting in immunodeficiency. HIV is diagnosed by a positive confirmatory antibody test or positive/detectable viral detection test.



**Race and Ethnicity:** Ethnicity categories include Hispanic and not Hispanic. Data for all not Hispanic persons are displayed in combination with their racial groupings, which include:

- White
- Black or African American
- Asian
- Native Hawaiian or other Pacific Islander
- American Indian or Alaska Native

Kentucky's HIV data are collected for all racial and ethnic groupings. However, due to small numbers, data for the following racial groups are aggregated into the "other" designation: American Indian or Alaska Native, Native Hawaiian or Pacific Islander and persons of multiple races.

**Sex:** Sex designations in this report are based on a person's sex assignment at birth. In May 2012, CDC issued guidance to state and local programs on methods for collecting data on transgender persons and working with transgender-specific data. However, characterization of HIV infection among transgender persons in Kentucky would require supplemental data from special studies.

**Transmission Category:** Classification used to summarize the behavior or event most likely responsible for disease transmission. Each case is only included in a single transmission route.

**Male-to-Male Sexual Contact (MMSC):** Men who report having sexual contact with other men.

**Injection Drug Use (IDU):** Individuals who report injecting nonprescription drugs.

**MMSC/IDU:** Men who report having sex with other men and also inject nonprescription drugs.

**Heterosexual Contact:** A person reporting specific heterosexual contact with a person known to have, or to be at high risk for HIV infection, such as an injection drug user, a bisexual male (females only), or a person with hemophilia/coagulation disorder.

**Female Heterosexual Contact (FHC):** A female who does not fit in the heterosexual contact category above, with no reported injection drug use, but reported sexual contact with a male and no additional information about the male's HIV status or behaviors.

**Hemophilia:** Individuals receiving clotting factor for hemophilia/coagulation disorder.

**Perinatal:** Individuals born to a mother with HIV or a mother with an exposure history listed in the transmission category hierarchy.

**Blood Transfusion/Organ Transplant:** Individuals who received blood transfusions or organ transplants. Individuals with a transfusion date listed after March 1985 are considered Cases of Public Health Importance (COPHI) and are followed to verify the mode of transmission.

**Other:** Individuals who had a transfusion/transplant, hemophilia/coagulation disorder or pediatric cases diagnosed as adults.

**Undetermined/No Identified Risk (NIR):** Individuals reporting no exposure history to HIV through any of the modes listed in the transmission category hierarchy above.



## Technical Notes

**Reporting Delays:** Delays exist between the time HIV infection is diagnosed and the time the infection is reported to the HIV/AIDS Surveillance Program. As a result of reporting delays, case statistics for the most recent years of diagnosis may not be complete. Therefore, the data for 2023 and 2024 are considered provisional and will not be presented in the analysis of trends. The data presented in this report have not been adjusted for reporting delays.

**Place of Residence:** HIV data are presented based on residence at the time the initial HIV infection was diagnosed. Data presented on living cases reflect those originally diagnosed while living in Kentucky that are still presumed to be living, regardless of their current residence.

**Vital Status:** Cases are presumed to be alive unless the HIV/AIDS Surveillance Program has received notification of death. Current vital status information for cases is ascertained through routine site visits with major reporting sites, reports of death from providers, reports of death from other states' surveillance programs, routine matches with Kentucky death certificates (vital statistics registry) and Social Security Death Master Files (SSDMF).

**Transmission Route:** Despite the possible existence of multiple methods through which HIV was transmitted, cases are assigned a single most likely transmission route based on a hierarchy developed by the CDC. A limitation of the dataset is the large number of cases reported with an undetermined transmission route. Currently, surveillance data are collected through hard copy case reports, telephone reports and chart reviews which occasionally result in missing information. Enhanced surveillance activities have been implemented to resolve case reports with missing risk factor information, including the re-classification of females into the FHC category.

**Routine Interstate Duplicate Review (RIDR):** Case duplication between states can occur and has become more of an issue due to the mobility of our society. To help resolve duplicate reporting, CDC initiated the RIDR project in 2004. RIDR compares patient records throughout the nation in order to identify duplicate cases. The states with duplicate cases contact one another to compare patient profiles to assign the case one state residency based on residence at the earliest date of diagnosis. Due to this process, the cumulative number of cases within Kentucky may change, but the process has increased the accuracy of Kentucky's data by reducing the chance that a case has been counted more than once nationally.

**Small Numbers:** Data release limitations are set to ensure that the information cannot be used to inadvertently identify an individual. Data suppression rules are applied based on the population denominators for analyses below the state level. Additional numerator suppression rules are applied for groups or geographic areas that have <50,000 population. Rates are not released when the numerator is fewer than ten cases because of the low reliability of rates based on the small number of cases.

### **Difference between HIV Infection/HIV Disease, HIV without AIDS, and concurrent diagnosis of HIV with AIDS:**

HIV infection includes all individuals diagnosed with HIV regardless of the stage of disease progression. This term is used interchangeably with HIV disease. The data are presented based on the date of the first diagnosis reported to the HIV/AIDS Surveillance Program. HIV without AIDS includes individuals who were diagnosed with HIV and had not progressed to AIDS as of the report date. Concurrent diagnosis with AIDS includes those who were diagnosed with AIDS within 30 days of initial HIV diagnosis.



## Section I: Cumulative and Living HIV Infections Diagnosed as of December 31, 2023, Kentucky

<b>Table 1. Cumulative <sup>(1)</sup> HIV Disease Cases by Age at Diagnosis*, Race/Ethnicity and Sex at Birth as of December 31, 2023, Kentucky</b>											
	Age Group	White, Not Hispanic		Black, Not Hispanic		Hispanic		Other/Unknown		TOTAL	
		No.	%**	No.	%**	No.	%**	No.	%**	No.	%**
<b>MALE</b>	<13	25	<1	29	1	0	0	4	1	58	1
	13-19	146	2	181	6	13	2	26	6	366	4
	20-29	1,802	29	1,074	37	250	41	170	42	3,296	32
	30-39	2,186	35	839	29	220	36	108	27	3,353	33
	40-49	1,415	22	541	18	89	14	63	16	2,108	21
	50+	735	12	267	9	45	7	31	8	1,078	11
	<b>TOTAL</b>	<b>6,309</b>	<b>100</b>	<b>2,931</b>	<b>100</b>	<b>617</b>	<b>100</b>	<b>402</b>	<b>100</b>	<b>10,259</b>	<b>100</b>
<b>FEMALE</b>	<13	12	1	19	2	4	4	2	2	37	2
	13-19	48	5	57	6	7	7	2	2	114	5
	20-29	274	27	269	29	43	43	38	31	624	29
	30-39	333	33	293	31	20	20	42	34	688	32
	40-49	205	20	178	19	18	18	28	23	429	20
	50+	132	13	115	12	9	9	11	9	267	12
	<b>TOTAL</b>	<b>1,004</b>	<b>100</b>	<b>931</b>	<b>100</b>	<b>101</b>	<b>100</b>	<b>123</b>	<b>100</b>	<b>2,159</b>	<b>100</b>

(1) Includes HIV disease cases diagnosed from the beginning of the epidemic as of December 31, 2023.

\*Age at initial HIV diagnosis.

\*\*Percentages may not total 100% due to rounding.

Since the beginning of the HIV epidemic in 1982, the majority (83%) of HIV cases diagnosed among Kentuckians have been reported among males (10,259 cases). In terms of age at time of diagnosis, more male HIV cases were diagnosed at ages 30-39 (3,353 or 33%) than any other age grouping. Among white males, the highest percentage of cumulative cases were aged 30-39 years at the time of diagnosis (35%). Among black males, 37% of cases were aged 20-29 years and 29% were aged 30-39 years at time of diagnosis. The percentage of Hispanic males aged 20-29 at time of diagnosis (41%) was higher when compared to blacks (37%) and whites (29%). Conversely, Hispanic males had the lowest percentage of cases diagnosed at ages 40-49 years (14%) as compared to black males and white males (18% and 22% respectively). Six percent (6%) of black males were teenagers at time of diagnosis compared to 2% of white males and 2% of Hispanic males.

Similar patterns exist among females with HIV disease. More females were diagnosed with HIV disease at ages 30-39 (688 or 32%) than in any other age category. For female cases, age at identification was nearly identical across age groups for black and white females, while Hispanic females were most often identified in the 20–29-year age group (43%).



Table 2. Cumulative <sup>(1)</sup> Adult/Adolescent* HIV Disease Cases By Transmission Route, Race/Ethnicity and Sex at Birth as of December 31, 2023, Kentucky											
	Transmission Category	White, Not Hispanic		Black, Not Hispanic		Hispanic		Other/Unknown		TOTAL	
		No.	%	No.	%	No.	%	No.	%	No.	%
<b>MALE</b>	MMSC	4,425	70	1,684	58	411	67	267	67	6,787	67
	IDU	469	7	335	12	34	6	24	6	862	8
	MMSC/IDU	563	9	179	6	27	4	25	6	794	8
	Heterosexual	223	4	231	8	46	7	30	8	530	5
	Other <sup>(2)</sup>	85	1	14	<1	1	<1	0	0	100	1
	Undetermined <sup>(3)</sup>	519	8	459	16	98	16	52	13	1,128	11
	<b>TOTAL <sup>(4)</sup></b>	<b>6,284</b>	<b>100</b>	<b>2,902</b>	<b>100</b>	<b>617</b>	<b>100</b>	<b>398</b>	<b>100</b>	<b>10,201</b>	<b>100</b>
<b>FEMALE</b>	IDU	321	32	173	19	11	11	18	15	523	25
	Heterosexual	429	43	431	47	54	56	67	55	981	46
	Female Heterosexual	174	18	252	28	26	27	29	24	481	23
	Other <sup>(2)</sup>	12	1	4	<1	0	0	1	1	17	1
	Undetermined <sup>(3)</sup>	56	6	52	6	6	6	6	5	120	6
	<b>TOTAL <sup>(4)</sup></b>	<b>992</b>	<b>100</b>	<b>912</b>	<b>100</b>	<b>97</b>	<b>100</b>	<b>121</b>	<b>100</b>	<b>2,122</b>	<b>100</b>

\*Cases are classified as adult/adolescent if they were 13 years of age or older at time of HIV diagnosis.

(1) Includes HIV disease cases diagnosed from the beginning of the epidemic as of December 31, 2023.

(2) Other includes persons who had a transfusion/transplant, hemophilia/coagulation disorder or pediatric cases diagnosed as adults.

(3) Undetermined refers to persons whose route of exposure to HIV is unknown. This includes persons who are under investigation, dead, lost to investigation, refused interview and persons whose mode of exposure remains undetermined after investigation.

(4) Percentages may not total 100% due to rounding.


See terminology on page 4 for additional definition by transmission category.

Among adult/adolescent males, the majority of cumulative HIV cases reported the primary route of exposure as MMSC (67%), while among adult/adolescent women, most (46%) were exposed through heterosexual contact with a person with HIV or at high risk for HIV infection (e.g., a person who injects drugs). Adult/adolescent black males (12%) reported higher percentages of IDU as the route of HIV transmission in comparison to adult/adolescents white (7%) and Hispanic males (6%). Conversely, a higher percentage of adult/adolescent white males (70%) reported MMSC as the primary route of transmission as compared to 58% of all adult/adolescent black males and 67% of all adult/adolescent Hispanic males.

The most reported risk factor for adult/adolescent female cases in each racial/ethnic group was heterosexual contact (46%). When including female heterosexual contact as a risk category, only 6% of adult/adolescent females have undetermined routes of transmission compared to 11% of adult/adolescent males. Adult/adolescent Hispanic and black males (each 16%) have higher percentages of cases without an identified risk factor than adult/adolescent white males (8%). The existence of large percentages of cases without known routes of transmission poses a barrier to the provision of effective responses to the epidemic within these groups. Risk factor information forms the basis for program planning, service provision and guides resource allocation.








\* Percentages may not total 100% due to rounding.


Figure 1 shows the distribution of cumulative Kentucky HIV cases by age at diagnosis. One-third (33%) of cumulative HIV cases in Kentucky were aged 30-39 years at time of diagnosis. Persons aged 20-29 years also account for almost a third of cumulative cases (32%). Children (aged <13 years at diagnosis) and teenagers (aged 13-19 years) account for the smallest percentages of cases at less than 5% each.



\* Percentages may not total 100% due to rounding.


Figure 2 shows the percentage of HIV cases by age group and sex. Cumulatively, 10,259 male HIV cases have been diagnosed, of which 33% were aged 30-39 years at time of diagnosis. Similarly, females aged 30-39 years at time of diagnosis accounted for the highest percentage of cumulative HIV cases by age group among females (32%). The mean age at diagnosis for both males and females is 35 years.





\* Percentages may not total 100% due to rounding.


Figure 3 shows that 59% of cumulative HIV cases diagnosed in Kentucky are in white populations, 31% are in black populations, and 6% are in Hispanic populations.



\* Percentages may not total 100% due to rounding.

Figure 4 shows the percentages of cumulative HIV cases within each sex group by race/ethnicity. Among males, the majority are white (62%) with black males accounting for 29% of cumulative cases. The distribution among females by racial/ethnic grouping differs from males with both black, as well as white females accounting for almost equal percentage of cases at 43% and 47% respectively.

### Cumulative Adult/Adolescent HIV Diagnoses by Transmission Route, Kentucky



**Table 3. Cumulative Adult/Adolescent HIV Disease Cases by Transmission Route as of December 31, 2023, Kentucky**

Transmission Route	No.	%
MMSC	6,787	55
IDU	1,385	11
MMSC/IDU	794	6
Heterosexual	1,511	12
Female Heterosexual*	481	4
Other†	117	1
Undetermined	1,248	10
<b>Total**</b>	<b>12,323</b>	<b>100</b>

\* Female Heterosexual = A female not reporting drug use but reporting sex with male. See terminology on page 4 for additional definition.

\*\* Percentages may not total 100% due to rounding.

† Other includes persons with transfusion/transplant, hemophilia/coagulation listed as mode of transmission or persons with perinatal exposure who were diagnosed as an adult. See Table 12 for perinatal data.

In Kentucky, 55% of cumulative adult/adolescent HIV cases identified their primary transmission route as MMSC as shown in Figure 5. Twelve percent (12%) of adult/adolescent HIV cases reported heterosexual contact as their primary transmission route, 11% reported IDU and 6% reported both MMSC and IDU. Ten percent (10%) of cumulative adult/adolescent HIV cases were reported without a risk factor identified. Cumulative adult/adolescent HIV case frequencies for each route of exposure are displayed in Table 3.



## Cumulative HIV Diagnoses by Residential Area Development District (ADD) and County at Time of Diagnosis, Kentucky

**Table 4. Cumulative and Living HIV Disease Cases by Residential ADD and County at Time of Diagnosis as of December 31, 2023, Kentucky<sup>(1)</sup>**

ADD/County	Total HIV Disease Cases <sup>(2)</sup>	Total Living with HIV Disease <sup>(3)</sup>	ADD/County	Total HIV Disease Cases <sup>(2)</sup>	Total Living with HIV Disease <sup>(3)</sup>
<b>Barren River</b>	<b>431</b>	<b>271</b>	<b>Buffalo Trace</b>	<b>67</b>	<b>42</b>
Allen	26	13	Bracken, Fleming and Robertson*	18	10
Barren	52	30	Lewis	17	8
Butler	17	14	Mason	32	24
Edmonson	11	8			
Hart and Metcalfe*	20	8			
Logan	33	21			
Monroe	17	9	<b>Cumberland Valley</b>	<b>251</b>	<b>155</b>
Simpson	27	18	Bell	30	21
Warren	228	150	Clay	36	26
			Harlan	25	10
			Jackson	18	12
			Knox	28	19
<b>Big Sandy</b>	<b>102</b>	<b>65</b>	Laurel	55	34
Floyd	29	20	Rockcastle	13	8
Johnson and Magoffin*	19	10	Whitley	46	25
Martin	13	12			
Pike	41	23			
			<b>FIVCO</b>	<b>173</b>	<b>103</b>
			Boyd	99	58
			Carter	24	16
<b>Bluegrass</b>	<b>2,395</b>	<b>1,695</b>	Elliott and Lawrence*	21	10
Anderson	37	25	Greenup	29	19
Bourbon	37	25			
Boyle	39	26			
Clark	64	46	<b>Gateway</b>	<b>123</b>	<b>84</b>
Estill	13	7	Bath	15	11
Fayette	1,633	1,153	Menifee	12	11
Franklin	127	89	Montgomery	32	24
Garrard	15	9	Morgan	35	17
Harrison	14	10	Rowan	29	21
Jessamine	93	73			
Lincoln	17	9	<b>Green River</b>	<b>338</b>	<b>205</b>
Madison	139	107	Daviess	164	92
Mercer	37	19	Hancock and Webster*	21	13
Nicholas	7	6	Henderson	74	44
Powell	12	7	McLean	11	8
Scott	67	51	Ohio	14	9
Woodford	44	33	Union	54	39

(1) One case was missing residential county at time of diagnosis.  
 (2) Total cases with HIV disease regardless of progression to AIDS, both living and deceased.  
 (3) Living cases regardless of current residence.

(Continued on page 12)  
 \* Cases combined due to confidentiality guidelines.

## Cumulative HIV Diagnoses by Residential Area Development District (ADD) and County at Time of Diagnosis, Kentucky

**Table 4 (continued). Cumulative and Living HIV Disease Cases By Residential ADD and County at Time of Diagnosis as of December 31, 2023, Kentucky <sup>(1)</sup>**

ADD/County	Total HIV Disease Cases <sup>(2)</sup>	Total Living with HIV Disease <sup>(3)</sup>	ADD/County	Total HIV Disease Cases <sup>(2)</sup>	Total Living with HIV Disease <sup>(3)</sup>
<b>Kentucky River</b>	<b>100</b>	<b>58</b>	<b>Northern Kentucky</b>	<b>1,068</b>	<b>723</b>
Breathitt, Owsley and Wolfe*	14	6	Boone	190	140
Knott	17	11	Campbell	214	143
Lee and Leslie*	10	5	Carroll	18	13
Letcher	23	9	Gallatin and Owen*	17	14
Perry	36	27	Grant	37	24
			Kenton	580	379
			Pendleton	12	10
<b>KIPDA/North Central</b>	<b>5,982</b>	<b>3,849</b>	<b>Pennyriple</b>	<b>392</b>	<b>226</b>
Bullitt	128	100	Caldwell	23	13
Henry	32	24	Christian	186	126
Jefferson	5,502	3,557	Crittenden and Lyon*	28	8
Oldham	203	86	Hopkins	57	33
Shelby	92	67	Livingston	15	6
Spencer and Trimble*	25	15	Muhlenberg	38	19
			Todd	30	14
			Trigg	15	7
<b>Lake Cumberland</b>	<b>208</b>	<b>145</b>	<b>Purchase</b>	<b>365</b>	<b>218</b>
Adair and Cumberland*	13	7	Ballard and Carlisle*	17	11
Casey	10	6	Calloway	46	28
Clinton	14	11	Fulton	12	9
Green	10	8	Graves	61	34
McCreary	22	20	Hickman	9	7
Pulaski	73	44	Marshall	33	20
Russell	17	11	McCracken	187	109
Taylor	32	27			
Wayne	17	11			
<b>Lincoln Trail</b>	<b>422</b>	<b>293</b>			
Breckinridge	20	9			
Grayson	21	13			
Hardin	247	175			
Larue	9	8			
Marion	23	14			
Meade	33	24			
Nelson	59	43			
Washington	10	7			

(1) One case was missing residential county at time of diagnosis.


(2) Total cases with HIV disease regardless of progression to AIDS, both living and deceased.

(3) Living cases regardless of current residence.

\* Cases combined due to confidentiality guidelines.




**Figure 6. Cumulative HIV Disease Cases Diagnosed by Residential County at Time of Diagnosis as of December 31, 2023, Kentucky\***



\* One case was missing residential county at time of diagnosis.

**Figure 7. HIV Disease Prevalence Rates by Residential County at Time of Diagnosis as of December 31, 2023, Kentucky**




\* Rates not published when cell size is less than 10.





**Figure 8. Cumulative HIV Disease Diagnoses by ADD of Residence at Time of HIV Diagnosis as of December 31, 2023, Kentucky\***

**Cumulative HIV Disease Diagnoses by ADD**




\*One case was missing residential county information at time of diagnosis.

Figure 8 indicates that the highest number of cumulative HIV cases, 5,982 (48%), resided in the KIPDA ADD at the time of diagnosis, which includes the city of Louisville. The Bluegrass ADD, which includes the city of Lexington, had the second highest number of HIV cases diagnosed, 2,395 (19%), followed by the Northern Kentucky ADD, including a portion of the Cincinnati metropolitan area, with 1,068 (9%) of cumulative cases.

**Figure 9. Percentage of Cumulative HIV Disease Diagnoses by ADD of Residence at Time of HIV Diagnosis as of December 31, 2023, Kentucky\***

**Cumulative % HIV Disease Diagnoses by ADD**



\*One case was missing residential county information at time of diagnosis.

Figure 9 shows the percentage of the cumulative (12,418) HIV cases statewide that were diagnosed within each ADD. The percentage of diagnoses by ADD ranged from 1% of total statewide cases residing in each of Buffalo Trace, Gateway, FIVCO, Big Sandy and Kentucky River ADDs to almost half (48%) residing in the KIPDA ADD at time of diagnosis.



## Persons Living with HIV Disease by Demographics, Kentucky

Table 5. Living HIV Disease Diagnoses by Transmission Route, Race/Ethnicity and Sex at Birth as of December 31, 2023, Kentucky <sup>(1)</sup>											
	Transmission Category	White, Not Hispanic		Black, Not Hispanic		Hispanic		Other/Unknown		TOTAL	
		No.	%	No.	%	No.	%	No.	%	No.	%
MALE	MMSC	2,763	71	1,209	64	379	69	248	69	4,599	69
	IDU	262	7	116	6	20	4	19	5	417	6
	MMSC/IDU	358	9	80	4	23	4	19	5	480	7
	Heterosexual	116	3	135	7	40	7	24	7	315	5
	Perinatal	12	<1	20	1	0	0	4	1	36	1
	Other <sup>(2)</sup>	12	<1	2	<1	1	<1	0	0	15	<1
	Undetermined <sup>(3)</sup>	359	9	316	17	83	15	44	12	802	12
	Male Subtotal <sup>(4)</sup>	<b>3,882</b>	<b>100</b>	<b>1,878</b>	<b>100</b>	<b>546</b>	<b>100</b>	<b>358</b>	<b>100</b>	<b>6,664</b>	<b>100</b>
FEMALE	IDU	219	33	71	12	6	7	13	12	309	21
	Heterosexual	281	42	289	48	50	57	55	52	675	46
	Female Heterosexual	131	19	199	33	23	26	28	27	381	26
	Perinatal	8	1	13	2	4	5	2	2	27	2
	Other <sup>(2)</sup>	0	0	1	<1	0	0	1	1	2	<1
	Undetermined <sup>(3)</sup>	33	5	30	5	5	6	6	6	74	5
	Female Subtotal <sup>(4)</sup>	<b>672</b>	<b>100</b>	<b>603</b>	<b>100</b>	<b>88</b>	<b>100</b>	<b>105</b>	<b>100</b>	<b>1,468</b>	<b>100</b>
ALL LIVING	MMSC	2,763	61	1,209	49	379	60	248	54	4,599	57
	IDU	481	11	187	8	26	4	32	7	726	9
	MMSC/IDU	358	8	80	3	23	4	19	4	480	6
	Heterosexual	397	9	424	17	90	14	79	17	990	12
	Female Heterosexual	131	3	199	8	23	4	28	6	381	5
	Perinatal	20	<1	33	1	4	<1	6	1	63	1
	Other <sup>(2)</sup>	12	<1	3	<1	1	<1	1	<1	17	<1
	Undetermined <sup>(3)</sup>	392	9	346	14	88	14	50	11	876	11
	TOTAL <sup>(4)</sup>	<b>4,554</b>	<b>100</b>	<b>2,481</b>	<b>100</b>	<b>634</b>	<b>100</b>	<b>463</b>	<b>100</b>	<b>8,132</b>	<b>100</b>

(1) Includes living HIV disease cases diagnosed from beginning of the epidemic as of December 31, 2023.

(2) Other includes persons who had exposure through hemophilia/coagulation disorder, transfusion/transplant or pediatric cases diagnosed as adults.

(3) Undetermined refers to persons whose route of exposure to HIV is unknown. This includes persons who are under investigation, dead, lost to investigation, refused interview and persons whose route of exposure remains undetermined after investigation.

(4) Percentages may not total 100% due to rounding.

See terminology on page 4 for additional definition by transmission category.

Table 5 shows living HIV cases diagnosed through December 31, 2023, by demographic and behavioral characteristics. There are 8,132 Kentuckians reported to be living with HIV (prevalence rate: 179.7 cases per 100,000). The distribution of behavioral characteristics varied by race/ethnicity and sex, but the majority of Kentucky males living with HIV contracted the disease through MMSC (69%), whereas the majority of Kentucky females contracted HIV through heterosexual contact (46%). An additional 26% of females reported female heterosexual contact which is different than heterosexual contact in that the behavioral risk or serostatus of the male partner is unknown.





## Section II: New HIV Infections Diagnosed among Kentuckians, as of December 31, 2023


As of December 31, 2023, a total of 12,418 cumulative HIV infections among Kentuckians had been reported to the Department for Public Health’s HIV/AIDS Surveillance Program since AIDS reporting started in 1982. Of these infections, 59% have progressed to AIDS. The number of new HIV infections diagnosed since 2013 are presented in Table 6 along with the percentage from each year that have progressed to AIDS. Of the 3,924 HIV infections diagnosed since 2013, 1,204 (31%) had progressed to AIDS as of December 31, 2023.

Year of HIV Diagnosis	TOTAL HIV/AIDS*	Percentage that Progressed to AIDS <sup>†</sup>
	No.	%
2013	356	38%
2014	349	36%
2015	336	32%
2016	341	38%
2017	363	32%
2018	379	27%
2019	333	27%
2020	303	27%
2021	394	24%
2022	405	27%
2023 <sup>†</sup>	365	30%
<b>TOTAL</b>	<b>3,924</b>	<b>31%</b>

\*Total HIV infections regardless of disease progression.

<sup>†</sup>Data reported as of December 31, 2023.

Figure 10 displays annual HIV diagnosis rates among Kentuckians. The annual HIV diagnosis rate has remained fairly steady from 2013 to 2022 with slight fluctuations between 6.7 to 9.0 cases per 100,000 population.



\*Data are current as of December 31, 2023. Data for 2023 and 2024 are considered preliminary due to reporting delays and not included in trend analysis.



## A Comparison of Kentucky to Other States and Washington, DC., Using National Data from the Centers for Disease Control and Prevention (CDC), 2022<sup>(1)</sup>

**Table 7. Estimated\* Annual HIV Disease Diagnosis Rates per 100,000 Population by Residence at Time of Diagnosis, 2022**

Rank	Area of Residence	Rate
1	Washington, DC	31.4
2	Georgia	23.0
3	Florida	19.3
4	Louisiana	18.6
5	Nevada	16.7
6	Texas	16.3
7	Mississippi	15.3
8	Alabama	13.8
9	South Carolina	13.6
10	North Carolina	12.7
11	Delaware	12.6
12	California	12.5
13	Maryland	12.2
13	Tennessee	12.2
15	New Jersey	11.8
16	Arizona	11.7
17	New York	11.3
18	Illinois	10.4
19	Arkansas	10.0
20	Oklahoma	9.8
21	Virginia	9.6
22	Indiana	9.0
<b>22</b>	<b>Kentucky**</b>	<b>9.0</b>
24	Missouri	8.3
25	West Virginia	7.7
26	Colorado	7.4

Rank	Area of Residence	Rate
26	Pennsylvania	7.4
28	Ohio	7.3
29	Massachusetts	6.3
29	Michigan	6.3
29	Rhode Island	6.3
29	Washington	6.3
33	Connecticut	6.1
34	Oregon	5.9
35	Hawaii	5.4
36	Alaska	5.2
37	North Dakota	4.9
37	Wisconsin	4.9
39	Minnesota	4.6
39	Nebraska	4.6
39	Utah	4.6
42	Kansas	4.5
42	South Dakota	4.5
44	New Mexico	4.4
45	Iowa	3.8
46	Maine	3.0
47	Wyoming	2.2
48	Idaho	2.1
49	New Hampshire	2.0
50	Montana	1.0
51	Vermont	0.6

Centers for Disease Control and Prevention. HIV Surveillance Report, 2022; vol.35 <http://www.cdc.gov/hiv-data/nhss/hiv-diagnoses-deaths-prevalence.html/>. Published May 2024. Accessed May 2024.

\*Estimated numbers resulted from statistical adjustment that accounted for reporting delays, but not incomplete reporting.

\*\*Kentucky's rate is estimated by CDC using a different methodology and should not be compared directly to reported data elsewhere in this report.

**Estimated National HIV Diagnosis Rate per 100,000, 2022: 11.3**

In 2022, the annual estimated national HIV diagnosis rate was 11.3 per 100,000 population. The diagnosis rates among the 50 States and Washington, DC ranged from 0.6 per 100,000 population (Vermont) to 31.4 per 100,000 (Washington, DC). Kentucky ranked 22nd with an estimated diagnosis rate of 9.0 per 100,000.



## New HIV Diagnosis: Kentucky vs. the United States, 2022

Characteristics	Number of New Cases	% of New HIV cases <sup>(1)</sup>
<b>SEX</b>		
Male (adult/adolescent)	333	82
Female (adult/adolescent)	72	18
Child (<13 yrs)	0	0
<b>TOTAL</b>	405	100
<b>AGE AT DIAGNOSIS‡</b>		
<13	0	0
13-24	75	19
25-44	231	57
45-64	92	23
65+	7	2
<b>TOTAL</b>	405	100
<b>RACE/ETHNICITY</b>		
White, Not Hispanic	210	52
Black, Not Hispanic	120	30
Hispanic	57	14
Other/Unknown	18	4
<b>TOTAL</b>	405	100
<b>TRANSMISSION ROUTE</b>		
MSM	212	52
IDU	69	17
MSM/IDU	24	6
Heterosexual	27	7
Perinatal	0	0
Other/Undetermined <sup>(2)</sup>	73	18
<b>TOTAL</b>	405	100

Characteristics	Number of New Cases <sup>(6)</sup>	% of New HIV cases <sup>(1)</sup>
<b>SEX</b>		
Male (adult/adolescent)	30,621	81
Female (adult/adolescent)	6,980	19
Child (<13 yrs)	62	<1
<b>TOTAL<sup>†</sup></b>	37,663	100
<b>AGE AT DIAGNOSIS‡</b>		
<13	62	<1
13-24	7,049	19
25-44	22,314	59
45-64	7,366	20
65+	872	2
<b>TOTAL<sup>†</sup></b>	37,663	100
<b>RACE/ETHNICITY</b>		
White, Not Hispanic	9,110	24
Black, Not Hispanic	14,582	39
Hispanic	11,814	31
Other	2,157	6
<b>TOTAL<sup>†</sup></b>	37,663	100
<b>TRANSMISSION ROUTE</b>		
MSM	25,193	67
IDU	2,621	7
MSM/IDU	1,323	4
Heterosexual	8,380	22
Perinatal	118	<1
Other/Undetermined <sup>(2)</sup>	29	<1
<b>TOTAL<sup>†</sup></b>	37,664	100

\*HIV diagnoses regardless of disease progression.

(1) Percentages may not always total 100% due to rounding.

(2) Includes hemophilia, blood transfusion, and risk not reported or not identified.

(3) U.S. cases from CDC. HIV Surveillance Report: Diagnoses of HIV Infection in the United States and Dependent Areas, 2022: 35.

‡ Age at initial HIV diagnosis.

† Totals among subpopulations may be different because values were calculated independently.

Kentucky’s distribution of HIV cases by sex and age at diagnosis (Table 8) closely parallels that of the U.S. (Table 9). The percentage of new HIV cases in Kentuckians that are white, not Hispanic, is much greater than in the U.S. population (52% vs. 24%, respectively). This can be partially attributed to the greater percentage of white, not Hispanic, persons in Kentucky’s general population (84%) as compared to the U.S. population (60%)<sup>1</sup>. U.S. cases have been adjusted for reporting delays and missing risk factors. Kentucky cases have not been adjusted.

<sup>1</sup><https://www.census.gov/quickfacts/fact/table/KY,US/PST045218>



## Adult/Adolescent HIV Diagnoses Regardless of Progression to AIDS†, Kentucky

Characteristics	1982-17		2018		2019		2020		2021		2022		2023 <sup>(2)</sup>		Total	
SEX	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male	8,411	83	310	82	274	83	257	85	321	81	333	82	295	81	10,201	83
Female	1,735	17	69	18	58	17	45	15	73	19	72	18	70	19	2,122	17
<b>TOTAL<sup>(3)</sup></b>	<b>10,146</b>	<b>100</b>	<b>379</b>	<b>100</b>	<b>332</b>	<b>100</b>	<b>302</b>	<b>100</b>	<b>394</b>	<b>100</b>	<b>405</b>	<b>100</b>	<b>365</b>	<b>100</b>	<b>12,323</b>	<b>100</b>
<b>AGE AT DIAGNOSIS*</b>																
13-19	390	4	28	7	23	7	9	3	10	3	11	3	9	2	480	4
20-29	3,168	31	138	36	127	38	109	36	130	33	138	34	110	30	3,920	32
30-39	3,405	34	108	28	81	24	88	29	135	34	115	28	109	30	4,041	33
40-49	2,156	21	54	14	56	17	58	19	64	16	68	17	81	22	2,537	21
50+	1,027	10	51	13	45	14	38	13	55	14	73	18	56	15	1,345	11
<b>TOTAL<sup>(3)</sup></b>	<b>10,146</b>	<b>100</b>	<b>379</b>	<b>100</b>	<b>332</b>	<b>100</b>	<b>302</b>	<b>100</b>	<b>394</b>	<b>100</b>	<b>405</b>	<b>100</b>	<b>365</b>	<b>100</b>	<b>12,323</b>	<b>100</b>
<b>RACE/ETHNICITY</b>																
White, Not Hispanic	6,050	60	214	56	195	59	183	61	223	57	210	52	201	55	7,276	59
Black, Not Hispanic	3,218	32	109	29	88	27	67	22	115	29	120	30	97	27	3,814	31
Hispanic	484	5	24	6	32	10	29	10	31	8	57	14	57	16	714	6
Other/Unknown	394	4	32	8	17	5	23	8	25	6	18	4	10	3	519	4
<b>TOTAL<sup>(3)</sup></b>	<b>10,146</b>	<b>100</b>	<b>379</b>	<b>100</b>	<b>332</b>	<b>100</b>	<b>302</b>	<b>100</b>	<b>394</b>	<b>100</b>	<b>405</b>	<b>100</b>	<b>365</b>	<b>100</b>	<b>12,323</b>	<b>100</b>
<b>TRANSMISSION ROUTE</b>																
MSM	5,686	56	205	54	180	54	153	51	181	46	212	52	170	47	6,787	55
IDU	1,045	10	53	14	50	15	49	16	83	21	69	17	36	10	1,385	11
MSM/IDU	654	6	24	6	32	10	20	7	26	7	24	6	14	4	794	6
Heterosexual	1,349	13	27	7	23	7	26	9	26	7	27	7	33	9	1,511	12
Female Heterosexual	357	4	21	6	18	5	15	5	19	5	22	5	29	8	481	4
Other <sup>(4)</sup>	116	1	0	0	0	0	0	0	0	0	0	0	1	0	117	1
Undetermined <sup>(5)</sup>	939	9	49	13	29	9	39	13	59	15	51	13	82	22	1,248	10
<b>TOTAL<sup>(3)</sup></b>	<b>10,146</b>	<b>100</b>	<b>379</b>	<b>100</b>	<b>332</b>	<b>100</b>	<b>302</b>	<b>100</b>	<b>394</b>	<b>100</b>	<b>405</b>	<b>100</b>	<b>365</b>	<b>100</b>	<b>12,323</b>	<b>100</b>

†HIV disease cases include both persons with HIV alone and those who have progressed to AIDS.

\*Age at time of initial HIV diagnosis.

(1) Cases are classified as Adult/Adolescent if they were 13 years of age or older at time of diagnosis.

(2) Data reported as of December 31, 2023. Data from 2023 are not used in trend analyses due to reporting delays.

(3) Percentages may not total 100% due to rounding.

(4) Other includes persons who had exposure through hemophilia/coagulation disorder, transfusion/transplant or perinatal diagnosed as an adult.

(5) Undetermined refers to persons whose route of exposure to HIV is unknown. This includes persons who are under investigation, deceased, lost to investigation, refused interview and persons whose route of exposure remains undetermined after investigation.

See terminology on page 4 for additional definition by transmission category.

Table 10 shows a breakdown of new adult/adolescent HIV diagnoses by year of diagnosis and demographic characteristics. Cumulative data are presented through December 31, 2023. New diagnoses over the most recent years for which data are complete, 2018-2022, have been predominantly among males, white persons and males reporting sexual contact with other males. New HIV cases over the five-year period (2018-2022) were also highest among 20–29-year-olds in comparison to other age groups. This shows a change in trends as compared to total cases, where highest number of cases are among 30-39 years old group.



### Adult/Adolescent HIV Diagnoses that have Progressed to AIDS†, Kentucky

Table 11. Adult/Adolescent <sup>(1)</sup> HIV Disease Cases with AIDS by Year of Initial HIV Diagnosis, Sex, Age at Diagnosis, Race/Ethnicity, and Transmission Route, Kentucky																
Characteristics	1982-17		2018		2019		2020		2021		2022		2023 <sup>(2)</sup>		Total	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
<b>SEX</b>																
Male	5,563	84	85	83	73	83	65	78	79	82	96	87	89	82	6,050	83
Female	1,099	16	17	17	15	17	18	22	17	18	14	13	19	18	1,199	17
<b>TOTAL<sup>(3)</sup></b>	6,662	100	102	100	88	100	83	100	96	100	110	100	108	100	7,249	100
<b>AGE AT DIAGNOSIS*</b>																
13-19	203	3	5	5	3	3	1	1	0	0	2	2	1	1	215	3
20-29	1,847	28	29	28	23	26	16	19	17	18	25	23	26	24	1,983	27
30-39	2,423	36	26	25	22	25	30	36	29	30	35	32	29	27	2,594	36
40-49	1,479	22	20	20	18	20	16	19	27	28	25	23	26	24	1,611	22
50+	710	11	22	22	22	25	20	24	23	24	23	21	26	24	846	12
<b>TOTAL<sup>(3)</sup></b>	6,662	100	102	100	88	100	83	100	96	100	110	100	108	100	7,249	100
<b>RACE/ETHNICITY</b>																
White, Not Hispanic	4,068	61	62	61	50	57	52	63	56	58	62	56	59	55	4,409	61
Black, Not Hispanic	2,054	31	24	24	19	22	17	20	26	27	29	26	26	24	2,195	30
Hispanic	307	5	7	7	9	10	7	8	13	14	15	14	20	19	378	5
Other/Unknown	233	3	9	9	10	11	7	8	1	1	4	4	3	3	267	4
<b>TOTAL<sup>(3)</sup></b>	6,662	100	102	100	88	100	83	100	96	100	110	100	108	100	7,249	100
<b>TRANSMISSION ROUTE</b>																
MMSC	3,650	55	51	50	41	47	36	43	39	41	58	53	51	47	3,926	54
IDU	804	12	15	15	8	9	17	20	16	17	18	16	9	8	887	12
MMSC/IDU	459	7	4	4	7	8	4	5	6	6	5	5	1	1	486	7
Heterosexual	982	15	3	3	11	13	10	12	7	7	9	8	7	6	1,029	14
Female Heterosexual	168	3	7	7	4	5	4	5	6	6	3	3	10	9	202	3
Other <sup>(4)</sup>	112	2	0	0	0	0	0	0	0	0	0	0	1	1	113	2
Undetermined <sup>(5)</sup>	487	7	22	22	17	19	12	14	22	23	17	15	29	27	606	8
<b>TOTAL<sup>(3)</sup></b>	6,662	100	102	100	88	100	83	100	96	100	110	100	108	100	7,249	100

†HIV disease cases that have progressed to AIDS include only persons reported with an AIDS diagnosis as of December 31, 2023.

\*Age at time of initial HIV diagnosis.

(1) Cases are classified as Adult/Adolescent if they were 13 years of age or older at time of diagnosis.

(2) Data reported as of December 31, 2023. Data from 2023 are not used in trend analyses due to reporting delays.

(3) Percentages may not total 100% due to rounding.

(4) Other includes persons who had exposure through hemophilia/coagulation disorder, transfusion/transplant, or perinatal diagnosed as an adult.

(5) Undetermined refers to persons whose route of exposure to HIV is unknown. This includes persons who are under investigation, deceased, lost to investigation, refused interview, and persons whose route of exposure remains undetermined after investigation.

See terminology on page 4 for additional definition by transmission category.

Table 11 shows a breakdown of adult/adolescent HIV diagnoses that have progressed to AIDS by year of initial HIV diagnosis and demographic characteristics. Newly diagnosed cases that had progressed to AIDS as of December 31, 2023, were predominantly male, white persons and males reporting sexual contact with other males.

**Table 12. Number and Percentage of Cumulative Pediatric<sup>(1)</sup> HIV Disease Cases by Transmission Route and Race/Ethnicity as of December 31, 2023, Kentucky**

Transmission Route	White, Not Hispanic		Black, Not Hispanic		Other <sup>(2)</sup> Unknown		TOTAL	
	No.	%	No.	%	No.	%	No.	%
	Pediatric Hemophilia/Coagulation Disorder	10	27	1	2	0	0	11
Perinatal Exposure, Mother with HIV	24	65	41	85	10	100	75	79
Pediatric Transfusion/Transplant	2	5	0	0	0	0	2	2
Pediatric Risk Not Identified or Reported	1	3	6	13	0	0	7	7
<b>TOTAL<sup>(3)</sup></b>	<b>37</b>	<b>100</b>	<b>48</b>	<b>100</b>	<b>10</b>	<b>100</b>	<b>95</b>	<b>100</b>

(1) Cases are classified as pediatric if they are less than 13 years of age at time of diagnosis.  
 (2) Other includes Hispanic persons and persons of other races.  
 (3) Percentages may not total 100% due to rounding.

**Table 13. Number and Percentage of Cumulative Pediatric<sup>(1)</sup> HIV Disease Cases by Disease Status and Year of Diagnosis, Kentucky**



Disease Status	1982-2017		2018		2019		2020		2021		2022		2023 <sup>(2)</sup>		Total	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
HIV infections <i>without</i> AIDS	43	46	0	0	0	0	1	100	0	0	0	0	0	0	44	46
HIV infections <i>with</i> AIDS	50	54	0	0	1	100	0	0	0	0	0	0	0	0	51	54
<b>Total<sup>(3)</sup></b>	<b>93</b>	<b>100</b>	<b>0</b>	<b>0</b>	<b>1</b>	<b>100</b>	<b>1</b>	<b>100</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>95</b>	<b>100</b>

(1) Cases are classified as pediatric if they are less than 13 years of age at time of diagnosis.  
 (2) Data reported as of December 31, 2023.  
 (3) Percentages may not total 100% due to rounding.

There have been 95 pediatric HIV cases reported to the Kentucky HIV/AIDS Surveillance Program (Table 12 and Table 13) since reporting began in 1982. The majority of reported pediatric cases (79%) were due to perinatal transmission through an HIV-infected mother, 11 cases were reported with a primary exposure route of pediatric hemophilia or coagulation disorders, and two cases were due to pediatric transfusion or transplant (Table 12). Since 1991, there have been no pediatric HIV cases with hemophilia or coagulation disorders reported as the route of exposure. The two pediatric cases reported with pediatric transfusion or transplant as the risk factor were diagnosed in 1987 or earlier. Eighty-five percent (85%) of the 48 pediatric HIV cases among black populations were due to perinatal exposure as compared to 65% of the 37 pediatric HIV cases among white populations. The majority (55%) of the 75 cumulative perinatal exposures from a mother with HIV were in black mothers.

Table 13 shows disease progression to AIDS as of December 31, 2023. Ninety-two of the cumulative 95 pediatric cases (97%) in Kentucky were diagnosed prior to 2018. One or no new pediatric HIV case have been reported during each of the most recent five years.

## New HIV Disease Cases by Race/Ethnicity, Kentucky



\*\* Percentages may not total 100% due to rounding.

Figure 11 shows the race/ethnicity percentage distribution for newly diagnosed HIV cases among Kentuckians in 2022, the latest year data are considered complete. The majority of cases diagnosed in 2022 were white (52%), followed by black cases (30%).

Figure 12 shows the percentage race/ethnicity distribution of Kentucky’s population based on the 2022 population estimates. The majority of Kentuckians are white, not Hispanic. Persons who identify with multiple races were grouped under the “other” category.

HIV racial disparities are highlighted by these two graphs, showing higher percentages of new cases among black and Hispanic populations in relation to their representation in the general population. Black populations accounted for 30% of new HIV cases diagnosed in 2022 yet comprised just 8% of Kentucky's population in 2022. Similarly, Hispanic populations accounted for 14% of newly diagnosed HIV cases in 2022 yet comprised only 4% of Kentucky’s population in that same year. Rates of new diagnoses by race/ethnicity and sex are presented in Table 14.

**Table 14. Number and Rate of New HIV Diagnoses by Race/Ethnicity and Sex, Kentucky, 2022**

Race/Ethnicity	Male		Female		Total No of Cases	Total Rate
	No of Cases	Rate*	No of Cases	Rate*		
Hispanic	50	42.8	7	†	57	26.0
Black, not Hispanic	98	51.9	22	11.6	120	31.8
White, not Hispanic	170	9.2	40	2.1	210	5.6
Other	15	16.6	3	†	18	9.8
<b>Total**</b>	333	14.9	72	3.2	405	9.0



\*Rate per 100,000 based on census data estimates for racial and gender distribution for Kentucky in 2022.

†Rates are not published when cell size is less than 10.

\*\* Percentages may not total 100% due to rounding.



**New HIV Disease Cases by Age at Diagnosis, Kentucky**



\*\* Percentages may not total 100% due to rounding.

Figure 13 shows the percentage age distribution of newly diagnosed HIV cases among Kentuckians in 2022 at time of HIV diagnosis. The highest percentage of new diagnoses was reported among Kentuckians aged 20-29 years (34%). Kentuckians aged 30-39 and 40-49 years accounted for 28% and 17% of new cases, respectively. Kentuckians aged 50+ years accounted for 18% of new cases diagnosed in 2022.

Figure 14 shows the percentage distribution of Kentucky’s population based on 2022 estimates, which can be directly compared to the percentages in each age group that were newly diagnosed in 2022. HIV-related disparities by age are highlighted by these two graphs. Higher percentages of new diagnoses occurred among persons in age groups 20-29, 30-39, and 40-49 years in comparison to the proportion of these groups in the general population.

**Table 15. Number and Rate of New HIV Diagnoses by Age at Diagnosis and Race/Ethnicity§, Kentucky, 2022**

Age at Diagnosis	Black not Hispanic		White not Hispanic	
	No. of Cases	Rate*	No. of Cases	Rate*
20-29	49	79.6	52	10.9
30-39	35	64.7	60	12.7
40-49	13	28.4	46	9.9
50+	19	17.1	49	3.3

§Rates among pediatric cases (<13 years), teens and Hispanics by age at diagnosis not published due to small numbers.

\*Rate per 100,000 based on census data estimates for racial and age distribution for Kentucky in 2022.

Rates of new diagnoses in 2022 (Table 15) were higher among black populations across all age groups in comparison to white populations. These relative rates were highest among 20-year-olds at the time of diagnosis. However, the rates among black populations in all age groups were at least about three times higher than the rates among their white counterparts of the same age group. Rates among Hispanic populations are not presented due to small numbers.



**Table 16. HIV Disease Cases and Diagnosis Rates by Year of HIV Diagnosis and ADD of Residence at Time of HIV Diagnosis, 1982-2023<sup>(2)</sup>, Kentucky**

AREA DEVELOPMENT DISTRICT	CASES & RATES <sup>(1)</sup>	1982-2017*	2018	2019	2020	2021	2022	2023 <sup>(2)</sup>	TOTAL CASES <sup>(3)</sup>	% of Total
1. Barren River	Cases	360	14	13	9	12	15	8	431	3%
	Rate per 100,000		4.6	4.2		3.8	4.7			
2. Big Sandy	Cases	80	6	4	3	3	5	1	102	1%
	Rate per 100,000									
3. Bluegrass	Cases	1,997	78	47	50	74	79	70	2,395	19%
	Rate per 100,000		9.4	5.6	6.0	8.9	9.4			
4. Buffalo Trace	Cases	56	4	1	2	0	0	4	67	1%
	Rate per 100,000									
5. Cumberland Valley	Cases	204	6	6	5	11	11	8	251	2%
	Rate per 100,000					4.8	4.8			
6. FIVCO	Cases	140	4	8	3	4	4	10	173	1%
	Rate per 100,000									
7. Gateway	Cases	104	5	5	0	6	3	0	123	1%
	Rate per 100,000									
8. Green River	Cases	291	7	8	8	7	9	8	338	3%
	Rate per 100,000									
9. KIPDA	Cases	4,928	169	154	150	201	200	180	5,982	48%
	Rate per 100,000		16.8	15.3	14.6	19.7	19.6			
10. Kentucky River	Cases	86	1	3	2	4	3	1	100	1%
	Rate per 100,000									
11. Lake Cumberland	Cases	177	3	5	3	5	6	9	208	2%
	Rate per 100,000									
12. Lincoln Trail	Cases	326	23	13	14	16	14	16	422	3%
	Rate per 100,000		8.3	4.7	5.0	5.6	4.9			
13. Northern KY	Cases	847	49	42	38	27	29	36	1,068	9%
	Rate per 100,000		10.6	9.0	8.1	5.7	6.1			
14. Pennyriple	Cases	327	4	13	8	15	18	7	392	3%
	Rate per 100,000			6.1		7.0	8.4			
15. Purchase	Cases	315	6	11	8	9	9	7	365	3%
	Rate per 100,000			5.6						
<b>TOTAL CASES <sup>(3)</sup></b>		<b>10,238</b>	<b>379</b>	<b>333</b>	<b>303</b>	<b>394</b>	<b>405</b>	<b>365</b>	<b>12,417</b>	<b>100%</b>

(1) Rates are only listed for years of diagnosis 2018-2022. Data for 2023 are provisional due to reporting delays and are subject to change. Due to the small numbers of HIV cases reported in some ADDs, please interpret the corresponding rates with caution. Rates are not published when cell size is less than 10.

(2) Data reported as of December 31, 2023. Rates are not published for 2023 because data are not complete.

(3) Total HIV disease cases both living and deceased, regardless of progression to AIDS; Total HIV cases reported are 12,418— 1 HIV case had unknown residential information.


\*Rates are not published due to multi-year aggregation of data.



Table 16 shows the HIV disease cases and diagnosis rates by year of HIV diagnosis and ADD of residence at time of HIV diagnosis. The majority of the cases can be accounted for by the three urban ADDs, i.e. KIPDA, Bluegrass, and Northern Kentucky ADDs. The rates are higher in general for KIPDA ADD followed by Bluegrass ADD, which includes the cities of Louisville and Lexington respectively. The KIPDA ADD showed a distinct increase in 2021 and 2022, which on investigation was found to be mainly associated with people who inject drugs.

### Trends in HIV Disease Diagnosis Rates by Sex, 2018-2022, Kentucky

The trends in HIV diagnosis rates among Kentuckians by sex assigned at birth from 2018 to 2022 are presented in figure 15.




\*Data for 2023 and 2024 are not included in trend analyses since they are considered provisional due to reporting delays.

Males represent the majority (83%) of cumulative HIV cases diagnosed among Kentuckians. The yearly diagnosis rates among males varied from 11.5 to 14.9 cases per 100,000 population over the five-year period shown. From 2018 to 2022, the HIV diagnosis rates among males fluctuated between 4.5 to 5.8 times higher than the rate for females (Figure 15).

The female HIV diagnosis rates have remained fairly stable over the most recent five years, between 2.0 to 3.2 cases per 100,000 females. The highest HIV diagnosis rate among females within the most recent five years was in 2021 and 2022 at 3.2 newly diagnosed cases per 100,000 females.

### Trends in HIV Disease Diagnosis Rates by Age at HIV Diagnosis, 2018-2022, Kentucky




\* Due to the small numbers of HIV cases reported, rates are not presented for age groups <13 and 13-19 years old.  
 \*\*Data for 2023 and 2024 are not included in trend analyses since they are considered provisional due to reporting delays

Figure 16 shows HIV diagnosis rates by age category over the most recent five years (2018-2022) with complete data. The diagnosis rates among Kentuckians in the 20-29- and 30-39-year age groups reveal a downward trend from 2018 to 2019, while rates in the other year age groups stayed almost stable during this time. Between 2019 and 2020, the rate decreased among the 20–29-year age group. From 2020 to 2021 the rate increased among 20-29- and 30-39-years old categories. From 2021 to 2022 the rate of diagnosis showed a decrease among 20-29- and 30–39-year-old age groups. The yearly diagnosis rates among those 40-49 and 50 years and over remained almost stable over the five-year period shown.

HIV Diagnosis Year	Mean Age	Age Range
2018	34.1	15-84
2019	34.3	0-73
2020	35	0-77
2021	35.7	15-79
2022	36.2	17-70


Table 17 shows the mean ages and actual age ranges at time of HIV diagnosis from 2018-2022. The mean ages of Kentuckians at time of HIV diagnosis in the five-year period ranged between 34.1-36.2 years (age range 0-84 years).

### Trends in HIV Disease Diagnosis Rates by Race/Ethnicity, 2018-2022, Kentucky



\*Data for 2023 and 2024 are not included in trend analyses since they are considered provisional due to reporting delays.

Figure 17 shows that between 2018 and 2022, the HIV diagnosis rates for black populations fluctuated between 3.7 to 5.7 times higher than white populations. The diagnosis rates for Hispanic populations were between 2.4 to 4.6 times higher than whites over the same five-year period. The trends among white populations have remained steady. The rates for black populations decreased between 2018 and 2020, then increased abruptly between 2020 and 2021, with another slight increase in 2022. The rates for Hispanic populations increased between 2018 and 2019, decreased in 2019, then slightly increased to 14.5 in 2021 with a major increase to 26.0 cases per 100,000 in 2022.




\*Data for 2023 and 2024 are not included in trend analyses since they are considered provisional due to reporting delays.

<sup>a</sup>Rates for Hispanic cases by sex are not presented due to the small number of cases reported.


Figure 18 presents diagnosis rates from 2018 through 2022 for black and white cases by sex. Black males and black females had consistently higher rates of new diagnoses in comparison to their white counterparts. The HIV diagnosis rates among black males fluctuated between 3.6 to 5.6 times higher than that of white males. The rates among black females were 4.2 to 6.6 times higher than those of white females over the five-year period.

### Trends in HIV Disease Diagnosis Rates by Route of Transmission and Sex, 2018-2022, Kentucky



\*Data for 2023 and 2024 are not included in trend analyses since they are considered provisional due to reporting delays.  
 §Female Heterosexual Contact is a female not reporting drug use but reporting sex with male with unknown HIV status or risk.  
 See terminology on page 4.

Figure 19 shows Kentucky’s adult/adolescent female HIV cases by transmission route and year of diagnosis. The largest number of new female cases reported IDU as their primary route of transmission followed by FHC over the five-year period. The number of new female cases reporting IDU decreased from 2018 to 2020 with an abrupt increase in 2021 due to a Jefferson County cluster and slight decrease in 2022. IDU as route of transmission accounted for largest number of cases diagnosed among females from 2018 to 2022, except for 2020.



\*Data for 2023 and 2024 are not included in trend analyses since they are considered provisional due to reporting delays.

In Figure 20, which depicts trends for adult/adolescent males by transmission route, the largest number of cases diagnosed each year from 2018 to 2022 reported MMSC as their primary risk factor. The second largest number of cases were those with an undetermined risk. The number of males reporting IDU as a risk factor constantly increased between 2018 (30 cases) and 2021 (45 cases) and slightly decreased in 2022 (39 cases).

## Section III: HIV Infections Diagnosed Concurrently with AIDS among Kentuckians as of December 31, 2023

During the most recent 10-year period for which data are available (January 1, 2014, to December 31, 2023), a total of 3,568 HIV disease cases were diagnosed among Kentuckians. Of these, 1,070 (30%) had progressed to AIDS as of December 31, 2023.

<b>Time to AIDS Diagnosis (Days)</b>	<b>No.</b>	<b>%</b>
≤30 Days †	770	72.0
31-60 Days	62	5.8
61-90 Days	25	2.3
91-365 Days	77	7.2
>365 Days	136	12.7
Total	1,070	100

†Cases diagnosed with AIDS within 30 days of initial HIV diagnosis are considered concurrent diagnoses.

Note: 2,498 HIV-only cases diagnosed in the same timeframe are not included in the table as they had not progressed to AIDS as of December 31, 2023.

During the most recent 10-year period, 770 (21.6%) of the 3,568 newly diagnosed HIV cases were diagnosed with AIDS within 30 days of the initial HIV diagnosis - also known as a concurrent diagnosis.

The distribution of progression to AIDS (in days) for the 1,070 AIDS cases is shown in Table 18. About 72% of the 1,070 AIDS cases diagnosed in the most recent 10 years were diagnosed with AIDS within 30 days of the initial HIV diagnosis.

According to the CDC\* late testers are those who have an AIDS diagnosis within one year of initial HIV diagnosis. During the presented time period, 934 (26.2%) of the 3,568 Kentuckians diagnosed with HIV disease were late testers.

\*CDC. Late versus early testing of HIV—16 sites, United States, 2000-2003. MMWR 2003; 52(25): 581-586.



## Concurrent Diagnoses by Selected Characteristics, 2014-2023\*, Kentucky

**Table 19. HIV Infections Diagnosed in the Most Recent 10 Year Period (January 1, 2014-December 31, 2023) that were Diagnosed Concurrently with AIDS (within 30 Days of HIV Diagnosis) and those without a Concurrent Diagnosis\*\* by Sex, Age at Diagnosis, Race/Ethnicity and Transmission Category, Kentucky**

Characteristics	HIV With Concurrent AIDS Diagnosis*		HIV Without Concurrent AIDS Diagnosis**		Total HIV Disease Diagnoses***	
	No.	% <sup>(1)</sup>	No.	% <sup>(1)</sup>	No.	% <sup>(1)</sup>
<b>SEX</b>						
Male	646	84	2,301	82	2,947	83
Female	124	16	497	18	621	17
<b>AGE AT DIAGNOSIS</b>						
<13	1	0	11	0	12	0
13-19	10	1	134	5	144	4
20-29	154	20	1,105	39	1,259	35
30-39	218	28	776	28	994	28
40-49	190	25	447	16	637	18
50+	197	26	325	12	522	15
<b>RACE/ETHNICITY- Female</b>						
White, Not Hispanic	58	47	279	56	337	54
Black, Not Hispanic	49	40	175	35	224	36
Hispanic	11	9	16	3	27	4
Other/Unknown	6	5	27	5	33	5
<b>RACE/ETHNICITY- Male</b>						
White, Not Hispanic	375	58	1,278	56	1,653	56
Black, Not Hispanic	150	23	660	29	810	27
Hispanic	84	13	230	10	314	11
Other/Unknown	37	6	133	6	170	6
<b>TRANSMISSION CATEGORY</b>						
MMSC	374	49	1,531	55	1,905	53
IDU	66	9	373	13	439	12
MMSC/IDU	28	4	210	8	238	7
Heterosexual	66	9	191	7	257	7
Female Heterosexual	64	8	165	6	229	6
Perinatal	1	0	11	0	12	0
Other <sup>(2)</sup>	2	<1	0	<1	2	<1
Undetermined <sup>(3)</sup>	169	22	317	11	486	14
<b>TOTAL</b>	<b>770</b>	<b>100</b>	<b>2,798</b>	<b>100</b>	<b>3,568</b>	<b>100</b>

\*Concurrent diagnosis is defined as being diagnosed with both HIV and AIDS within a 30-day period.

\*\*Without AIDS diagnosis 30 days after initial HIV diagnosis. Includes both HIV (non-AIDS) cases and those with an AIDS diagnosis more than 30 days after initial HIV diagnosis.

\*\*\*Total diagnoses January 1, 2014, through December 31, 2023, with HIV, regardless of AIDS diagnosis status.

(1) Percentages may not total to 100% due to rounding.

(2) Other includes persons who had exposure through hemophilia, transfusion/transplant or perinatal diagnosed as an adult.

(3) Undetermined refers to persons whose route of exposure to HIV is unknown. This includes persons who are under investigation, deceased, lost to investigation, refused interview, and persons whose route of exposure remains undetermined after investigation. See terminology on page 4 for additional definition by transmission category.



## Concurrent Diagnoses by Selected Characteristics, 2013-2022, Kentucky (Narrative)

Table 19 (page 30) examines the distribution of HIV cases among Kentuckians diagnosed between January 1, 2014, and December 31, 2023 by sex at birth, age at diagnosis, race/ethnicity and transmission route. Data are presented for cases diagnosed concurrently with AIDS (diagnosed with AIDS within a 30-day period after an initial HIV diagnosis), cases without a concurrent HIV/AIDS diagnosis (anyone who did not have an AIDS diagnosis within 30 days of the initial HIV diagnosis, whether they developed AIDS or not) and for all cases diagnosed with HIV (regardless of AIDS diagnosis status) within the 10-year period.

Of the 3,568 Kentuckians diagnosed with HIV disease during the 10-year period, about one-fifth (770 or 21.6%) were diagnosed with HIV and AIDS concurrently (within 30 days).

During the 10-year period presented, males consistently represent the highest number of diagnosed cases of HIV, with (84%) and without (82%) a concurrent AIDS diagnosis.

The distribution of with and without concurrent diagnosis by age at diagnosis differs between the two groups, with the highest percentages of concurrent cases aged 30-39 (28%), followed by 50 plus (26%) and 40-49 years (25%), while the highest percentages among non-concurrently diagnosed cases were aged 20-29 years (39%), followed by 30-39 years (28%) and 40-49 years at 16%.

The racial/ethnic distribution of cases diagnosed concurrently with AIDS differs by sex. Among females, the highest percentage of concurrent diagnoses were among white females (47%), followed by black females (40%) and Hispanic females at 9%. However, among males, the majority of concurrent diagnoses were among white males (58%). Twenty-three percent (23%) of concurrently diagnosed cases in males were among black males and 13% were among Hispanic males. The percentages of concurrent diagnoses among Hispanic males and Hispanic females are much lower compared to white and black Kentuckians. Caution should be taken when interpreting the data for the other and unknown race/ethnicity categories as the numbers of cases are small.

Data by route of transmission show HIV cases diagnosed concurrently with AIDS have a similar distribution to those without a concurrent diagnosis, with the majority of cases among those with a concurrent diagnosis reporting MMSC as the mode of transmission (49%), followed by both persons reporting heterosexual exposure and IDU at 9%, and female presumed heterosexual at 8%. Twenty-two percent (22%) of concurrently diagnosed HIV and AIDS cases have an undetermined transmission route, which creates challenges for prevention initiatives aimed at increasing early testing and engagement in care.





## HIV Diagnoses by ADD, January 1, 2014-December 31, 2023

Figure 21. Number of HIV Disease Diagnoses by ADD of Residence at Time of HIV Diagnosis, for Most Recent 10 years, January 1, 2014 – December 31, 2023, Kentucky





Figure 21 displays the total number of HIV infections (3,568) diagnosed between January 1, 2014, and December 31, 2023, by ADD of residence at time of HIV diagnosis. Data represent the total number of HIV cases in each ADD, regardless of disease progression status. The highest number of cases (1,672 or 47%) diagnosed during this time period were among residents of the KIPDA ADD, which includes the city of Louisville. The second highest number of cases (669 or 19%) resided in the Bluegrass ADD, which includes the city of Lexington.

Figure 22. Percentage of All HIV Disease Diagnoses within each ADD of Residence at Time of Diagnosis, who have a Concurrent Diagnosis of AIDS, for the Most Recent 10 Years, January 1, 2014 – December 31, 2023, Kentucky



Note: The percentages presented in Figure 22 represent the proportion of concurrent diagnoses out of the total for each individual ADD. Totals for each ADD are presented in Figure 21.

Figure 22 shows the percentage of total HIV cases within each ADD that were concurrently diagnosed with AIDS (within 30 days of initial HIV diagnosis), between January 1, 2014, and December 31, 2023. The percentage of concurrent HIV and AIDS diagnoses within each ADD ranged from 6% to 34%. Big Sandy and Purchase ADDs (34%) had the highest proportion of concurrent HIV and AIDS cases, followed by Cumberland Valley ADD (33%).



## HIV Diagnoses by Care Coordinator Region, January 1, 2014-December 31, 2023

Figure 23. Number of New HIV Disease Diagnoses within each Care Coordinator Region of Residence at Time of Diagnosis, for the Most Recent 10 Years, January 1, 2014—December 31, 2023, Kentucky





Figure 23 shows the total number of new HIV infections (regardless of disease progression status) diagnosed between January 1, 2014, and December 31, 2023, by Care Coordinator Region based on place of residence at time of HIV diagnosis. The highest number of cases (1,672 or 47%) diagnosed in this period occurred among residents of the Louisville Region. The second highest number of diagnoses (771 or 22%) occurred in residents of the Lexington Region.

Figure 24. Percentage of New HIV Cases with Concurrent Diagnosis within each Care Coordinator Region of Residence at Time of Diagnosis, for the Most Recent 10 Years, January 1, 2014—December 31, 2023, Kentucky



Note: The percentages presented in Figure 24 represent the proportion of concurrent diagnoses out of the total for each individual region. Totals for each region are presented in Figure 23. Owsley County is currently being served by both the Lake Cumberland and KY River District Health Departments (HD). In Figures 23 & 24 Owsley County is included only in the KY River District HD, and Graves and Todd Counties are included in Purchase District HD.

Figure 24 shows the percentage of total HIV cases within each Care Coordinator Region that were concurrently diagnosed with AIDS (within 30 days of an initial HIV diagnosis) between January 1, 2014, and December 31, 2023. In all regions, approximately one-fifth or more of cases diagnosed within each jurisdiction were concurrent diagnoses with the highest proportions of concurrent HIV and AIDS cases residing in the Kentucky River Region and Lake Cumberland Region (29% each).



## HIV Care Coordinator Regions, Kentucky

Map for Counties Covered	Region Name and Address	Counties Covered			
	<b>Purchase Region:</b> LivWell Community Health Services 1903 Broadway Street Paducah, KY 42001 <b>(270) 444-8183, (877) 444-8183</b> <b>Fax: (270) 444-8147</b>	Ballard Caldwell Calloway Carlisle Christian	Crittenden Fulton Graves* Hickman	Hopkins Livingston Lyon Marshall	McCracken Muhlenberg Todd** Trigg
	<b>Barren Region:</b> Matthew 25 452 Old Corydon Road Henderson, KY 42420 <b>(270) 826-0200, (866) 607-6590</b> <b>Fax: (270) 826-0212</b>	Allen Barren Breckinridge Butler Daviess Edmonson Grayson	Hancock Hardin Hart Henderson Larue Logan Marion	McLean Meade Metcalfe Monroe Nelson Ohio Simpson	Union Warren Washington Webster
	<b>Louisville Region:</b> ULSD KCCP 1212 S. 4 <sup>th</sup> Street Suite 101 Louisville, KY 40203 <b>(502) 852-2008</b> <b>Fax: (502) 852-2510</b>	Bullitt Henry	Jefferson Oldham	Shelby Spencer	Trimble
	<b>Northern Kentucky Region:</b> Northern KY Distt HD 8001 Veterans Memorial Drive Florence, KY 41042 <b>(859) 341-4264</b> <b>Fax: (859) 578-3689</b>	Boone Campbell	Carroll Gallatin	Grant Kenton	Owen Pendleton
	<b>Lexington Region:</b> UK Bluegrass Care Clinic 3101 Beaumont Ctr Circle Suite 300 Lexington, KY 40513 <b>(859) 323-5544, (866) 761-0206</b> <b>Fax: (859) 257-3477</b>	Anderson Bath Bourbon Boyd Boyle Bracken Carter Clark	Elliott Estill Fayette Fleming Franklin Garrard Greenup Harrison	Jessamine Lawrence Lewis Lincoln Madison Mason Menifee Mercer	Montgomery Morgan Nicholas Powell Robertson Rowan Scott Woodford
	<b>Lake Cumberland Region:</b> Lake Cumberland Distt HD 500 Bourne Avenue Somerset, KY 42501 <b>(606) 678-4761, (800) 928-4416</b> <b>Fax: (606) 678-2708</b>	Adair Bell Breathitt Casey Clay Clinton	Cumberland Floyd Green Harlan Jackson Johnson	Knox Laurel Magoffin Martin McCreary Pike	Pulaski Rockcastle Russell Taylor Wayne Whitley
	<b>Kentucky River Region:</b> Kentucky River Distt HD 441 Gorman Hollow Road Hazard, KY 41701 <b>(606) 439-2361</b> <b>Fax: (606) 439-0870</b>	Knott Lee	Leslie Letcher	Owsley Perry	Wolfe
	<b>Graves County HD</b> 416 Central Ave Mayfield, KY 42066 <b>(270) 247-3553</b>	Graves			
	<b>Todd County HD</b> 205 Public Square Elkton, KY 42220 <b>(270) 265-2362</b>	Todd			
	<b>Mountain Comprehensive Care</b> PO Box Whitesburg, KY 41858 <b>606-633-4823</b>	Bell Clay Floyd Harlan Jackson Johnson	Knott Knox Leslie Letcher Magoffin Owsley	Perry Pike	







## State Sponsored HIV Counseling and Testing Sites, Kentucky

County / Agency	City	Phone	County / Agency	City	Phone
Adair County Health Department	Columbia	(270) 384-2286	Jessamine County Health Department	Nicholasville	(859) 885-4149
Allen County Health Department	Scottsville	(270) 237-4423	Johnson County Health Department	Paintsville	(606) 789-2590
Anderson County Health Department	Lawrenceburg	(502) 839-4551	Kenton County Health Department	Covington	(859) 431-3345
Ballard County Health Department	La Center	(270) 665-5432	Knott County Health Department	Hindman	(606) 785-3144
Barren County Health Department	Glasgow	(270) 651-8321	Knox County Health Department	Barbourville	(606) 546-3486
Bath County Health Department	Owingsville	(606) 674-2731	Larue County Health Department	Hodgenville	(270) 358-3844
Bell County Health Department	Pineville	(606) 248-2862	Laurel County Health Department	London	(606) 864-5187
Boone County Health Department	Florence	(859) 363-2060	Lawrence County Health Department	Louisia	(606) 638-4389
Bourbon County Health Department	Paris	(859) 987-1915	Lee County Health Department	Beattyville	(606) 464-2492
Boyd County Health Department	Ashland	(606) 324-7181	Leslie County Health Department	Hyden	(606) 672-2393
Boyle County Health Department	Danville	(859) 236-2053	Letcher County Health Department	Whitesburg	(606) 633-2945
Bracken County Health Department	Brooksville	(606) 735-2157	Lyon County Health Department	Vanceburg	(606) 796-2632
Breathitt County Health Department	Jackson	(606) 666-7755	Lincoln County Health Department	Stanford	(606) 365-3106
Breckinridge County Health Department	Hardinsburg	(270) 756-5121	Livingston County Health Department	Smithland	(270) 928-2193
Bullitt County Health Department	Shepherdsville	(502) 543-2415	Logan County Health Department	Russellville	(270) 726-8341
Butler County Health Department	Morgantown	(270) 526-3221	Lyon County Health Department	Eddyville	(270) 692-9363
Caldwell County Health Department	Princeton	(270) 365-6571	Madison County Health Department	Richmond	(859) 626-4241
Calloway County Health Department	Murray	(270) 753-3381	Madison County Health Department - Berea	Berea	(859) 986-1192
Campbell County Health Department	Newport	(859) 431-1704	Magoffin County Health Department	Salyersville	(606) 349-6212
Carlisle County Health Department	Bardwell	(270) 628-5431	Marion County Health Department	Lebanon	(270) 692-3393
Carroll County Health Department	Carrollton	(502) 732-6641	Marshall County Health Department	Benton	(270) 527-1496
Carter County Health Department	Grayson	(606) 474-5109	Martin County Health Department	Inez	(606) 298-7752
Casey County Health Department	Liberty	(606) 787-6911	Mason County Health Department	Maysville	(606) 564-9447
Christian County Health Department	Hopkinsville	(270) 887-4160	(McCracken Co.) Heartland Cares Clinic	Paducah	(270) 444-8183
Clark County Health Department	Winchester	(859) 744-4482	McCracken County Health Department	Paducah	(270) 444-9631
Clay County Health Department	Manchester	(606) 598-2425	McCreary County Health Department	Whitley City	(606) 376-2412
Clinton County Health Department	Albany	(606) 387-5711	McLean County Health Department	Calhoun	(270) 273-3062
Crittenden County Health Department	Marion	(270) 965-5215	Meade County Health Department	Brandenburg	(270) 422-3988
Cumberland County Health Department	Burkesville	(270) 864-2206	Menifee County Health Department	Frenchburg	(606) 768-2151
Daviess County Health Department	Owensboro	(270) 686-7744	Mercer County Health Department	Harrodsburg	(859) 734-4522
Edmonson County Health Department	Brownsville	(270) 597-2194	Metcalfe County Health Department	Edmonton	(270) 432-3214
Elliott County Health Department	Sandy Hook	(606) 738-5205	Monroe County Health Department	Tompkinsville	(270) 487-6782
Estill County Health Department	Irvine	(606) 723-5181	Montgomery County Health Department	Mount Sterling	(859) 498-3808
(Fayette Co.) AVOL (AIDS Volunteers, Inc.)	Lexington	(859) 225-3000	Morgan County Health Department	West Liberty	(606) 743-3744
(Fayette Co.) Bluegrass Community Health Center	Lexington	(859) 259-2635	Muhlenberg County Health Department	Central City	(270) 754-3200
(Fayette Co.) Lex-Fayette Health Department	Lexington	(859) 252-2371	Nelson County Health Department	Bardstown	(502) 348-3222
(Fayette Co.) Moveable Feast Lexington	Lexington	(859) 252-2867	Nicholas County Health Department	Carlisle	(859) 289-2188
Fleming County Health Department	Flemingsburg	(606) 845-6511	Ohio County Health Department	Hartford	(270) 298-3663
Floyd County Health Department	Prestonsburg	(606) 886-2788	Oldham County Health Department	LaGrange	(502) 222-3516
Franklin County Health Department	Frankfort	(502) 564-7647	Owen County Health Department	Owenton	(502) 484-5736
Fulton County Health Department	Fulton	(270) 472-1982	Owsley County Health Department	Booneville	(606) 593-5181
Fulton County Health Department – Hickman	Hickman	(270) 236-2825	Pendleton County Health Department	Falmouth	(859) 654-6985
Gallatin County Health Department	Warsaw	(859) 567-2844	Perry County Health Department	Hazard	(606) 436-2196
Garrard County Health Department	Lancaster	(859) 792-2153	Pike County Health Department	Pikeville	(606) 437-5500
Grant County Health Department	Williamstown	(859) 824-5074	Powell County Health Department	Stanton	(606) 663-4360
Graves County Health Department	Mayfield	(270) 247-3553	Pulaski County Health Department	Somerset	(606) 679-4416
Grayson County Health Department	Leitchfield	(270) 259-3141	Robertson County Health Department	Mount Olivet	(606) 724-5222
Green County Health Department	Greensburg	(270) 932-4341	Rockcastle County Health Department	Mt. Vernon	(606) 256-2242
Greenup County Health Department	Greenup	(606) 473-9838	Rowan County Health Department	Morehead	(606) 784-8954
Hancock County Health Department	Hawesville	(270) 927-8803	Russell County Health Department	Jamestown	(270) 343-2181
Hardin County Health Department	Elizabethtown	(270) 765-6196	Scott County Health Department	Georgetown	(502) 863-3971
Harlan County Health Department	Harlan	(606) 573-4820	Shelby County Health Department	Shelbyville	(502) 633-1231
Harrison County Health Department	Cynthiana	(859) 234-2842	Simpson County Health Department	Franklin	(270) 586-8261
Hart County Health Department	Munfordville	(270) 524-2511	Spencer County Health Department	Taylorsville	(502) 477-8146
(Henderson Co.) Matthew 25 AIDS Services	Henderson	(270) 826-0200	Taylor County Health Department	Campbellsville	(270) 465-4191
Henderson County Health Department	Henderson	(270) 826-3951	Todd County Health Department	Elkton	(270) 265-2362
Henry County Health Department	New Castle	(502) 845-2882	Trigg County Health Department	Cadiz	(270) 522-8121
Hickman County Health Department	Clinton	(270) 653-6110	Trimble County Health Department	Bedford	(502) 255-7701
Hopkins County Health Department	Madisonville	(270) 821-5242	Union County Health Department	Morganfield	(270) 389-1230
Jackson County Health Department	McKee	(606) 287-8421	Warren County Health Department	Bowling Green	(270) 781-2490
(Jefferson Co.) Dixie Health Center	Louisville	(502) 574-5380	Washington County Health Department	Springfield	(859) 336-3989
(Jefferson Co.) Harambee Health Center, Inc.	Louisville	(502) 593-5939	Wayne County Health Department	Monticello	(606) 348-9349
(Jefferson Co.) Lou.-Metro HD - Fam Plan/Methadone	Louisville	(502) 574-6660	Webster County Health Department	Dixon	(270) 639-9315
(Jefferson Co.) Louisville-Metro HD - Specialty	Louisville	(502) 574-6697	Whitley County Health Department	Corbin	(606) 549-3380
(Jefferson Co.) Louisville-Metro HD - TB Clinic	Louisville	(502) 574-6617	Wolfe County Health Department	Campton	(606) 668-3185
(Jefferson Co.) Newburg Health Center	Louisville	(502) 458-0778	Woodford County Health Department	Versailles	(859) 873-4541
(Jefferson Co.) The More Center	Louisville	(502) 574-6414			
(Jefferson Co.) Volunteers of America – Louisville	Louisville	(502) 636-4540			

