

Table 4A
2002 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Racial and Ethnic Distribution by Clinic							
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan %	Asian/Pacific Islander %	All Other / Unknown %
Adair Coun	515	93.2	4.9	1.9	0.0	0.0	0.0
Allen Coun	600	96.8	1.0	1.0	0.0	1.2	0.0
Anderson C	345	91.3	5.2	3.5	0.0	0.0	0.0
Ashland-Bo	1,063	95.9	2.1	1.7	0.2	0.2	0.0
Ballard Co	268	89.2	10.1	0.4	0.0	0.4	0.0
Bar-Riv-Wa	2,281	68.7	15.3	13.3	0.3	2.5	0.0
Barren Cty	995	90.8	6.6	1.8	0.2	0.6	0.0
Bath Count	427	97.9	1.2	0.5	0.5	0.0	0.0
Belfry	432	99.3	0.7	0.0	0.0	0.0	0.0
Bell Cty M	832	96.6	0.1	2.9	0.4	0.0	0.0
Bell Cty P	800	94.6	4.6	0.6	0.0	0.1	0.0
Berea	435	97.2	2.1	0.2	0.0	0.5	0.0
Betsy Lane	424	100.0	0.0	0.0	0.0	0.0	0.0
Bledsoe	92	*	*	*	*	*	*
Boone Coun	1,980	88.5	3.3	7.6	0.0	0.5	0.0
Bourbon Ct	474	78.1	9.1	11.6	0.2	1.1	0.0
Boyle Coun	567	94.2	5.5	0.0	0.0	0.4	0.0
Bracken Co	230	99.6	0.0	0.4	0.0	0.0	0.0
Breathitt	687	99.3	0.6	0.0	0.1	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4A
2002 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution by Clinic							
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan %	Asian/Pacific Islander %	All Other / Unknown %
Breckinrid	492	95.7	3.9	0.4	0.0	0.0	0.0
Bridges	497	11.9	85.5	1.4	0.0	1.2	0.0
Bullitt Co	1,122	99.1	0.5	0.1	0.0	0.3	0.0
Butler Cty	456	96.9	1.1	0.0	0.0	2.0	0.0
C & Y	1,237	4.6	91.1	0.3	1.5	2.4	0.0
Caldwell C	421	92.2	7.4	0.2	0.0	0.2	0.0
Calloway C	919	86.6	7.9	3.4	0.1	2.0	0.0
Campbell C	1,425	92.7	6.0	0.9	0.1	0.3	0.0
Cannonsbur	362	99.7	0.3	0.0	0.0	0.0	0.0
Carlisle C	190	93.7	5.8	0.5	0.0	0.0	0.0
Carroll Ct	405	84.9	1.2	13.8	0.0	0.0	0.0
Carter Cty	692	99.1	0.1	0.0	0.6	0.1	0.0
Casey Cty	552	97.3	0.0	2.7	0.0	0.0	0.0
Christian	2,076	57.5	40.5	1.0	0.1	0.9	0.0
Clark Coun	823	89.6	6.6	3.3	0.0	0.6	0.0
Clay Cty H	905	97.7	1.3	1.0	0.0	0.0	0.0
Clinton Ct	506	95.3	0.2	4.2	0.0	0.4	0.0
Corbin	1,165	99.7	0.2	0.0	0.0	0.2	0.0
Crittenden	256	99.6	0.4	0.0	0.0	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4A
2002 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution by Clinic							
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan %	Asian/Pacific Islander %	All Other / Unknown %
Cumberland	296	93.6	4.1	2.4	0.0	0.0	0.0
Daviess Co	2,226	87.6	8.6	3.1	0.1	0.6	0.0
Dixie	1,546	86.0	12.4	1.0	0.3	0.3	0.0
Edmonson C	362	100.0	0.0	0.0	0.0	0.0	0.0
Elliott Ct	298	99.3	0.0	0.3	0.3	0.0	0.0
Estill Cty	595	98.8	0.7	0.0	0.0	0.5	0.0
Etown	1,297	88.0	9.7	1.7	0.2	0.3	0.0
Evarts	353	98.9	1.1	0.0	0.0	0.0	0.0
Fairdale	924	92.6	5.0	1.4	0.4	0.5	0.0
Fleming Co	551	97.5	1.6	0.9	0.0	0.0	0.0
Floyd Cty	1,142	99.9	0.0	0.1	0.0	0.0	0.0
Foust Hlth	217	62.2	33.2	4.6	0.0	0.0	0.0
Franklin C	1,011	77.0	16.2	4.5	0.3	2.1	0.0
Ft Campbel	2,905	68.8	22.8	7.6	0.1	0.7	0.0
Fulton	203	53.2	46.8	0.0	0.0	0.0	0.0
Gallatin C	264	84.1	2.3	13.3	0.0	0.4	0.0
Garrard Ct	551	94.7	2.4	2.7	0.2	0.0	0.0
Goose Rock	167	98.8	0.0	0.6	0.6	0.0	0.0
Grant Cty	1,061	97.8	0.9	0.4	0.0	0.8	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4A
2002 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution by Clinic							
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan %	Asian/Pacific Islander %	All Other / Unknown %
Graves Cty	1,199	75.6	10.3	11.8	0.1	2.3	0.0
Grayson Ct	901	98.6	0.9	0.4	0.1	0.0	0.0
Green Cty	332	95.2	3.6	0.9	0.0	0.3	0.0
Greenup	813	98.9	1.0	0.0	0.0	0.1	0.0
H & Env.	148	81.8	16.2	1.4	0.0	0.7	0.0
Hancock Ct	258	97.7	0.8	1.2	0.4	0.0	0.0
Harlan Cty	766	97.9	1.6	0.4	0.0	0.1	0.0
Harrison C	536	92.4	2.8	4.9	0.0	0.0	0.0
Hart Cty H	627	95.4	4.3	0.3	0.0	0.0	0.0
Henderson	1,152	84.5	11.9	0.9	0.4	2.3	0.0
Henry Coun	370	88.4	3.5	8.1	0.0	0.0	0.0
Hickman	170	67.1	32.9	0.0	0.0	0.0	0.0
Hickman Ct	127	80.3	17.3	1.6	0.0	0.8	0.0
Highview	1,174	78.4	9.3	10.9	0.1	1.3	0.0
Hindman	720	99.4	0.6	0.0	0.0	0.0	0.0
Hopkins Ct	1,513	84.6	13.0	2.0	0.1	0.3	0.0
Iroquois	1,826	52.4	28.9	11.6	0.1	7.1	0.0
Jackson Ct	640	100.0	0.0	0.0	0.0	0.0	0.0
Jessamine	1,053	90.8	5.2	2.6	0.1	1.3	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4A
2002 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution by Clinic							
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan %	Asian/Pacific Islander %	All Other / Unknown %
Johnson Co	1,038	100.0	0.0	0.0	0.0	0.0	0.0
JT Arnett	63	*	*	*	*	*	*
Kenton Cty	2,018	83.8	14.8	0.9	0.2	0.2	0.0
Knox Cty	1,520	98.8	1.1	0.1	0.0	0.0	0.0
L & N	1,416	20.5	75.2	2.2	0.1	2.0	0.0
Larue Cty	434	89.2	5.5	4.6	0.7	0.0	0.0
Laurel Cty	1,640	98.7	0.7	0.3	0.0	0.3	0.0
Lawrence C	600	99.8	0.2	0.0	0.0	0.0	0.0
Lee Cty He	347	99.7	0.3	0.0	0.0	0.0	0.0
Leslie Cty	650	99.5	0.5	0.0	0.0	0.0	0.0
Letcher Ct	866	99.4	0.5	0.0	0.0	0.1	0.0
Lewis Cty	500	99.6	0.4	0.0	0.0	0.0	0.0
Lex-Fayett	2,174	33.8	29.3	28.9	0.4	7.7	0.0
Lex-Fayett	593	50.3	41.1	3.2	0.3	5.1	0.0
Lex-Fayett	638	63.9	20.2	10.0	0.5	5.3	0.0
Lex-Fayett	839	55.9	36.6	2.9	0.2	4.4	0.0
Lex-Fayett	778	63.8	31.1	1.7	0.4	3.1	0.0
Liberty	53	*	*	*	*	*	*
Lincoln Ct	874	93.6	3.0	3.3	0.0	0.1	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4A
2002 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Racial and Ethnic Distribution by Clinic							
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan %	Asian/Pacific Islander %	All Other / Unknown %
Livingston	198	97.5	0.0	1.0	0.0	1.5	0.0
Logan Cty	716	80.7	16.2	2.4	0.7	0.0	0.0
Lynch	336	92.9	6.8	0.3	0.0	0.0	0.0
Lyon Cty	172	98.8	1.2	0.0	0.0	0.0	0.0
Magoffin C	664	99.7	0.0	0.3	0.0	0.0	0.0
Marion Cty	677	83.3	11.5	4.7	0.0	0.4	0.0
Marshall C	744	100.0	0.0	0.0	0.0	0.0	0.0
Marshall C	115	96.5	3.5	0.0	0.0	0.0	0.0
Martin Cty	711	99.9	0.0	0.1	0.0	0.0	0.0
Mason Cty	592	90.9	7.3	0.7	0.5	0.7	0.0
McCracken	1,988	63.5	33.0	1.6	0.5	1.4	0.0
McCreary C	884	99.8	0.2	0.0	0.0	0.0	0.0
McLean Ct	452	97.3	0.7	1.8	0.0	0.2	0.0
Meade Cty	697	97.1	2.9	0.0	0.0	0.0	0.0
Menifee Ct	286	99.0	0.0	1.0	0.0	0.0	0.0
Mercer Cty	592	89.2	4.7	5.7	0.0	0.3	0.0
Metcalfe C	350	97.7	1.1	0.9	0.3	0.0	0.0
Middletown	918	74.2	21.5	3.5	0.1	0.8	0.0
Monroe Cnt	408	88.7	5.6	5.4	0.2	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4A
2002 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution by Clinic							
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan %	Asian/Pacific Islander %	All Other / Unknown %
Montgomery	872	95.5	1.3	2.6	0.0	0.6	0.0
Morgan Cty	564	99.8	0.0	0.2	0.0	0.0	0.0
Muhlenberg	1,033	96.6	3.0	0.3	0.0	0.1	0.0
Neighborhood	1,155	64.6	30.0	2.3	0.1	3.0	0.0
Nelson Cty	1,025	89.7	8.5	1.4	0.2	0.3	0.0
Newburg	1,156	32.7	55.1	10.4	0.1	1.7	0.0
Nicholas C	285	98.9	0.0	1.1	0.0	0.0	0.0
Oasis Hlth	203	86.2	12.8	1.0	0.0	0.0	0.0
Ohio Cty	830	95.3	0.4	3.7	0.0	0.6	0.0
Oldham Cty	588	82.8	5.1	11.7	0.0	0.3	0.0
Olive Hill	564	99.3	0.7	0.0	0.0	0.0	0.0
Oneida Ele	67	*	*	*	*	*	*
Owen Cty	280	92.9	0.0	6.4	0.4	0.4	0.0
Owsley Cty	272	99.6	0.4	0.0	0.0	0.0	0.0
Park Duval	1,528	13.4	78.7	3.3	0.9	3.7	0.0
Pendleton	404	96.0	2.0	1.0	0.7	0.2	0.0
Perry Cty	1,115	97.8	1.9	0.3	0.0	0.1	0.0
Phelps	152	100.0	0.0	0.0	0.0	0.0	0.0
Pike Cty	1,487	99.5	0.3	0.0	0.1	0.1	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4A
2002 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Racial and Ethnic Distribution by Clinic							
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan %	Asian/Pacific Islander %	All Other / Unknown %
Portland	1,183	47.4	51.7	0.6	0.1	0.2	0.0
Powell Cty	681	97.8	0.6	0.9	0.0	0.7	0.0
Prenatal	94	*	*	*	*	*	*
Prenatal	33	*	*	*	*	*	*
Pulaski Ct	1,901	97.4	0.9	1.5	0.1	0.1	0.0
Radcliffe	2,179	71.6	25.4	2.2	0.0	0.8	0.0
Richmond	1,350	87.6	9.5	1.3	0.1	1.5	0.0
Robertson	111	94.6	0.9	4.5	0.0	0.0	0.0
Rockcastle	632	99.7	0.2	0.2	0.0	0.0	0.0
Rowan Cty	690	97.1	1.0	1.2	0.4	0.3	0.0
Russell Ct	607	95.6	0.2	3.1	0.2	1.0	0.0
Scott Cty	723	86.7	7.6	4.8	0.4	0.4	0.0
Shelby Cty	791	52.2	13.5	34.3	0.0	0.0	0.0
Siler	49	*	*	*	*	*	*
Simpson Ct	446	83.6	14.6	0.9	0.2	0.7	0.0
South Shor	163	100.0	0.0	0.0	0.0	0.0	0.0
Spencer Ct	275	92.7	1.5	5.1	0.4	0.4	0.0
TAPP-South	58	*	*	*	*	*	*
TAPP-West	67	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4A
2002 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Racial and Ethnic Distribution by Clinic							
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan %	Asian/Pacific Islander %	All Other / Unknown %
Taylor Cty	783	92.3	7.4	0.1	0.1	0.0	0.0
Todd Cty H	460	81.5	11.5	5.4	0.2	1.3	0.0
Trigg Cty	301	82.7	15.6	0.0	0.7	1.0	0.0
Trimble Ct	247	96.4	0.4	3.2	0.0	0.0	0.0
Union Cty	494	85.4	13.6	0.2	0.0	0.8	0.0
Wallins	111	100.0	0.0	0.0	0.0	0.0	0.0
Washington	374	75.7	12.6	11.5	0.0	0.3	0.0
Wayne Cty	879	92.3	1.6	6.1	0.0	0.0	0.0
Webster Ct	428	85.7	4.2	7.2	0.0	2.8	0.0
Wheelbrigh	288	97.2	2.1	0.7	0.0	0.0	0.0
Williamsbu	964	99.8	0.1	0.1	0.0	0.0	0.0
Wolfe Cty	449	99.3	0.7	0.0	0.0	0.0	0.0
Woodford C	533	78.2	4.7	16.5	0.2	0.4	0.0
Unknown Clinics	1,324	83.8	12.8	2.7	0.0	0.6	0.0
Kentucky	120,031	82.9	12.6	3.4	0.1	0.9	0.0
Nation (Prior Year)	5,174,210	40.2	21.4	31.4	1.4	2.8	2.9

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2002 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Adair Coun	515	29.1	25.6	45.2
Allen Coun	600	24.8	24.3	50.8
Anderson C	345	31.0	25.2	43.8
Ashland-Bo	1,063	33.8	25.6	40.6
Ballard Co	268	25.7	28.7	45.5
Bar-Riv-Wa	2,281	39.5	23.3	37.2
Barren Cty	995	32.3	26.3	41.4
Bath Count	427	25.5	27.6	46.8
Belfry	432	28.9	25.5	45.6
Bell Cty M	832	24.5	24.9	50.6
Bell Cty P	800	22.1	25.9	52.0
Berea	435	30.3	26.0	43.7
Betsy Lane	424	19.6	27.4	53.1
Bledsoe	92	*	*	*
Boone Coun	1,980	31.7	25.5	42.8
Bourbon Ct	474	27.0	32.3	40.7
Boyle Coun	567	35.1	26.3	38.6
Bracken Co	230	33.9	23.5	42.6

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2002 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Breathitt	687	24.6	21.1	54.3
Breckinrid	492	27.2	26.4	46.3
Bridges	497	34.2	29.0	36.8
Bullitt Co	1,122	36.5	25.6	37.9
Butler Cty	456	28.9	25.2	45.8
C & Y	1,237	36.3	23.1	40.6
Caldwell C	421	29.9	27.3	42.8
Calloway C	919	30.5	26.7	42.9
Campbell C	1,425	35.2	23.4	41.3
Cannonsbur	362	34.3	26.0	39.8
Carlisle C	190	20.0	25.8	54.2
Carroll Ct	405	28.9	28.4	42.7
Carter Cty	692	27.7	25.9	46.4
Casey Cty	552	29.0	23.2	47.8
Christian	2,076	37.5	23.8	38.7
Clark Coun	823	36.6	23.6	39.9
Clay Cty H	905	24.4	27.7	47.8
Clinton Ct	506	23.3	27.7	49.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2002 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Corbin	1,165	28.2	24.5	47.4
Crittenden	256	26.2	28.1	45.7
Cumberland	296	23.6	28.4	48.0
Daviess Co	2,226	35.7	24.3	39.9
Dixie	1,546	39.3	23.9	36.8
Edmonson C	362	39.0	19.6	41.4
Elliott Ct	298	21.8	26.5	51.7
Estill Cty	595	30.3	24.2	45.5
Etown	1,297	37.5	22.5	39.9
Evarts	353	21.8	30.0	48.2
Fairdale	924	40.0	23.7	36.3
Fleming Co	551	24.0	25.4	50.6
Floyd Cty	1,142	28.7	26.0	45.3
Foust Hlth	217	18.9	26.7	54.4
Franklin C	1,011	40.7	22.8	36.5
Ft Campbel	2,905	32.8	21.3	45.8
Fulton	203	36.5	25.6	37.9
Gallatin C	264	32.2	23.5	44.3

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2002 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Garrard Ct	551	25.0	27.8	47.2
Goose Rock	167	21.6	21.0	57.5
Grant Cty	1,061	22.2	27.0	50.7
Graves Cty	1,199	27.9	26.4	45.6
Grayson Ct	901	25.0	26.1	48.9
Green Cty	332	30.1	28.0	41.9
Greenup	813	34.2	23.4	42.4
H & Env.	148	23.6	29.7	46.6
Hancock Ct	258	27.5	23.3	49.2
Harlan Cty	766	27.5	21.5	50.9
Harrison C	536	27.1	27.4	45.5
Hart Cty H	627	33.2	23.6	43.2
Henderson	1,152	38.6	22.3	39.1
Henry Coun	370	30.0	30.0	40.0
Hickman	170	25.9	22.4	51.8
Hickman Ct	127	30.7	26.8	42.5
Highview	1,174	35.9	22.4	41.7
Hindman	720	25.3	21.0	53.8

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2002 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Hopkins Ct	1,513	31.0	23.5	45.5
Iroquois	1,826	37.7	23.3	38.9
Jackson Ct	640	21.9	29.5	48.6
Jessamine	1,053	32.0	24.3	43.7
Johnson Co	1,038	24.6	26.0	49.4
JT Arnett	63	*	*	*
Kenton Cty	2,018	32.4	26.5	41.1
Knox Cty	1,520	25.6	24.5	49.9
L & N	1,416	41.2	21.0	37.9
Larue Cty	434	31.3	24.0	44.7
Laurel Cty	1,640	36.3	23.3	40.4
Lawrence C	600	27.2	28.3	44.5
Lee Cty He	347	23.3	27.1	49.6
Leslie Cty	650	24.2	22.0	53.8
Letcher Ct	866	31.8	23.0	45.3
Lewis Cty	500	27.0	26.8	46.2
Lex-Fayett	2,174	41.2	25.3	33.5
Lex-Fayett	593	32.9	29.5	37.6

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2002 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Lex-Fayette	638	33.5	25.9	40.6
Lex-Fayette	839	34.4	26.3	39.2
Lex-Fayette	778	35.5	24.3	40.2
Liberty	53	*	*	*
Lincoln Ct	874	33.9	22.2	43.9
Livingston	198	25.3	31.3	43.4
Logan Cty	716	33.7	28.6	37.7
Lynch	336	25.9	21.1	53.0
Lyon Cty	172	29.1	27.3	43.6
Magoffin C	664	23.0	23.8	53.2
Marion Cty	677	28.8	25.1	46.1
Marshall C	744	29.3	27.2	43.5
Marshall C	115	31.3	25.2	43.5
Martin Cty	711	23.5	23.1	53.4
Mason Cty	592	30.2	26.4	43.4
McCracken	1,988	29.3	28.0	42.7
McCreary C	884	24.3	26.5	49.2
McLean Ct	452	24.3	27.7	48.0

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2002 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Meade Cty	697	31.7	23.4	44.9
Menifee Ct	286	22.7	26.2	51.0
Mercer Cty	592	33.8	22.6	43.6
Metcalfe C	350	31.1	23.7	45.1
Middletown	918	39.9	26.3	33.9
Monroe Cnt	408	31.4	22.5	46.1
Montgomery	872	32.7	22.9	44.4
Morgan Cty	564	23.2	25.5	51.2
Muhlenberg	1,033	27.0	28.1	44.9
Neighborho	1,155	35.8	25.3	39.0
Nelson Cty	1,025	32.4	23.6	44.0
Newburg	1,156	40.3	26.9	32.8
Nicholas C	285	32.3	23.9	43.9
Oasis Hlth	203	19.7	20.7	59.6
Ohio Cty	830	30.1	22.5	47.3
Oldham Cty	588	29.6	27.0	43.4
Olive Hill	564	24.3	25.2	50.5
Oneida Ele	67	*	*	*

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2002 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Owen Cty	280	32.9	22.9	44.3
Owsley Cty	272	21.7	26.1	52.2
Park Duval	1,528	33.7	22.0	44.3
Pendleton	404	27.5	30.0	42.6
Perry Cty	1,115	26.0	26.5	47.4
Phelps	152	33.6	13.2	53.3
Pike Cty	1,487	33.4	22.8	43.8
Portland	1,183	37.5	24.3	38.1
Powell Cty	681	27.6	24.1	48.3
Prenatal	94	*	*	*
Prenatal	33	*	*	*
Pulaski Ct	1,901	35.7	24.1	40.1
Radcliffe	2,179	31.4	21.5	47.1
Richmond	1,350	37.7	22.8	39.5
Robertson	111	25.2	26.1	48.6
Rockcastle	632	33.7	22.8	43.5
Rowan Cty	690	31.9	24.1	44.1
Russell Ct	607	29.7	28.7	41.7

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2002 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Scott Cty	723	35.8	27.2	36.9
Shelby Cty	791	38.2	26.9	34.9
Siler	49	*	*	*
Simpson Ct	446	34.3	27.4	38.3
South Shor	163	26.4	23.9	49.7
Spencer Ct	275	28.7	28.0	43.3
TAPP-South	58	*	*	*
TAPP-West	67	*	*	*
Taylor Cty	783	27.2	26.6	46.2
Todd Cty H	460	23.7	30.2	46.1
Trigg Cty	301	28.9	26.9	44.2
Trimble Ct	247	24.7	29.6	45.7
Union Cty	494	31.2	25.9	42.9
Wallins	111	26.1	31.5	42.3
Washington	374	25.9	29.1	44.9
Wayne Cty	879	29.8	22.1	48.1
Webster Ct	428	37.4	20.3	42.3
Wheelbrigh	288	25.0	21.9	53.1

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2002 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Williamsbu	964	28.0	22.1	49.9
Wolfe Cty	449	24.1	26.3	49.7
Woodford C	533	30.0	30.8	39.2
Unknown Clinics	1,324	89.7	3.2	7.1
Kentucky	120,031	32.4	24.5	43.1
Nation (Prior Year)	5,174,210	37.3	21.6	41.1

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2002 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Adair Coun	135	6.7 (13)	6.7 (39)	501	6.8(67)	5.2(10)	12.4(14)	223	13.9(19)	11.7(13)	305	5.2 (18)
Allen Coun	138	10.9 (80)	10.1 (97)	585	7.4(80)	3.4(41)	17.1(91)	290	22.8(13)	19.0(10)	368	8.4 (61)
Anderson C	81	* *	* *	327	9.2(12)	5.8(11)	20.8(13)	147	19.0(90)	23.8(13)	253	5.9 (29)
Ashland-Bo	319	12.5(102)	6.9 (45)	1,014	7.7(85)	9.9(15)	15.8(60)	422	19.0(90)	17.8(87)	654	10.9 (85)
Ballard Co	62	* *	* *	250	7.6(82)	6.0(12)	16.4(77)	119	9.2 (2)	10.9(10)	198	4.5 (12)
Bar-Riv-Wa	728	11.0 (84)	6.6 (35)	2,202	7.2(75)	3.4(41)	15.7(58)	836	17.5(66)	16.3(68)	1,337	13.4(108)
Barren Cty	250	9.6 (52)	7.2 (57)	949	9.1(11)	3.6(52)	18.2(10)	397	17.6(69)	17.4(82)	671	9.1 (68)
Bath Count	100	6.0 (8)	16.0(115)	416	12.3(15)	2.4(18)	20.4(12)	200	19.5(10)	20.5(11)	272	9.2 (69)
Belfry	115	11.3 (89)	10.4 (99)	413	1.5 (1)	10.4(15)	14.3(33)	187	14.4(25)	13.9(46)	280	8.2 (56)
Bell Cty M	188	11.7 (93)	5.9 (21)	789	7.2(75)	3.7(57)	21.2(13)	411	20.9(11)	21.2(11)	546	2.7 (4)
Bell Cty P	146	6.2 (10)	7.5 (63)	774	7.6(82)	3.5(47)	18.6(11)	408	13.5(17)	13.0(28)	535	12.5(103)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2002 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic												(3,4)
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number
Berea	109	8.3 (32)	11.0 (103)	428	7.2 (75)	4.4 (87)	13.1 (20)	183	24.6 (14)	16.4 (71)	318	8.2 (56)
Betsy Lane	81	* *	* *	409	2.7 (4)	10.0 (15)	12.5 (15)	220	9.5 (5)	11.8 (17)	310	9.0 (66)
Bledsoe	23	* *	* *	88	* *	* *	* *	52	* *	* *	53	* *
Boone Coun	487	10.3 (67)	9.2 (91)	1,895	8.6 (10)	4.9 (10)	18.3 (10)	826	18.9 (87)	18.3 (92)	1,384	14.7 (118)
Bourbon Ct	117	11.1 (86)	9.4 (93)	450	8.9 (11)	2.0 (11)	21.1 (13)	178	22.5 (13)	21.9 (12)	342	29.8 (149)
Boyle Coun	181	12.7 (105)	6.6 (35)	527	9.1 (11)	5.3 (11)	16.3 (74)	199	19.1 (94)	18.1 (89)	325	8.3 (60)
Bracken Co	69	* *	* *	220	8.2 (95)	6.8 (13)	22.7 (14)	95	* *	* *	146	4.1 (8)
Breathitt	154	8.4 (33)	9.1 (88)	639	6.4 (56)	3.3 (36)	20.0 (12)	361	14.4 (25)	17.2 (80)	506	5.3 (19)
Breckinrid	113	8.0 (27)	6.2 (27)	481	7.5 (81)	3.3 (36)	15.0 (49)	226	21.2 (11)	13.3 (34)	367	3.3 (6)
Bridges	161	10.6 (73)	6.2 (27)	473	12.1 (14)	4.7 (95)	16.1 (66)	183	16.9 (53)	11.5 (12)	316	6.0 (30)
Bullitt Co	368	8.2 (30)	9.0 (86)	1,067	8.2 (95)	3.6 (52)	16.8 (86)	410	15.4 (38)	17.6 (84)	625	19.2 (136)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2002 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Butler Cty	115	6.1 (9)	7.0 (51)	446	5.4(33)	4.7(95)	19.5(12)	204	19.1(94)	24.5(14)	306	7.2 (44)
C & Y	389	10.0 (62)	5.7 (20)	1,174	8.2(95)	5.7(11)	16.3(74)	491	18.9(87)	20.4(10)	300	21.0(140)
Caldwell C	108	2.8 (1)	7.4 (61)	405	6.7(62)	7.2(13)	16.3(74)	178	15.2(36)	16.3(68)	298	15.8(125)
Calloway C	243	8.6 (38)	13.2(112)	892	7.0(72)	3.5(47)	14.9(47)	388	17.3(62)	15.2(55)	582	16.2(127)
Campbell C	385	11.2 (87)	8.1 (70)	1,340	8.4(99)	3.3(36)	18.1(10)	563	20.2(10)	23.3(13)	873	14.7(118)
Cannonsbur	106	7.5 (19)	9.4 (93)	343	4.1(18)	11.4(15)	9.6 (4)	139	15.1(34)	13.7(41)	226	11.1 (87)
Carlisle C	35	* *	* *	183	6.0(46)	2.7(21)	22.4(14)	102	18.6(80)	24.5(14)	138	24.6(147)
Carroll Ct	106	8.5 (36)	4.7 (8)	396	11.9(14)	3.8(63)	20.7(13)	171	25.1(14)	24.0(13)	295	11.5 (89)
Carter Cty	175	6.9 (14)	9.7 (95)	653	3.4 (9)	8.6(14)	10.4 (6)	302	9.3 (3)	11.3(11)	438	8.9 (64)
Casey Cty	136	10.3 (67)	3.7 (5)	532	9.4(12)	4.7(95)	13.2(23)	254	15.0(33)	13.0(28)	337	11.6 (92)
Christian	516	12.0 (98)	5.0 (11)	1,971	8.8(10)	5.8(11)	16.4(77)	794	13.6(18)	12.5(22)	1,383	35.2(151)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2002 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic												(3,4)
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Clark Coun	255	14.1 (110)	8.6 (76)	791	7.7(85)	3.7(57)	15.7(58)	315	21.9(12)	15.2(55)	495	12.7(105)
Clay Cty H	211	11.4 (90)	6.6 (35)	876	6.7(62)	5.5(11)	21.5(13)	420	20.7(11)	20.5(11)	658	8.2 (56)
Clinton Ct	111	9.0 (46)	12.6(111)	486	6.0(46)	4.3(81)	16.7(83)	239	16.3(43)	14.2(49)	308	5.8 (28)
Corbin	243	6.2 (10)	9.1 (88)	1,139	5.4(33)	5.0(10)	15.5(54)	530	14.2(23)	15.1(53)	874	5.7 (25)
Crittenden	62	* *	* *	248	5.6(38)	6.5(12)	12.1(11)	110	16.4(44)	6.4 (2)	181	11.0 (86)
Cumberland	66	* *	* *	290	3.4 (9)	3.8(63)	14.8(46)	140	17.1(57)	13.6(38)	160	13.8(114)
Daviess Co	652	8.4 (33)	5.2 (12)	2,141	11.0(13)	2.0(11)	21.9(14)	877	16.4(44)	21.0(11)	1,484	13.3(107)
Dixie	509	10.6 (73)	7.7 (66)	1,471	8.8(10)	3.7(57)	14.2(30)	543	13.4(16)	15.8(62)	952	12.6(104)
Edmonson C	102	9.8 (56)	5.9 (21)	350	4.9(28)	4.9(10)	13.1(20)	146	18.5(78)	15.1(53)	219	4.6 (13)
Elliott Ct	55	* *	* *	287	3.1 (6)	4.9(10)	14.3(33)	151	17.2(60)	13.2(32)	213	9.9 (77)
Estill Cty	158	6.3 (12)	7.6 (64)	578	3.8(15)	4.8(99)	15.9(62)	267	18.0(74)	18.4(93)	421	7.1 (42)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2002 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic												(3,4)
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Etown	382	10.5 (71)	7.9 (68)	1,228	9.2(12)	2.2(15)	15.9(62)	512	20.7(11)	13.5(36)	884	7.6 (50)
Evarts	71	* *	* *	340	5.3(32)	9.7(14)	22.1(14)	166	14.5(27)	21.7(12)	272	9.6 (74)
Fairdale	331	9.1 (48)	8.8 (80)	877	5.8(43)	4.4(87)	14.4(36)	323	14.9(31)	12.7(25)	580	5.0 (17)
Fleming Co	127	7.1 (16)	11.8(108)	539	6.7(62)	5.2(10)	14.5(38)	275	16.4(44)	16.0(64)	418	16.7(129)
Floyd Cty	297	7.4 (18)	6.4 (31)	1,091	4.9(28)	10.3(15)	13.0(18)	485	12.0(13)	13.6(38)	819	4.9 (16)
Foust Hlth	37	* *	* *	209	8.1(94)	3.8(63)	19.1(11)	119	16.8(49)	16.0(64)	178	21.3(141)
Franklin C	308	8.1 (29)	6.2 (27)	968	8.9(11)	7.0(13)	18.6(11)	363	18.7(82)	21.5(12)	683	13.2(106)
Ft Campbel	715	6.9 (14)	11.2(105)	2,817	3.4 (9)	7.6(14)	6.0 (1)	1,298	11.3 (8)	5.5 (1)	1,951	20.7(138)
Fulton	53	* *	* *	191	6.3(52)	4.2(79)	22.5(14)	79	* *	* *	128	7.0 (40)
Gallatin C	74	* *	* *	255	5.5(36)	5.1(10)	17.3(93)	115	10.4 (6)	20.9(11)	173	11.6 (92)
Garrard Ct	119	8.4 (33)	6.7 (39)	527	4.9(28)	3.8(63)	16.1(66)	255	16.9(53)	18.4(93)	340	11.5 (89)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2002 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Goose Rock	35	* *	* *	157	2.5 (3)	9.6(14)	14.6(42)	90	* *	* *	132	6.1 (31)
Grant Cty	207	9.7 (55)	7.2 (57)	1,026	5.6(38)	5.6(11)	11.2 (8)	521	8.1 (1)	7.7 (5)	836	5.5 (22)
Graves Cty	313	8.6 (38)	8.9 (83)	1,145	7.9(89)	5.9(12)	17.9(10)	532	20.5(10)	18.2(90)	771	15.6(123)
Grayson Ct	201	9.0 (46)	6.5 (34)	879	13.2(15)	2.2(15)	20.4(12)	432	19.0(90)	17.6(84)	680	2.4 (3)
Green Cty	93	* *	* *	318	6.3(52)	1.9 (9)	23.3(14)	138	22.5(13)	24.6(14)	207	7.2 (44)
Greenup	214	11.7 (93)	8.9 (83)	786	3.3 (8)	8.5(14)	13.0(18)	343	19.0(90)	14.3(51)	535	7.1 (42)
H & Env.	21	* *	* *	145	5.5(36)	5.5(11)	19.3(11)	70	* *	* *	122	23.8(146)
Hancock Ct	58	* *	* *	249	6.8(67)	4.8(99)	16.1(66)	127	16.5(47)	14.2(49)	196	4.1 (8)
Harlan Cty	168	7.7 (23)	12.5(110)	732	2.3 (2)	6.3(12)	19.7(12)	372	15.9(41)	21.0(11)	504	6.3 (33)
Harrison C	123	13.8(109)	8.1 (70)	509	4.1(18)	6.1(12)	9.6 (4)	245	11.4(10)	6.9 (4)	384	13.8(114)
Hart Cty H	184	7.6 (22)	8.7 (77)	613	6.4(56)	3.6(52)	15.5(54)	270	14.8(30)	17.0(79)	401	5.7 (25)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2002 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic												(3,4)
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Henderson	362	10.5 (71)	5.2 (12)	1,120	8.9(11)	4.5(91)	15.9(62)	438	22.1(13)	22.1(12)	271	1.8 (1)
Henry Coun	86	* *	* *	350	16.9(15)	3.4(41)	25.1(14)	141	20.6(10)	22.7(12)	273	10.3 (80)
Hickman	42	* *	* *	162	6.8(67)	1.2 (5)	17.3(93)	86	* *	* *	120	23.3(145)
Hickman Ct	36	* *	* *	121	4.1(18)	5.8(11)	10.7 (7)	54	* *	* *	88	* *
Highview	329	9.1 (48)	6.4 (31)	1,120	6.4(56)	4.6(94)	15.6(57)	473	15.4(38)	10.6 (9)	744	13.4(108)
Hindman	142	10.6 (73)	4.2 (6)	681	10.7(13)	4.3(81)	20.4(12)	370	11.9(12)	13.8(44)	570	8.2 (56)
Hopkins Ct	346	14.5(111)	3.2 (2)	1,429	13.7(15)	1.5 (6)	21.6(13)	664	21.1(11)	18.7(97)	1,087	12.3 (97)
Iroquois	582	9.8 (56)	6.2 (27)	1,748	6.2(51)	2.7(21)	16.4(77)	682	18.2(75)	22.0(12)	1,203	12.3 (97)
Jackson Ct	131	11.5 (91)	10.7(100)	627	6.7(62)	1.6 (8)	17.7(99)	301	14.0(21)	15.9(63)	502	3.0 (5)
Jessamine	263	5.3 (3)	8.7 (77)	1,013	8.5(10)	2.3(17)	17.5(96)	449	18.9(87)	17.6(84)	742	9.6 (74)
Johnson Co	233	9.9 (61)	10.3 (98)	998	5.4(33)	5.9(12)	17.8(10)	505	14.9(31)	16.8(75)	769	16.6(128)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2002 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic												(3,4)
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
JT Arnett	10	* *	* *	58	* *	* *	* *	25	* *	* *	49	* *
Kenton Cty	486	12.8 (107)	8.4 (74)	1,922	8.5(10)	3.5(47)	20.0(12)	816	22.2(13)	23.5(13)	1,469	13.7 (112)
Knox Cty	334	12.6 (104)	6.0 (24)	1,471	6.1 (49)	3.2(34)	17.9(10)	740	17.7(72)	20.0(10)	1,116	11.6 (92)
L & N	492	12.0 (98)	6.9 (45)	1,329	6.9(70)	6.6(13)	13.8(27)	507	18.7(82)	15.6(60)	847	12.3 (97)
Larue Cty	110	5.5 (4)	7.3 (59)	413	9.0(11)	4.4(87)	20.6(13)	189	17.5(66)	18.0(88)	323	9.0 (66)
Laurel Cty	481	9.8 (56)	9.1 (88)	1,591	6.9(70)	5.2(10)	14.5(38)	635	20.6(10)	19.2(10)	1,019	14.6(117)
Lawrence C	154	11.7 (93)	4.5 (7)	575	5.2(31)	4.9(10)	20.3(12)	262	18.7(82)	21.0(11)	402	7.5 (48)
Lee Cty He	63	* *	* *	338	9.2(12)	0.9 (2)	26.6(15)	174	24.1(13)	21.8(12)	280	15.4(122)
Leslie Cty	128	7.8 (25)	11.7(107)	621	3.9(16)	3.1(30)	22.7(14)	343	19.2(97)	23.0(12)	513	8.4 (61)
Letcher Ct	204	7.8 (25)	6.9 (45)	829	11.9(14)	3.1(30)	25.8(15)	381	17.3(62)	21.3(11)	638	4.1 (8)
Lewis Cty	125	10.4 (70)	6.4 (31)	480	7.1(74)	2.9(27)	16.7(83)	218	22.0(12)	20.2(10)	315	4.8 (15)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2002 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Lex-Fayette	746	11.0 (84)	6.7 (39)	2,083	7.8(87)	4.0(73)	15.3(51)	710	20.3(10)	19.2(10)	1,270	5.4 (21)
Lex-Fayette	150	12.7 (105)	6.0 (24)	561	9.3(12)	3.7(57)	14.6(42)	220	17.3(62)	13.2(32)	364	31.0 (150)
Lex-Fayette	174	10.9 (80)	10.9 (101)	610	7.2(75)	3.6(52)	14.3(33)	256	19.5(10)	13.7(41)	411	17.5 (132)
Lex-Fayette	245	15.1 (112)	7.3 (59)	741	6.3(52)	4.3(81)	14.2(30)	303	21.8(12)	13.5(36)	475	13.9 (116)
Lex-Fayette	239	15.9 (113)	8.8 (80)	796	4.6(24)	4.4(87)	11.8(10)	321	16.8(49)	12.8(26)	553	28.9 (148)
Liberty	33	* *	* *	48	* *	* *	* *	9	* *	* *	17	* *
Lincoln Ct	227	11.9 (96)	8.4 (74)	834	7.6(82)	5.0(10)	18.5(10)	369	17.1(57)	19.8(10)	626	7.7 (52)
Livingston	44	* *	* *	193	4.7(26)	7.8(14)	16.1(66)	83	* *	* *	153	6.5 (35)
Logan Cty	199	7.5 (19)	3.5 (3)	691	6.5(59)	3.3(36)	16.2(72)	257	21.8(12)	16.3(68)	465	17.6 (133)
Lynch	81	* *	* *	322	9.0(11)	4.3(81)	16.1(66)	175	14.3(24)	18.9(98)	245	2.0 (2)
Lyon Cty	38	* *	* *	162	9.3(12)	4.3(81)	18.5(10)	74	* *	* *	123	6.5 (35)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2002 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Magoffin C	137	8.8 (44)	8.8 (80)	645	11.2(14)	4.0(73)	27.1(15)	338	16.6(48)	23.7(13)	462	21.4(142)
Marion Cty	159	11.9 (96)	6.9 (45)	660	11.5(14)	3.8(63)	22.6(14)	308	20.1(10)	17.2(80)	478	10.5 (82)
Marshall C	196	7.7 (23)	7.1 (55)	716	11.3(14)	7.1(13)	13.8(27)	317	11.4(10)	6.6 (3)	478	6.9 (39)
Marshall C	32	* *	* *	106	10.4(13)	3.8(63)	18.9(11)	48	* *	* *	80	* *
Martin Cty	149	9.4 (51)	4.7 (8)	683	5.9(44)	3.1(30)	17.6(98)	363	17.6(69)	18.5(96)	544	5.5 (22)
Mason Cty	169	5.9 (6)	8.3 (73)	573	7.9(89)	7.5(13)	16.2(72)	254	16.9(53)	13.4(35)	389	11.8 (95)
McCracken	537	12.5(102)	7.8 (67)	1,840	8.2(95)	4.2(79)	21.5(13)	815	17.4(65)	16.8(75)	1,396	18.1(134)
McCreary C	184	10.9 (80)	10.9(101)	855	9.1(11)	3.4(41)	14.5(38)	420	21.2(11)	13.8(44)	663	8.1 (55)
McLean Ct	99	* *	* *	428	11.2(14)	2.8(23)	21.0(13)	202	21.8(12)	24.3(13)	329	4.0 (7)
Meade Cty	172	8.7 (41)	3.5 (3)	677	10.8(13)	3.4(41)	19.1(11)	310	25.2(14)	18.4(93)	518	13.7(112)
Menifee Ct	62	* *	* *	280	4.6(24)	3.9(71)	12.9(17)	140	18.6(80)	13.6(38)	189	10.1 (78)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2002 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic												(3,4)
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Mercer Cty	165	11.5 (91)	5.5 (17)	575	8.7(10)	4.3(81)	16.0(65)	249	23.7(13)	18.9(98)	410	7.8 (53)
Metcalfe C	92	* *	* *	333	5.7(41)	3.6(52)	13.2(23)	149	21.5(12)	16.8(75)	225	9.3 (71)
Middletown	321	12.1 (101)	5.9 (21)	882	7.3(79)	7.0(13)	12.7(16)	304	16.1(42)	15.5(58)	530	15.7(124)
Monroe Cnt	99	* *	* *	398	8.0(91)	3.8(63)	13.6(25)	185	15.1(34)	11.9(18)	286	11.5 (89)
Montgomery	229	9.6 (52)	7.0 (51)	831	6.7(62)	3.2(34)	14.2(30)	379	13.2(15)	12.9(27)	622	9.2 (69)
Morgan Cty	106	8.5 (36)	11.3(106)	544	6.1(49)	3.7(57)	23.2(14)	281	19.2(97)	22.4(12)	444	20.3(137)
Muhlenberg	256	9.8 (56)	9.0 (86)	1,004	4.3(21)	7.2(13)	12.1(11)	466	14.6(28)	13.7(41)	661	9.5 (72)
Neighborho	330	8.2 (30)	7.9 (68)	1,113	8.6(10)	4.0(73)	16.7(83)	439	23.0(13)	16.2(67)	798	5.3 (19)
Nelson Cty	271	10.0 (62)	8.1 (70)	996	8.7(10)	3.5(47)	18.8(11)	445	17.5(66)	14.6(52)	715	7.0 (40)
Newburg	383	11.2 (87)	6.8 (43)	1,079	10.4(13)	5.9(12)	14.6(42)	360	13.9(19)	12.5(22)	709	14.8(120)
Nicholas C	78	* *	* *	274	4.7(26)	10.6(15)	11.7 (9)	128	9.4 (4)	11.7(13)	202	12.4(100)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2002 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)													
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)			
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)						
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Oasis Hlth	33	* *	* *	201	8.0(91)	1.0 (4)	16.9(88)	116	21.6(12)	10.3 (8)		169	8.9 (64)
Ohio Cty	209	9.6 (52)	11.0(103)	800	6.0(46)	2.8(23)	15.5(54)	389	18.3(77)	16.5(72)		601	10.5 (82)
Oldham Cty	150	10.7 (78)	6.7 (39)	558	12.7(15)	2.9(27)	18.3(10)	241	22.0(12)	16.6(74)		373	15.8(125)
Olive Hill	125	5.6 (5)	13.6(113)	552	4.5(23)	5.6(11)	14.5(38)	279	21.1(11)	14.0(47)		355	10.4 (81)
Oneida Ele	11	* *	* *	65	* *	* *	* *	41	* *	* *		56	* *
Owen Cty	86	* *	* *	271	3.0 (5)	8.1(14)	14.4(36)	123	17.1(57)	15.4(57)		191	6.3 (33)
Owsley Cty	51	* *	* *	259	9.7(12)	1.5 (6)	29.3(15)	137	21.2(11)	24.1(13)		219	21.5(143)
Park Duval	428	18.7(115)	6.1 (26)	1,462	11.8(14)	4.1(78)	15.4(53)	673	19.9(10)	15.6(60)		1,062	9.5 (72)
Pendleton	102	9.8 (56)	6.9 (45)	397	9.3(12)	4.0(73)	13.6(25)	170	18.8(85)	14.1(48)		297	7.4 (47)
Perry Cty	263	10.6 (73)	5.3 (15)	1,034	11.2(14)	2.1(13)	32.8(15)	512	18.2(75)	21.9(12)		813	8.0 (54)
Phelps	42	* *	* *	143	7.0(72)	15.4(15)	14.7(45)	79	* *	* *		94	* *

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2002 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic												(3,4)	
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)			
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Pike Cty	397	10.1 (66)	7.6 (64)	1,404	3.7(12)	7.5(13)	18.4(10)	622	20.6(10)	24.3(13)		989	5.7 (25)
Portland	362	10.8 (79)	5.5 (17)	1,126	10.2(13)	4.5(91)	19.6(12)	442	19.2(97)	20.1(10)		799	18.5(135)
Powell Cty	173	7.5 (19)	5.2 (12)	659	3.9(16)	3.3(36)	15.3(51)	330	15.2(36)	15.5(58)		501	6.2 (32)
Prenatal	94	* *	* *	75	* *	* *	* *	0	* *	* *		0	* *
Prenatal	32	* *	* *	32	* *	* *	* *	0	* *	* *		0	* *
Pulaski Ct	533	8.6 (38)	6.8 (43)	1,818	8.9(11)	4.7(95)	12.2(13)	741	15.8(40)	13.0(28)		1,131	6.5 (35)
Radcliffe	539	10.0 (62)	7.1 (55)	2,110	6.3(52)	2.5(19)	14.9(47)	1,005	23.0(13)	16.5(72)		1,585	7.2 (44)
Richmond	415	9.2 (50)	9.2 (91)	1,308	8.9(11)	6.0(12)	13.9(29)	516	20.9(11)	20.2(10)		901	12.4(100)
Robertson	25	* *	* *	108	13.9(15)	0.9 (2)	18.5(10)	53	* *	* *		85	* *
Rockcastle	183	8.7 (41)	6.6 (35)	611	6.5(59)	7.5(13)	17.3(93)	279	16.8(49)	12.5(22)		416	20.7(138)
Rowan Cty	177	7.3 (17)	15.3(114)	681	3.7(12)	3.4(41)	15.0(49)	298	18.5(78)	13.1(31)		445	16.9(130)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2002 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% >95th (Rank)	Number
Russell Ct	157	8.9 (45)	8.9 (83)	580	8.6(10)	2.1(13)	17.1(91)	241	19.1(94)	20.7(11)	368	10.1 (78)
Scott Cty	230	8.7 (41)	7.0 (51)	684	8.8(10)	5.7(11)	17.5(96)	256	11.3 (8)	12.1(19)	464	11.4 (88)
Shelby Cty	231	10.0 (62)	5.6 (19)	738	22.1(15)	1.9 (9)	37.1(15)	265	21.9(12)	32.8(14)	515	13.6(110)
Siler	5	* *	* *	44	* *	* *	* *	25	* *	* *	40	* *
Simpson Ct	124	12.9(108)	4.8 (10)	426	9.6(12)	2.8(23)	16.4(77)	165	29.1(14)	18.2(90)	275	22.2(144)
South Shor	34	* *	* *	156	3.2 (7)	7.7(14)	9.0 (3)	75	* *	* *	127	5.5 (22)
Spencer Ct	69	* *	* *	258	10.9(13)	3.1(30)	18.2(10)	112	17.0(56)	16.1(66)	188	10.6 (84)
TAPP-South	27	* *	* *	53	* *	* *	* *	4	* *	* *	30	* *
TAPP-West	26	* *	* *	65	* *	* *	* *	8	* *	* *	34	* *
Taylor Cty	195	3.6 (2)	8.7 (77)	759	6.6(61)	2.8(23)	16.9(88)	356	21.3(12)	17.4(82)	492	4.7 (14)
Todd Cty H	101	5.9 (6)	6.9 (45)	448	8.0(91)	4.0(73)	16.5(82)	205	14.1(22)	8.8 (6)	351	17.4(131)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2002 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic												(3,4)
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Trigg Cty	75	* *	* *	290	5.9(44)	4.5(91)	13.1(20)	131	16.8(49)	12.2(21)	199	7.5 (48)
Trimble Ct	57	* *	* *	232	5.6(38)	8.6(14)	7.3 (2)	111	10.8 (7)	9.0 (7)	184	4.3 (11)
Union Cty	119	10.9 (80)	2.5 (1)	474	12.0(14)	2.5(19)	16.9(88)	205	17.6(69)	11.7(13)	369	7.6 (50)
Wallins	28	* *	* *	109	3.7(12)	8.3(14)	19.3(11)	47	* *	* *	55	* *
Washington	88	* *	* *	369	9.8(12)	3.5(47)	19.8(12)	165	18.8(85)	12.1(19)	283	12.4(100)
Wayne Cty	216	12.0 (98)	7.4 (61)	844	4.3(21)	3.0(29)	16.8(86)	402	17.9(73)	21.4(11)	575	9.6 (74)
Webster Ct	113	18.6(114)	5.3 (15)	414	10.6(13)	3.9(71)	16.4(77)	177	12.4(14)	16.9(78)	304	15.1(121)
Wheelbrigh	62	* *	* *	279	5.7(41)	5.0(10)	15.8(60)	151	17.2(60)	19.2(10)	191	13.6(110)
Williamsbu	213	10.3 (67)	12.2(109)	918	9.9(13)	3.8(63)	21.0(13)	466	23.2(13)	23.4(13)	721	6.8 (38)
Wolfe Cty	100	8.0 (27)	7.0 (51)	431	10.9(13)	0.5 (1)	25.3(15)	218	19.7(10)	23.4(13)	340	8.5 (63)
Woodford C	132	10.6 (73)	9.8 (96)	515	7.8(87)	3.7(57)	16.1(66)	205	14.6(28)	11.7(13)	349	12.0 (96)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2002 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic											(3,4)	
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)			(8)		
	Number	%	%	Number	%	%	%	Number	%	%	Number	%
	<2500 g (Rank)	>4000 g (Rank)		<5th (Rank)	<5th (Rank)	≥95th (Rank)		85th- (Rank)	<95th (Rank)	>95th (Rank)		(Rank)
Kentucky	32,542	10.0	7.5	114,943	7.6	4.6	16.8	50,285	17.7	16.8	79,270	11.6
Nation (Prior Year)	1,260,117	9.0	7.9	4,846,643	6.4	5.4	13.1	2,031,241	15.3	13.4	3,265,122	13.3

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2002 Pediatric Nutrition Surveillance (1)
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)						
Clinic	Breastfeeding (6)					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Adair Coun	135	11.1 (74)	49	* *	98	* *
Allen Coun	138	24.6 (15)	51	* *	95	* *
Anderson C	82	* *	43	* *	57	* *
Ashland-Bo	319	13.2 (65)	96	* *	180	0.0 (1)
Ballard Co	62	* *	26	* *	58	* *
Bar-Riv-Wa	752	15.8 (48)	231	2.2 (7)	389	0.0 (1)
Barren Cty	252	15.1 (55)	104	1.9 (10)	188	0.0 (1)
Bath Count	100	8.0(104)	35	* *	79	* *
Belfry	115	16.5 (39)	46	* *	83	* *
Bell Cty M	188	8.5 (99)	82	* *	153	0.0 (1)
Bell Cty P	155	13.5 (62)	70	* *	149	0.0 (1)
Berea	109	22.9 (18)	40	* *	81	* *
Betsy Lane	81	* *	36	* *	72	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2002 Pediatric Nutrition Surveillance (1)
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)						
Clinic	Breastfeeding (6)					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Bledsoe	23	* *	6	* *	11	* *
Boone Coun	487	12.7 (69)	235	3.4 (3)	331	0.0 (1)
Bourbon Ct	119	17.6 (31)	63	* *	108	0.0 (1)
Boyle Coun	187	9.1 (94)	53	* *	104	0.0 (1)
Bracken Co	70	* *	20	* *	34	* *
Breathitt	154	9.7 (88)	52	* *	103	0.0 (1)
Breckinrid	114	16.7 (36)	41	* *	87	* *
Bridges	162	10.5 (83)	53	* *	102	0.0 (1)
Bullitt Co	374	16.6 (38)	102	0.0 (29)	213	0.0 (1)
Butler Cty	115	9.6 (92)	41	* *	80	* *
C & Y	391	15.3 (54)	107	0.9 (23)	200	0.0 (1)
Caldwell C	108	8.3(100)	55	* *	91	* *
Calloway C	243	28.4 (11)	94	* *	173	0.0 (1)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2002 Pediatric Nutrition Surveillance (1)
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)						
Clinic	Breastfeeding (6)					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Campbell C	404	7.7(106)	160	1.3 (20)	237	0.0 (1)
Cannonsbur	106	17.0 (33)	31	* *	58	* *
Carlisle C	35	* *	21	* *	38	* *
Carroll Ct	107	14.0 (59)	40	* *	82	* *
Carter Cty	176	18.8 (25)	58	* *	112	0.0 (1)
Casey Cty	136	16.2 (44)	43	* *	85	* *
Christian	568	10.2 (85)	231	1.7 (12)	278	0.0 (1)
Clark Coun	255	14.1 (58)	94	* *	140	0.0 (1)
Clay Cty H	212	9.0 (97)	83	* *	182	0.0 (1)
Clinton Ct	111	5.4(111)	41	* *	87	* *
Corbin	243	15.6 (52)	126	0.0 (29)	205	0.0 (1)
Crittenden	62	* *	22	* *	46	* *
Cumberland	66	* *	30	* *	55	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2002 Pediatric Nutrition Surveillance (1)
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)						
Clinic	Breastfeeding (6)					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Daviess Co	652	13.5 (62)	220	1.4 (19)	362	0.0 (1)
Dixie	511	13.3 (64)	172	0.6 (26)	264	0.0 (1)
Edmonson C	102	19.6 (23)	38	* *	44	* *
Elliott Ct	56	* *	30	* *	48	* *
Estill Cty	158	17.1 (32)	45	* *	91	* *
Etown	387	14.7 (56)	122	0.8 (25)	188	0.0 (1)
Evarts	71	* *	36	* *	76	* *
Fairdale	331	15.7 (51)	101	2.0 (8)	153	0.0 (1)
Fleming Co	127	11.0 (75)	58	* *	102	0.0 (1)
Floyd Cty	297	4.4(112)	108	0.0 (29)	202	0.0 (1)
Foust Hlth	37	* *	15	* *	38	* *
Franklin C	308	26.3 (12)	131	2.3 (5)	164	0.0 (1)
Ft Campbel	719	30.3 (8)	276	6.2 (2)	368	0.0 (1)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2002 Pediatric Nutrition Surveillance (1)
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)						
Clinic	Breastfeeding (6)					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Fulton	53	* *	25	* *	36	* *
Gallatin C	74	* *	20	* *	40	* *
Garrard Ct	124	17.7 (30)	50	* *	114	0.0 (1)
Goose Rock	35	* *	12	* *	25	* *
Grant Cty	210	9.0 (97)	116	0.0 (29)	196	0.0 (1)
Graves Cty	314	22.3 (21)	95	* *	215	0.0 (1)
Grayson Ct	203	29.6 (9)	67	* *	150	0.0 (1)
Green Cty	93	* *	29	* *	61	* *
Greenup	215	18.1 (27)	82	* *	115	0.0 (1)
H & Env.	21	* *	0	* *	20	* *
Hancock Ct	58	* *	30	* *	35	* *
Harlan Cty	168	7.7(106)	75	* *	128	0.0 (1)
Harrison C	124	16.9 (34)	58	* *	99	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2002 Pediatric Nutrition Surveillance (1)
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)						
Clinic	Breastfeeding (6)					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Hart Cty H	184	10.9 (78)	53	* *	91	* *
Henderson	362	10.2 (85)	115	0.0 (29)	156	0.0 (1)
Henry Coun	86	* *	38	* *	73	* *
Hickman	42	* *	15	* *	28	* *
Hickman Ct	36	* *	13	* *	26	* *
Highview	330	38.5 (3)	130	1.5 (14)	190	0.0 (1)
Hindman	142	4.2(113)	65	* *	101	0.0 (1)
Hopkins Ct	353	11.0 (75)	125	0.0 (29)	222	0.0 (1)
Iroquois	588	26.2 (13)	196	1.5 (14)	297	0.0 (1)
Jackson Ct	131	10.7 (82)	62	* *	140	0.0 (1)
Jessamine	264	16.7 (36)	110	1.8 (11)	166	0.0 (1)
Johnson Co	233	8.2(101)	101	0.0 (29)	183	0.0 (1)
JT Arnett	10	* *	7	* *	16	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2002 Pediatric Nutrition Surveillance (1)
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)						
Clinic	Breastfeeding (6)					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Kenton Cty	490	8.2(101)	247	1.2 (21)	344	0.0 (1)
Knox Cty	345	8.1(103)	136	0.0 (29)	257	0.0 (1)
L & N	495	9.7 (88)	135	1.5 (14)	195	0.0 (1)
Larue Cty	111	21.6 (22)	54	* *	73	* *
Laurel Cty	481	10.4 (84)	163	1.2 (21)	262	0.0 (1)
Lawrence C	154	9.7 (88)	59	* *	114	0.0 (1)
Lee Cty He	64	* *	32	* *	59	* *
Leslie Cty	128	10.9 (78)	51	* *	97	* *
Letcher Ct	209	12.0 (71)	97	* *	119	0.0 (1)
Lewis Cty	125	16.8 (35)	53	* *	104	0.0 (1)
Lex-Fayett	750	43.1 (1)	224	6.3 (1)	404	0.0 (1)
Lex-Fayett	152	29.6 (9)	72	* *	128	0.0 (1)
Lex-Fayett	174	37.4 (4)	61	* *	106	0.0 (1)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2002 Pediatric Nutrition Surveillance (1)
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)						
Clinic	Breastfeeding (6)					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Lex-Fayette	239	33.9 (6)	94	* *	158	0.0 (1)
Lex-Fayette	245	40.0 (2)	74	* *	136	0.0 (1)
Liberty	34	* *	3	* *	2	* *
Lincoln Ct	227	13.7 (61)	90	* *	126	0.0 (1)
Livingston	44	* *	19	* *	41	* *
Logan Cty	202	3.5(114)	80	* *	144	0.0 (1)
Lynch	81	* *	29	* *	46	* *
Lyon Cty	38	* *	20	* *	28	* *
Magoffin C	137	2.9(115)	73	* *	117	0.0 (1)
Marion Cty	159	16.4 (41)	57	* *	106	0.0 (1)
Marshall C	196	22.4 (20)	74	* *	129	0.0 (1)
Marshall C	34	* *	14	* *	24	* *
Martin Cty	153	2.6(116)	50	* *	121	0.0 (1)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2002 Pediatric Nutrition Surveillance (1)
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)						
Clinic	Breastfeeding (6)					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Mason Cty	169	13.0 (67)	57	* *	108	0.0 (1)
McCracken	538	13.9 (60)	196	0.0 (29)	385	0.0 (1)
McCreary C	187	9.1 (94)	92	* *	165	0.0 (1)
McLean Ct	99	* *	42	* *	77	* *
Meade Cty	172	25.6 (14)	59	* *	103	0.0 (1)
Menifee Ct	62	* *	21	* *	46	* *
Mercer Cty	165	10.9 (78)	54	* *	89	* *
Metcalfe C	92	* *	43	* *	57	* *
Middletown	322	32.3 (7)	97	* *	166	0.0 (1)
Monroe Cnt	100	16.0 (45)	48	* *	59	* *
Montgomery	231	11.7 (73)	84	* *	126	0.0 (1)
Morgan Cty	109	11.0 (75)	61	* *	104	0.0 (1)
Muhlenberg	256	9.4 (93)	99	* *	195	0.0 (1)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2002 Pediatric Nutrition Surveillance (1)
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)						
Clinic	Breastfeeding (6)					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Neighborho	330	23.3 (17)	125	2.4 (4)	195	0.0 (1)
Nelson Cty	271	15.9 (47)	104	0.0 (29)	148	0.0 (1)
Newburg	387	15.8 (48)	133	2.3 (5)	214	0.0 (1)
Nicholas C	78	* *	27	* *	39	* *
Oasis Hlth	33	* *	16	* *	33	* *
Ohio Cty	210	18.1 (27)	78	* *	131	0.0 (1)
Oldham Cty	150	24.0 (16)	61	* *	116	0.0 (1)
Olive Hill	125	16.0 (45)	51	* *	98	* *
Oneida Ele	11	* *	7	* *	10	* *
Owen Cty	86	* *	26	* *	40	* *
Owsley Cty	51	* *	36	* *	54	* *
Park Duval	429	15.6 (52)	130	1.5 (14)	205	0.0 (1)
Pendleton	102	22.5 (19)	48	* *	86	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2002 Pediatric Nutrition Surveillance (1)
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)						
Clinic	Breastfeeding (6)					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Perry Cty	267	7.5(108)	112	0.9 (23)	201	0.0 (1)
Phelps	42	* *	14	* *	15	* *
Pike Cty	399	12.8 (68)	154	0.6 (26)	232	0.0 (1)
Portland	363	13.2 (65)	130	1.5 (14)	188	0.0 (1)
Powell Cty	173	6.9(109)	61	* *	112	0.0 (1)
Prenatal	94	* *	2	* *	0	* *
Prenatal	32	* *	0	* *	0	* *
Pulaski Ct	537	19.2 (24)	229	1.7 (12)	352	0.0 (1)
Radcliffe	539	17.8 (29)	221	0.5 (28)	307	0.0 (1)
Richmond	416	14.4 (57)	153	2.0 (8)	214	0.0 (1)
Robertson	25	* *	16	* *	24	* *
Rockcastle	184	12.0 (71)	50	* *	84	* *
Rowan Cty	177	7.9(105)	60	* *	105	0.0 (1)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2002 Pediatric Nutrition Surveillance (1)
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)						
Clinic	Breastfeeding (6)					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Russell Ct	158	15.8 (48)	62	* *	124	0.0 (1)
Scott Cty	230	16.5 (39)	77	* *	131	0.0 (1)
Shelby Cty	231	9.1 (94)	85	* *	155	0.0 (1)
Siler	5	* *	13	* *	9	* *
Simpson Ct	129	16.3 (43)	36	* *	81	* *
South Shor	35	* *	21	* *	28	* *
Spencer Ct	69	* *	31	* *	57	* *
TAPP-South	29	* *	12	* *	15	* *
TAPP-West	27	* *	11	* *	18	* *
Taylor Cty	195	16.4 (41)	73	* *	137	0.0 (1)
Todd Cty H	101	18.8 (25)	46	* *	97	* *
Trigg Cty	75	* *	31	* *	50	* *
Trimble Ct	57	* *	21	* *	48	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2002 Pediatric Nutrition Surveillance (1)
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)						
Clinic	Breastfeeding (6)					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Union Cty	120	12.5 (70)	55	* *	88	* *
Wallins	28	* *	15	* *	27	* *
Washington	88	* *	33	* *	72	* *
Wayne Cty	223	9.9 (87)	61	* *	135	0.0 (1)
Webster Ct	113	9.7 (88)	48	* *	58	* *
Wheelbrigh	62	* *	33	* *	48	* *
Williamsbu	213	10.8 (81)	95	* *	141	0.0 (1)
Wolfe Cty	100	6.0(110)	53	* *	85	* *
Woodford C	132	34.8 (5)	63	* *	112	0.0 (1)
Unknown Clinics	1,175	1.9(117)	4	* *	20	* *
Kentucky	32,800	15.6	11,889	1.3	20,006	0.0
Nation (Prior Year)	874,598	51.1	493,552	20.3	484,228	13.3

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2000 - 2002 Pediatric Nutrition Surveillance⁽¹⁾
Kentucky
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Growth and Anemia Indicators by Clinic												(3,4)
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Adair Coun	423	7.6 (28)	7.8 (70)	1,453	4.5(43)	7.0(12)	11.1(11)	627	11.5(12)	11.6(21)	871	6.0 (20)
Allen Coun	426	12.2(141)	10.1(129)	1,747	6.6(10)	4.2(45)	14.9(83)	809	19.9(13)	15.9(95)	1,125	10.0 (76)
Anderson C	298	7.4 (23)	8.7(100)	1,072	7.5(13)	5.0(77)	17.2(12)	476	21.0(14)	18.7(12)	812	7.1 (36)
Ashland-Bo	909	10.2(100)	8.5 (91)	3,088	7.7(13)	9.6(15)	13.4(48)	1,319	15.0(46)	13.7(56)	2,019	11.9 (95)
Ballard Co	199	9.0 (59)	8.5 (91)	766	7.2(13)	6.8(12)	13.7(60)	354	13.0(22)	11.3(18)	587	6.3 (26)
Bar-Riv-Wa	2,130	10.5(105)	7.7 (65)	6,382	5.8(89)	5.1(81)	13.2(46)	2,327	17.0(91)	12.9(39)	3,821	13.6(114)
Barren Cty	856	8.8 (51)	8.8(105)	2,956	5.6(81)	4.7(68)	15.1(87)	1,209	16.1(68)	15.9(95)	1,909	9.2 (66)
Bath Count	312	5.8 (7)	14.1(153)	1,346	9.1(15)	4.3(51)	19.2(14)	658	16.6(79)	17.5(11)	855	10.2 (79)
Belfry	394	8.4 (40)	7.9 (73)	1,315	1.8 (3)	9.4(15)	12.7(37)	570	14.0(32)	11.9(27)	869	8.6 (60)
Bell Cty M	558	10.0 (90)	7.7 (65)	2,428	5.0(60)	3.4(21)	18.5(13)	1,261	18.0(11)	18.2(12)	1,803	2.6 (2)
Bell Cty P	511	9.2 (64)	8.6 (94)	2,331	4.7(52)	4.3(51)	15.0(84)	1,172	12.4(18)	12.3(33)	1,586	12.0 (96)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2000 - 2002 Pediatric Nutrition Surveillance⁽¹⁾
Kentucky
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years⁽²⁾

Comparison of Growth and Anemia Indicators by Clinic												(3,4)
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Berea	340	7.9 (34)	11.8(147)	1,374	5.8(89)	4.9(75)	11.9(23)	597	20.4(14)	14.4(66)	1,026	10.5 (82)
Betsy Lane	262	6.5 (11)	10.7(136)	1,237	1.9 (4)	11.9(16)	11.5(19)	635	8.3 (2)	10.9(14)	922	8.4 (56)
Bledsoe	66	* *	* *	279	2.9 (9)	11.5(15)	14.0(68)	151	16.6(79)	16.6(10)	172	8.7 (62)
Boone Coun	1,458	8.9 (54)	9.4(121)	5,209	6.8(12)	5.1(81)	14.4(75)	2,157	17.2(97)	13.6(51)	3,710	13.3(108)
Bourbon Ct	434	11.5(129)	9.0(109)	1,451	5.7(84)	2.8(11)	17.8(12)	604	21.7(15)	17.4(11)	1,024	29.7(156)
Boyle Coun	535	11.6(132)	5.6 (13)	1,635	5.7(84)	5.5(93)	13.7(60)	636	15.7(60)	15.4(81)	1,001	9.5 (70)
Bracken Co	197	12.2(141)	7.1 (46)	646	7.3(13)	7.0(12)	19.2(14)	264	17.0(91)	22.0(14)	431	6.0 (20)
Breathitt	448	9.2 (64)	8.7(100)	2,110	4.3(34)	4.0(38)	16.7(11)	1,188	13.0(22)	14.7(72)	1,712	6.9 (33)
Breckinrid	347	8.4 (40)	10.1(129)	1,415	6.6(10)	3.7(28)	13.1(44)	642	17.3(99)	11.2(16)	1,053	3.9 (5)
Bridges	474	12.7(146)	5.3 (7)	1,226	10.0(15)	4.6(65)	16.2(10)	418	15.8(63)	12.9(39)	787	8.3 (54)
Bullitt Co	1,071	8.1 (36)	7.3 (50)	3,130	6.4(10)	3.9(35)	14.5(77)	1,174	16.7(84)	14.0(60)	1,816	19.4(144)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2000 - 2002 Pediatric Nutrition Surveillance⁽¹⁾
Kentucky
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years⁽²⁾

Comparison of Growth and Anemia Indicators by Clinic												(3,4)
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)			(8)	
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number
Butler Cty	348	8.6 (47)	8.0 (78)	1,332	4.1(31)	4.2(45)	17.9(13)	598	17.7(11)	21.4(14)	894	5.9 (19)
C & Y	419	10.0 (90)	5.5 (11)	1,204	8.2(14)	5.6(95)	16.1(10)	493	19.1(12)	20.5(14)	302	20.9(148)
Caldwell C	320	5.9 (8)	8.8(105)	1,171	6.7(11)	8.3(14)	13.9(66)	485	14.4(36)	15.1(77)	862	15.4(130)
Caldwell E	1	* *	* *	35	* *	* *	* *	23	* *	* *	34	* *
Calloway C	725	8.1 (36)	10.9(139)	2,655	4.6(45)	4.3(51)	13.4(48)	1,134	15.3(51)	15.5(87)	1,706	16.9(136)
Campbell C	1,082	10.4(104)	7.6 (61)	3,601	6.2(98)	4.1(40)	15.7(98)	1,523	19.6(13)	19.6(13)	2,404	13.3(108)
Cannonsbur	303	10.6(107)	8.6 (94)	1,044	3.6(22)	9.7(15)	9.7 (7)	473	14.6(41)	14.0(60)	697	14.8(124)
Carlisle C	133	6.8 (14)	12.0(149)	581	5.0(60)	4.1(40)	20.7(15)	295	19.3(13)	26.1(15)	407	21.9(149)
Carroll Ct	354	9.6 (79)	6.5 (31)	1,216	9.7(15)	6.7(12)	16.2(10)	505	17.6(10)	17.8(12)	882	14.6(122)
Carter Cty	535	9.0 (59)	11.4(145)	2,024	3.2(15)	9.9(15)	9.6 (6)	915	9.1 (4)	8.7 (6)	1,391	8.9 (64)
Casey Cty	379	9.8 (84)	5.8 (18)	1,576	5.5(79)	5.6(95)	12.1(27)	737	16.4(74)	13.0(45)	988	7.5 (44)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2000 - 2002 Pediatric Nutrition Surveillance⁽¹⁾
Kentucky
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Growth and Anemia Indicators by Clinic												(3,4)	
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Christian	1,649	11.2 (125)	6.2 (29)	6,001	7.4(13)	6.0(10)	15.5(96)	2,371	14.4(36)	12.9(39)		4,013	31.9 (157)
Clark Coun	744	11.8 (135)	8.2 (83)	2,430	8.1(14)	3.4(21)	16.1(10)	964	19.9(13)	15.4(81)		1,658	12.0 (96)
Clay Cty H	668	12.1 (140)	6.7 (38)	2,697	5.2(72)	7.5(13)	17.2(12)	1,306	17.7(11)	17.1(11)		1,936	8.6 (60)
Clinton Ct	352	9.1 (61)	8.8(105)	1,443	4.0(30)	6.2(11)	15.0(84)	666	12.3(17)	11.9(27)		972	6.2 (24)
Corbin	846	9.2 (64)	9.0(109)	3,328	4.8(53)	4.1(40)	14.8(79)	1,488	15.5(55)	14.9(75)		2,432	6.0 (20)
Crittenden	197	5.6 (5)	10.7(136)	803	5.1(67)	6.4(11)	12.1(27)	362	14.4(36)	9.7 (9)		593	12.6(103)
Cumberland	211	3.8 (1)	9.0(109)	884	3.7(25)	6.3(11)	13.0(42)	411	15.3(51)	14.6(71)		444	14.9(126)
Daviess Co	2,109	8.9 (54)	5.8 (18)	6,665	8.0(13)	2.2 (5)	17.7(12)	2,616	16.6(79)	16.1(10)		4,608	11.0 (88)
Dixie	1,678	10.1 (96)	8.0 (78)	4,322	6.5(10)	5.6(95)	12.6(36)	1,532	13.6(28)	12.3(33)		2,570	12.7(105)
Edmonson C	312	9.9 (86)	8.3 (86)	1,098	3.6(22)	6.1(10)	11.7(22)	449	14.7(42)	13.8(57)		707	5.2 (10)
Elliott Ct	243	6.6 (13)	12.3(151)	959	2.8 (7)	6.0(10)	12.3(30)	458	14.4(36)	14.2(63)		671	5.7 (16)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2000 - 2002 Pediatric Nutrition Surveillance⁽¹⁾
Kentucky
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years⁽²⁾

Comparison of Growth and Anemia Indicators by Clinic												(3,4)
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Estill Cty	441	9.3 (71)	10.0 (127)	1,820	2.8 (7)	5.7 (98)	13.4 (48)	873	17.3 (99)	16.4 (10)	1,380	7.3 (39)
Etown	1,108	9.6 (79)	6.9 (41)	3,740	7.1 (12)	3.6 (25)	13.4 (48)	1,509	16.7 (84)	13.6 (51)	2,680	7.3 (39)
Evarts	237	8.4 (40)	11.0 (140)	1,095	3.7 (25)	9.3 (15)	18.4 (13)	558	15.2 (48)	19.5 (13)	877	8.4 (56)
Fairdale	983	9.6 (79)	8.4 (88)	2,571	4.6 (45)	6.1 (10)	11.2 (13)	933	14.9 (45)	10.9 (14)	1,653	6.3 (26)
Fleming Co	410	7.6 (28)	9.8 (126)	1,817	4.6 (45)	6.5 (12)	13.3 (47)	862	15.7 (60)	13.6 (51)	1,415	16.5 (135)
Floyd Cty	880	8.8 (51)	7.8 (70)	3,193	3.8 (27)	11.5 (15)	12.0 (25)	1,423	9.1 (4)	11.2 (16)	2,356	8.4 (56)
Foust Hlth	130	13.1 (148)	8.5 (91)	671	5.2 (72)	3.6 (25)	17.9 (13)	379	15.6 (58)	16.1 (10)	559	26.7 (155)
Franklin C	969	7.6 (28)	6.9 (41)	3,034	8.0 (13)	5.7 (98)	17.7 (12)	1,135	18.2 (11)	18.4 (12)	2,092	11.5 (90)
Ft Campbel	2,388	7.4 (23)	11.1 (141)	9,039	3.5 (19)	8.6 (14)	6.4 (2)	4,027	12.0 (16)	6.9 (2)	6,129	17.1 (138)
Fulton	181	11.0 (117)	5.5 (11)	601	7.0 (12)	4.8 (71)	19.6 (14)	255	15.7 (60)	20.4 (14)	391	10.0 (76)
Gallatin C	216	10.6 (107)	7.4 (56)	796	5.7 (84)	5.0 (77)	12.9 (40)	360	13.1 (25)	14.4 (66)	536	13.4 (111)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2000 - 2002 Pediatric Nutrition Surveillance⁽¹⁾
Kentucky
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Growth and Anemia Indicators by Clinic												(3,4)
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Garrard Ct	343	10.5 (105)	9.0 (109)	1,536	5.7(84)	3.7(28)	14.8(79)	749	17.1 (94)	17.9(12)	1,005	14.8 (124)
Goose Rock	94	* *	* *	494	1.4 (1)	13.2(16)	11.3(15)	286	6.6 (1)	8.4 (5)	424	6.1 (23)
Grant Cty	667	11.1 (120)	7.9 (73)	2,986	4.3(34)	7.3(13)	9.3 (5)	1,439	8.8 (3)	7.0 (3)	2,350	6.2 (24)
Graves Cty	896	9.9 (86)	9.4 (121)	3,377	5.7(84)	6.1(10)	14.8(79)	1,553	18.2(11)	15.3(80)	2,261	16.4 (134)
Grayson Ct	680	7.4 (23)	8.2 (83)	2,632	10.0(15)	3.1(16)	18.8(14)	1,165	16.6(79)	16.5(10)	1,966	3.0 (3)
Green Cty	270	8.5 (46)	9.6 (124)	956	4.6(45)	3.6(25)	17.7(12)	417	21.8(15)	17.7(11)	602	5.8 (17)
Greenup	734	9.9 (86)	8.3 (86)	2,659	2.9 (9)	8.9(14)	11.9(23)	1,167	15.5(55)	12.8(38)	1,806	8.2 (52)
H & Env.	567	9.7 (82)	4.8 (4)	1,721	4.9(56)	7.1(13)	12.3(30)	619	17.3(99)	14.5(69)	1,182	16.1 (133)
Hancock Ct	194	10.3 (102)	4.1 (2)	766	5.0(60)	4.2(45)	16.2(10)	349	14.3(34)	16.6(10)	583	7.2 (38)
Harlan Cty	558	8.4 (40)	12.0 (149)	2,500	3.2(15)	8.7(14)	17.1(12)	1,331	17.5(10)	18.0(12)	1,691	9.5 (70)
Harrison C	428	12.9 (147)	7.2 (48)	1,472	3.5(19)	6.7(12)	9.9 (9)	635	10.2 (7)	7.9 (4)	1,038	12.0 (96)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2000 - 2002 Pediatric Nutrition Surveillance⁽¹⁾
Kentucky
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Growth and Anemia Indicators by Clinic												(3,4)
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Hart Cty H	500	7.0 (15)	9.2 (118)	1,832	4.6(45)	4.4(58)	14.3(74)	807	15.2(48)	15.7(90)	1,236	7.3 (39)
Henderson	1,032	11.1 (120)	5.1 (5)	3,357	6.4(10)	6.1(10)	13.9(66)	1,307	20.0(14)	19.0(13)	1,249	7.7 (47)
Henry Coun	349	7.2 (19)	6.0 (24)	1,147	11.7(16)	4.7(68)	22.8(15)	445	19.3(13)	23.4(15)	830	15.3(127)
Hickman	110	11.8(135)	9.1(114)	497	6.6(10)	3.2(17)	20.3(15)	266	18.4(12)	20.3(14)	371	19.1(141)
Hickman Ct	99	* *	* *	394	4.3(34)	5.8(10)	15.2(89)	179	15.6(58)	20.1(13)	305	7.5 (44)
Highview	1,015	8.0 (35)	7.6 (61)	3,306	5.6(81)	5.1(81)	13.8(63)	1,353	13.7(29)	10.0(10)	2,101	13.7(116)
Hindman	459	11.5(129)	6.1 (26)	2,169	8.8(15)	6.0(10)	18.9(14)	1,134	11.7(15)	11.4(20)	1,789	8.8 (63)
Hopkins Ct	1,156	10.8(112)	5.4 (8)	4,279	9.9(15)	2.5 (7)	19.9(15)	1,882	19.3(13)	18.5(12)	3,086	13.8(117)
Iroquois	1,802	10.1 (96)	7.5 (58)	5,345	5.1(67)	4.9(75)	13.4(48)	2,048	16.3(71)	15.5(87)	3,445	15.6(131)
Jackson Ct	455	8.6 (47)	9.7(125)	1,823	5.9(92)	4.1(40)	15.8(10)	848	13.0(22)	15.8(92)	1,405	3.3 (4)
Jessamine	890	7.0 (15)	9.0(109)	3,026	6.6(10)	3.2(17)	15.3(94)	1,281	17.6(10)	15.4(81)	2,070	10.7 (84)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2000 - 2002 Pediatric Nutrition Surveillance⁽¹⁾
Kentucky
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Growth and Anemia Indicators by Clinic												(3,4)
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Johnson Co	754	7.0 (15)	10.3 (132)	3,002	3.6(22)	8.1(14)	15.0(84)	1,403	13.5(27)	14.4(66)	2,208	17.0 (137)
JT Arnett	28	* *	* *	159	4.4(41)	3.8(31)	18.9(14)	83	* *	* *	131	9.2 (66)
Kenton Cty	1,479	12.2 (141)	7.3 (50)	5,025	6.4(10)	4.6(65)	15.9(10)	2,054	20.4(14)	20.2(13)	3,571	11.8 (93)
Knox Cty	1,031	9.3 (71)	8.2 (83)	4,554	5.2(72)	3.9(35)	16.0(10)	2,244	16.5(76)	16.8(11)	3,528	14.1 (119)
L & N	1,137	9.2 (64)	7.3 (50)	3,009	6.7(11)	7.0(12)	12.3(30)	1,042	17.1(94)	15.0(76)	1,780	14.7 (123)
Larue Cty	343	8.7 (50)	6.7 (38)	1,173	7.7(13)	5.3(86)	16.5(11)	510	17.5(10)	14.7(72)	896	10.9 (87)
Laurel Cty	1,513	11.8 (135)	9.2 (118)	4,850	5.0(60)	6.5(12)	13.6(57)	1,890	19.6(13)	15.9(95)	3,008	13.3 (108)
Lawrence C	484	11.8 (135)	7.0 (44)	1,732	4.2(32)	8.0(14)	16.6(11)	767	15.8(63)	16.2(10)	1,161	9.1 (65)
Lee Cty He	206	8.3 (39)	11.2 (142)	1,026	6.1(97)	1.7 (3)	22.6(15)	528	19.1(12)	21.0(14)	841	18.2 (140)
Leslie Cty	405	7.2 (19)	11.9 (148)	1,966	2.4 (6)	4.8(71)	18.3(13)	1,067	17.5(10)	19.3(13)	1,580	6.6 (31)
Letcher Ct	686	9.2 (64)	4.5 (3)	2,662	7.2(13)	4.1(40)	21.3(15)	1,202	16.8(87)	20.2(13)	1,998	4.0 (6)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2000 - 2002 Pediatric Nutrition Surveillance⁽¹⁾
Kentucky
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Growth and Anemia Indicators by Clinic												(3,4)
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Lewis Cty	405	11.1 (120)	9.1 (114)	1,428	6.0(94)	5.3(86)	13.6(57)	615	21.0(14)	17.4(11)	968	5.3 (12)
Lex-Fayette	713	14.3 (152)	8.4 (88)	2,200	4.3(34)	5.3(86)	12.2(29)	859	16.1(68)	12.9(39)	1,455	23.2(152)
Lex-Fayette	1,965	10.8 (112)	6.8 (40)	5,393	5.2(72)	4.8(71)	14.0(68)	1,935	19.1(12)	17.9(12)	3,162	4.7 (9)
Lex-Fayette	879	10.1 (96)	6.4 (30)	2,195	4.8(53)	6.2(11)	11.3(15)	632	15.3(51)	11.9(27)	1,247	37.5(158)
Lex-Fayette	591	7.8 (32)	10.0(127)	1,793	4.9(56)	5.2(85)	12.8(38)	710	15.2(48)	12.3(33)	1,119	22.7(151)
Lex-Fayette	710	10.0 (90)	6.5 (31)	2,239	5.2(72)	4.3(51)	12.5(35)	909	18.4(12)	12.9(39)	1,375	12.4(100)
Liberty	46	* *	* *	71	* *	* *	* *	18	* *	* *	29	* *
Lincoln Ct	671	10.7(111)	7.6 (61)	2,433	5.6(81)	6.1(10)	15.5(96)	1,078	19.6(13)	17.3(11)	1,826	7.8 (49)
Livingston	174	10.3(102)	7.5 (58)	712	5.1(67)	7.7(13)	16.6(11)	336	16.1(68)	15.8(92)	550	8.2 (52)
Logan Cty	646	9.8 (84)	5.4 (8)	2,059	4.6(45)	3.9(35)	15.3(94)	794	20.2(14)	15.9(95)	1,317	19.3(143)
Lynch	226	11.5(129)	6.6 (36)	971	6.3(10)	5.8(10)	15.7(98)	504	16.9(89)	18.7(12)	779	2.4 (1)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2000 - 2002 Pediatric Nutrition Surveillance⁽¹⁾
Kentucky
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Growth and Anemia Indicators by Clinic												(3,4)
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Lyon Cty	123	5.7 (6)	10.6(135)	535	6.7(11)	5.4(91)	18.5(13)	258	16.3(71)	16.7(11)	418	14.4(120)
Magoffin C	447	8.9 (54)	10.5(134)	2,042	7.0(12)	5.3(86)	19.9(15)	1,020	15.9(65)	16.6(10)	1,414	19.9(147)
Marion Cty	464	10.6(107)	7.5 (58)	1,980	8.7(15)	4.7(68)	18.8(14)	899	16.9(89)	15.1(77)	1,413	9.8 (74)
Marshall C	577	9.4 (75)	7.3 (50)	2,106	6.6(10)	7.3(13)	13.0(42)	896	9.3 (6)	6.5 (1)	1,511	6.6 (31)
Marshall C	83	* *	* *	331	6.0(94)	4.2(45)	18.4(13)	159	17.0(91)	16.4(10)	255	10.6 (83)
Martin Cty	475	11.8(135)	5.9 (21)	2,154	4.9(56)	5.3(86)	16.3(11)	1,118	18.2(11)	17.8(12)	1,729	6.5 (30)
Mason Cty	498	7.2 (19)	8.0 (78)	1,682	8.3(14)	7.3(13)	14.7(78)	714	14.3(34)	11.8(25)	1,164	13.5(112)
McCracken	1,594	11.1(120)	8.1 (82)	5,560	6.6(10)	5.4(91)	18.1(13)	2,408	16.5(76)	13.8(57)	4,131	19.1(141)
McCreary C	607	11.0(117)	10.7(136)	2,678	5.0(60)	5.8(10)	13.4(48)	1,314	12.9(20)	11.6(21)	2,077	7.1 (36)
McLean Ct	316	11.4(128)	5.7 (16)	1,312	7.9(13)	3.3(19)	18.9(14)	597	18.8(12)	22.1(15)	1,011	5.2 (10)
Meade Cty	557	8.6 (47)	7.9 (73)	2,035	8.1(14)	4.3(51)	15.9(10)	917	17.6(10)	15.7(90)	1,527	19.4(144)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2000 - 2002 Pediatric Nutrition Surveillance⁽¹⁾
Kentucky
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Growth and Anemia Indicators by Clinic												(3,4)
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Menifee Ct	206	9.2 (64)	7.3 (50)	895	4.8(53)	5.5(93)	11.4(18)	425	17.4(10)	13.2(47)	580	10.7 (84)
Mercer Cty	486	9.5 (77)	6.6 (36)	1,770	6.7(11)	3.8(31)	13.6(57)	786	18.2(11)	15.4(81)	1,299	8.4 (56)
Metcalfe C	291	12.4(144)	11.3(144)	1,037	3.9(29)	4.5(62)	9.8 (8)	436	16.5(76)	13.1(46)	668	9.6 (72)
Middletown	857	10.9(116)	7.0 (44)	2,392	7.1(12)	6.8(12)	13.7(60)	828	12.4(18)	13.4(48)	1,441	13.5(112)
Monroe Cnt	332	5.1 (3)	8.7(100)	1,230	6.6(10)	2.8(11)	12.3(30)	500	16.6(79)	12.2(31)	820	13.0(107)
Montgomery	679	9.9 (86)	11.2(142)	2,448	5.3(78)	4.2(45)	13.4(48)	1,047	15.9(65)	12.3(33)	1,813	12.0 (96)
Morgan Cty	343	5.2 (4)	10.2(131)	1,661	4.3(34)	4.5(62)	19.6(14)	838	18.1(11)	19.5(13)	1,332	19.4(144)
Muhlenberg	854	9.7 (82)	7.8 (70)	3,007	2.9 (9)	7.5(13)	11.2(13)	1,294	11.6(13)	10.7(12)	1,898	11.1 (89)
Neighborhood	1,179	10.8(112)	7.7 (65)	3,685	5.5(79)	6.5(12)	13.5(55)	1,474	19.5(13)	15.4(81)	2,570	5.8 (17)
Nelson Cty	834	11.0(117)	7.2 (48)	3,102	6.9(12)	4.6(65)	14.2(73)	1,336	16.4(74)	12.1(30)	2,078	9.8 (74)
Newburg	1,168	11.2(125)	7.1 (46)	3,343	6.0(94)	7.6(13)	12.0(25)	1,162	11.1(11)	10.0(10)	2,130	26.0(154)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2000 - 2002 Pediatric Nutrition Surveillance⁽¹⁾
Kentucky
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Growth and Anemia Indicators by Clinic												(3,4)
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Nicholas C	222	13.5 (151)	5.4 (8)	843	4.5(43)	10.0(15)	10.1(10)	379	10.6 (9)	8.7 (6)	645	13.8(117)
Oasis Hlth	101	14.9(153)	5.9 (21)	567	8.1(14)	1.2 (1)	15.9(10)	318	16.7(84)	11.3(18)	471	11.7 (92)
Ohio Cty	627	10.0 (90)	8.6 (94)	2,448	4.4(41)	3.0(14)	15.2(89)	1,156	16.8(87)	16.1(10)	1,876	10.0 (76)
Oldham Cty	437	10.8(112)	8.7(100)	1,597	8.2(14)	3.3(19)	15.2(89)	691	18.5(12)	14.5(69)	1,029	12.6(103)
Olive Hill	378	6.1 (9)	13.0(152)	1,644	3.8(27)	7.1(13)	13.8(63)	826	21.7(15)	15.1(77)	1,120	7.7 (47)
Oneida Ele	42	* *	* *	244	7.0(12)	2.9(13)	22.1(15)	159	14.5(40)	26.4(15)	209	22.5(150)
Owen Cty	231	7.8 (32)	5.6 (13)	792	5.9(92)	6.4(11)	14.0(68)	339	15.0(46)	13.6(51)	553	6.9 (33)
Owsley Cty	163	7.4 (23)	8.0 (78)	814	5.8(89)	2.7(10)	25.3(16)	436	21.3(15)	21.1(14)	669	15.8(132)
Park Duval	461	18.2(154)	6.1 (26)	1,522	11.9(16)	4.3(51)	15.2(89)	696	20.0(14)	15.5(87)	1,090	9.4 (69)
Pendleton	321	10.6(107)	5.9 (21)	1,127	6.2(98)	4.4(58)	12.4(34)	479	18.2(11)	13.4(48)	812	8.3 (54)
Perry Cty	844	11.3(127)	7.3 (50)	3,324	8.0(13)	2.2 (5)	29.9(16)	1,596	17.5(10)	22.6(15)	2,614	5.3 (12)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2000 - 2002 Pediatric Nutrition Surveillance⁽¹⁾
Kentucky
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Growth and Anemia Indicators by Clinic												(3,4)
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number
Phelps	106	7.5 (27)	5.7 (16)	437	4.6(45)	14.0(16)	15.1(87)	224	11.6(13)	13.8(57)	305	14.4(120)
Pike Cty	1,237	9.3 (71)	8.6 (94)	4,712	3.1(14)	8.6(14)	16.8(11)	2,112	20.9(14)	20.8(14)	3,220	6.3 (26)
Portland	1,210	11.6(132)	5.8 (18)	3,603	8.0(13)	6.4(11)	16.8(11)	1,371	15.5(55)	14.7(72)	2,500	12.4(100)
Powell Cty	471	10.0 (90)	7.9 (73)	1,899	3.3(17)	3.8(31)	13.1(44)	939	14.8(44)	13.6(51)	1,477	5.6 (15)
Prenatal	254	7.1 (18)	9.1(114)	199	2.0 (5)	1.5 (2)	33.7(16)	0	* *	* *	0	* *
Prenatal	152	4.6 (2)	9.2(118)	148	1.4 (1)	12.8(16)	2.7 (1)	0	* *	* *	0	* *
Pulaski Ct	1,589	8.4 (40)	7.6 (61)	5,397	5.2(72)	5.8(10)	11.5(19)	2,241	12.9(20)	12.2(31)	3,445	7.3 (39)
Radcliffe	1,686	8.4 (40)	7.7 (65)	6,827	5.0(60)	3.0(14)	13.8(63)	3,131	20.5(14)	15.8(92)	5,168	8.0 (51)
Richmond	1,286	10.0 (90)	10.3(132)	4,081	6.8(12)	8.5(14)	11.6(21)	1,598	18.1(11)	16.5(10)	2,820	12.4(100)
Robertson	69	* *	* *	298	8.7(15)	4.0(38)	17.4(12)	137	10.9(10)	11.7(24)	237	4.2 (8)
Rockcastle	490	6.3 (10)	8.8(105)	1,787	5.0(60)	6.4(11)	16.6(11)	797	16.3(71)	13.4(48)	1,278	15.3(127)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2000 - 2002 Pediatric Nutrition Surveillance⁽¹⁾
Kentucky
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years⁽²⁾

Comparison of Growth and Anemia Indicators by Clinic												(3,4)
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number
Rowan Cty	561	8.9 (54)	14.6 (154)	1,987	3.4(18)	5.0(77)	13.5(55)	836	16.0(67)	14.1(62)	1,269	17.2 (139)
Russell Ct	507	8.1 (36)	7.7 (65)	1,732	6.2(98)	4.3(51)	16.2(10)	687	18.9(12)	17.9(12)	1,029	7.4 (43)
Scott Cty	658	9.3 (71)	6.5 (31)	1,826	6.5(10)	6.1(10)	17.1(12)	696	14.7(42)	12.6(37)	1,236	11.8 (93)
Shelby Cty	696	9.1 (61)	8.6 (94)	2,138	14.6(16)	2.6 (8)	32.2(16)	783	22.0(15)	27.8(15)	1,486	11.6 (91)
Siler	13	* *	* *	49	* *	* *	* *	25	* *	* *	40	* *
Simpson Ct	419	9.5 (77)	7.4 (56)	1,261	7.1(12)	4.4(58)	14.4(75)	450	21.8(15)	16.0(99)	758	25.6 (153)
South Shor	46	* *	* *	170	2.9 (9)	8.2(14)	8.8 (4)	75	* *	* *	127	5.5 (14)
Spencer Ct	216	13.4 (150)	5.6 (13)	746	10.2(15)	3.8(31)	18.2(13)	296	17.2(97)	16.2(10)	531	9.2 (66)
TAPP-South	124	7.3 (22)	6.5 (31)	203	3.0(13)	9.9(15)	11.3(15)	18	* *	* *	80	* *
TAPP-West	138	10.1 (96)	5.1 (5)	254	4.3(34)	9.1(14)	6.7 (3)	27	* *	* *	94	* *
Taylor Cty	582	7.6 (28)	7.9 (73)	2,173	4.9(56)	5.0(77)	14.1(71)	978	17.1(94)	14.3(65)	1,366	4.1 (7)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2000 - 2002 Pediatric Nutrition Surveillance⁽¹⁾
Kentucky
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Growth and Anemia Indicators by Clinic												(3,4)
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Todd Cty H	333	10.2 (100)	6.0 (24)	1,323	6.7(11)	4.2(45)	14.1(71)	592	13.9(31)	8.8 (8)	1,005	15.3 (127)
Trigg Cty	231	12.6 (145)	6.5 (31)	953	5.1(67)	4.8(71)	12.9(40)	429	13.8(30)	10.7(12)	717	10.2 (79)
Trimble Ct	202	8.9 (54)	6.9 (41)	667	6.3(10)	7.2(13)	11.1(11)	286	10.5 (8)	12.9(39)	481	6.9 (33)
Union Cty	398	11.1 (120)	2.3 (1)	1,468	10.4(16)	3.7(28)	14.8(79)	662	18.7(12)	11.8(25)	1,092	7.5 (44)
Wallins	45	* *	* *	138	4.3(34)	9.4(15)	18.1(13)	54	* *	* *	65	* *
Washington	263	6.5 (11)	8.7(100)	1,080	6.5(10)	3.4(21)	15.2(89)	481	17.9(11)	11.6(21)	813	9.7 (73)
Wayne Cty	626	8.8 (51)	8.6 (94)	2,537	3.5(19)	4.5(62)	16.4(11)	1,201	18.1(11)	20.3(14)	1,690	10.2 (79)
Webster Ct	347	13.3 (149)	6.1 (26)	1,170	10.0(15)	2.6 (8)	19.4(14)	470	13.4(26)	19.4(13)	845	12.9(106)
Wheelbrigh	179	11.7 (134)	8.4 (88)	834	4.2(32)	4.4(58)	12.8(38)	454	15.4(54)	15.4(81)	515	10.7 (84)
Williamsbu	654	9.2 (64)	11.6 (146)	2,784	8.3(14)	3.5(24)	20.3(15)	1,369	20.4(14)	23.0(15)	2,191	6.3 (26)
Wolfe Cty	287	9.1 (61)	9.4(121)	1,308	8.0(13)	1.8 (4)	21.9(15)	670	21.2(15)	20.1(13)	1,051	7.8 (49)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2000 - 2002 Pediatric Nutrition Surveillance⁽¹⁾
Kentucky
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Growth and Anemia Indicators by Clinic												(3,4)
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number
Woodford C	417	9.4 (75)	9.1 (114)	1,442	5.1 (67)	5.1 (81)	15.8 (10)	593	14.2 (33)	14.2 (63)	941	13.6 (114)
Kentucky	100,497	9.7	8.0	344,976	5.8	5.5	14.8	147,248	16.3	15.0	234,409	11.7

- (1) Reporting period is January 1 through December 31.
- (2) Analyses based on one record per child.
- (3) Excludes records with unknown data or errors.
- (4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
- (5) Infants born during the reporting period included in the analysis.
- (6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
- (7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
- (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2000 - 2002 Pediatric Nutrition Surveillance⁽⁴⁾
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)						
Clinic	Breastfeeding (6)					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Adair Coun	313	5.8 (94)	162	0.0 (70)	153	0.0 (1)
Allen Coun	322	12.4 (18)	178	0.0 (70)	172	0.0 (1)
Anderson C	208	12.5 (16)	121	0.8 (22)	102	0.0 (1)
Ashland-Bo	662	6.6 (78)	336	0.3 (61)	309	0.0 (1)
Ballard Co	161	6.2 (84)	93	* *	88	* *
Bar-Riv-Wa	1,504	8.6 (42)	727	0.7 (27)	625	0.0 (1)
Barren Cty	609	7.6 (64)	350	0.9 (17)	298	0.0 (1)
Bath Count	229	3.9(125)	127	0.0 (70)	123	0.0 (1)
Belfry	262	7.3 (67)	137	0.0 (70)	132	0.0 (1)
Bell Cty M	448	3.8(131)	257	0.0 (70)	250	0.0 (1)
Bell Cty P	422	5.7 (96)	263	0.0 (70)	251	0.0 (1)
Berea	273	12.5 (16)	151	2.0 (3)	138	0.0 (1)
Betsy Lane	215	1.4(147)	128	0.0 (70)	127	0.0 (1)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2000 - 2002 Pediatric Nutrition Surveillance⁽⁴⁾
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)						
Clinic	Breastfeeding (6)					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Bledsoe	48	* *	22	* *	22	* *
Boone Coun	1,233	6.8 (75)	689	1.3 (10)	574	0.0 (1)
Bourbon Ct	310	7.1 (71)	190	0.0 (70)	179	0.0 (1)
Boyle Coun	377	5.0(107)	181	0.0 (70)	175	0.0 (1)
Bracken Co	142	9.9 (26)	69	* *	64	* *
Breathitt	346	5.2(105)	180	0.0 (70)	167	0.0 (1)
Breckinrid	285	6.7 (76)	164	0.0 (70)	150	0.0 (1)
Bridges	340	5.6 (98)	168	0.0 (70)	163	0.0 (1)
Bullitt Co	729	9.7 (29)	338	0.0 (70)	319	0.0 (1)
Butler Cty	270	4.8(112)	145	0.0 (70)	138	0.0 (1)
C & Y	788	8.6 (42)	375	0.3 (61)	334	0.0 (1)
Caldwell C	259	3.5(135)	150	0.0 (70)	139	0.0 (1)
Caldwell E	0	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2000 - 2002 Pediatric Nutrition Surveillance⁽⁴⁾
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)						
Clinic	Breastfeeding (6)					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Calloway C	557	13.6 (14)	311	1.0 (15)	285	0.0 (1)
Campbell C	923	4.7(114)	484	0.6 (34)	404	0.0 (1)
Cannonsbur	236	8.1 (54)	124	0.0 (70)	107	0.0 (1)
Carlisle C	99	* *	58	* *	56	* *
Carroll Ct	253	6.3 (83)	145	0.0 (70)	137	0.0 (1)
Carter Cty	398	8.8 (38)	214	0.5 (38)	201	0.0 (1)
Casey Cty	305	7.5 (65)	163	0.6 (34)	150	0.0 (1)
Christian	1,346	5.5(100)	753	0.5 (38)	590	0.0 (1)
Clark Coun	533	7.9 (59)	268	0.4 (51)	236	0.0 (1)
Clay Cty H	510	3.9(125)	294	0.0 (70)	287	0.0 (1)
Clinton Ct	278	2.2(139)	165	0.0 (70)	159	0.0 (1)
Corbin	661	6.2 (84)	404	0.2 (65)	343	0.0 (1)
Crittenden	149	6.0 (89)	84	* *	81	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2000 - 2002 Pediatric Nutrition Surveillance⁽⁴⁾
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)						
Clinic	Breastfeeding (6)					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Cumberland	169	5.3(103)	98	* *	96	* *
Daviess Co	1,424	7.0 (73)	757	0.9 (17)	632	0.0 (1)
Dixie	1,048	7.5 (65)	502	0.2 (65)	428	0.0 (1)
Edmonson C	232	9.5 (31)	126	0.0 (70)	94	* *
Elliott Ct	160	3.8(131)	99	* *	91	* *
Estill Cty	344	9.0 (36)	178	0.0 (70)	163	0.0 (1)
Etown	814	8.4 (48)	399	0.5 (38)	327	0.0 (1)
Evarts	194	6.7 (76)	120	0.0 (70)	117	0.0 (1)
Fairdale	616	9.3 (32)	286	0.7 (27)	251	0.0 (1)
Fleming Co	283	4.9(110)	152	0.0 (70)	150	0.0 (1)
Floyd Cty	669	2.2(139)	358	0.0 (70)	341	0.0 (1)
Foust Hlth	112	3.6(133)	64	* *	60	* *
Franklin C	671	13.9 (12)	352	1.7 (4)	275	0.0 (1)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2000 - 2002 Pediatric Nutrition Surveillance⁽⁴⁾
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)						
Clinic	Breastfeeding (6)					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Ft Campbel	1,724	15.2 (7)	960	2.2 (2)	765	0.0 (1)
Fulton	132	4.5(116)	76	* *	62	* *
Gallatin C	157	4.5(116)	78	* *	70	* *
Garrard Ct	311	7.7 (60)	183	0.5 (38)	173	0.0 (1)
Goose Rock	76	* *	41	* *	41	* *
Grant Cty	599	5.0(107)	355	0.3 (61)	339	0.0 (1)
Graves Cty	694	11.4 (22)	370	0.5 (38)	351	0.0 (1)
Grayson Ct	495	14.1 (11)	272	0.0 (70)	265	0.0 (1)
Green Cty	205	11.7 (21)	114	0.0 (70)	107	0.0 (1)
Greenup	491	8.6 (42)	268	0.4 (51)	222	0.0 (1)
H & Env.	89	* *	54	* *	47	* *
Hancock Ct	141	9.2 (33)	81	* *	70	* *
Harlan Cty	406	3.0(137)	234	0.0 (70)	208	0.0 (1)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2000 - 2002 Pediatric Nutrition Surveillance⁽⁴⁾
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)						
Clinic	Breastfeeding (6)					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Harrison C	317	8.2 (52)	183	1.1 (14)	165	0.0 (1)
Hart Cty H	379	6.1 (86)	187	0.0 (70)	171	0.0 (1)
Henderson	749	5.6 (98)	368	0.0 (70)	306	0.0 (1)
Henry Coun	238	5.9 (91)	148	1.4 (6)	127	0.0 (1)
Hickman	93	* *	47	* *	46	* *
Hickman Ct	77	* *	42	* *	38	* *
Highview	716	20.7 (3)	368	0.8 (22)	310	0.0 (1)
Hindman	362	1.7(143)	218	0.0 (70)	190	0.0 (1)
Hopkins Ct	874	5.3(103)	491	0.4 (51)	424	0.0 (1)
Iroquois	1,196	14.6 (8)	580	0.7 (27)	498	0.0 (1)
Jackson Ct	361	4.4(118)	222	0.0 (70)	215	0.0 (1)
Jessamine	626	8.9 (37)	343	0.9 (17)	290	0.0 (1)
Johnson Co	568	3.9(125)	326	0.0 (70)	312	0.0 (1)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2000 - 2002 Pediatric Nutrition Surveillance⁽⁴⁾
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)						
Clinic	Breastfeeding (6)					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
JT Arnett	38	* *	26	* *	26	* *
Kenton Cty	1,280	4.3(119)	738	0.4 (51)	606	0.0 (1)
Knox Cty	812	3.9(125)	458	0.2 (65)	426	0.0 (1)
L & N	949	5.5(100)	419	0.5 (38)	354	0.0 (1)
Larue Cty	256	9.8 (28)	144	0.0 (70)	125	0.0 (1)
Laurel Cty	1,039	5.9 (91)	533	0.8 (22)	454	0.0 (1)
Lawrence C	350	4.9(110)	190	0.5 (38)	184	0.0 (1)
Lee Cty He	181	3.9(125)	115	0.9 (17)	104	0.0 (1)
Leslie Cty	335	4.8(112)	201	0.0 (70)	182	0.0 (1)
Letcher Ct	510	6.1 (86)	290	1.0 (15)	240	0.0 (1)
Lewis Cty	291	7.2 (69)	164	0.0 (70)	158	0.0 (1)
Lex-Fayette	1,545	25.4 (1)	728	2.6 (1)	624	0.0 (1)
Lex-Fayette	388	12.4 (18)	236	1.3 (10)	202	0.0 (1)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2000 - 2002 Pediatric Nutrition Surveillance⁽⁴⁾
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)						
Clinic	Breastfeeding (6)					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Lex-Fayette	398	18.8 (4)	220	1.4 (6)	193	0.0 (1)
Lex-Fayette	552	15.9 (6)	295	1.7 (4)	258	0.0 (1)
Lex-Fayette	519	21.6 (2)	251	0.8 (22)	230	0.0 (1)
Liberty	44	* *	8	* *	4	* *
Lincoln Ct	517	7.7 (60)	285	0.4 (51)	232	0.0 (1)
Livingston	118	8.5 (45)	72	* *	68	* *
Logan Cty	481	1.7(143)	269	0.0 (70)	246	0.0 (1)
Lynch	180	8.3 (50)	99	* *	91	* *
Lyon Cty	103	9.7 (29)	64	* *	56	* *
Magoffin C	344	1.7(143)	202	0.0 (70)	195	0.0 (1)
Marion Cty	377	7.7 (60)	204	0.0 (70)	188	0.0 (1)
Marshall C	449	10.9 (24)	246	0.4 (51)	232	0.0 (1)
Marshall C	71	* *	34	* *	30	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 7A
2000 - 2002 Pediatric Nutrition Surveillance⁽⁴⁾
Kentucky**

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years⁽³⁾** (2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic^(4,5)						
Clinic	Breastfeeding⁽⁶⁾					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Martin Cty	353	2.0(142)	191	0.0 (70)	185	0.0 (1)
Mason Cty	356	6.5 (80)	188	0.0 (70)	179	0.0 (1)
McCracken	1,237	6.5 (80)	678	0.0 (70)	644	0.0 (1)
McCreary C	486	3.9(125)	287	0.0 (70)	273	0.0 (1)
McLean Ct	251	9.2 (33)	149	1.3 (10)	144	0.0 (1)
Meade Cty	423	14.2 (9)	235	0.9 (17)	200	0.0 (1)
Menifee Ct	157	8.3 (50)	96	* *	95	* *
Mercer Cty	349	5.7 (96)	185	0.0 (70)	162	0.0 (1)
Metcalfe C	208	2.9(138)	117	0.0 (70)	101	0.0 (1)
Middletown	641	18.1 (5)	296	0.7 (27)	270	0.0 (1)
Monroe Cnt	242	8.7 (40)	142	1.4 (6)	112	0.0 (1)
Montgomery	517	6.0 (89)	285	0.4 (51)	242	0.0 (1)
Morgan Cty	299	4.3(119)	183	0.5 (38)	173	0.0 (1)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2000 - 2002 Pediatric Nutrition Surveillance⁽⁴⁾
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)						
Clinic	Breastfeeding (6)					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Muhlenberg	606	4.3(119)	344	0.0 (70)	329	0.0 (1)
Neighborho	743	11.4 (22)	407	0.7 (27)	348	0.0 (1)
Nelson Cty	609	9.2 (33)	328	0.0 (70)	285	0.0 (1)
Newburg	855	8.7 (40)	429	0.7 (27)	367	0.0 (1)
Nicholas C	171	5.8 (94)	88	* *	75	* *
Oasis Hlth	100	8.0 (55)	60	* *	55	* *
Ohio Cty	464	8.4 (48)	249	0.0 (70)	222	0.0 (1)
Oldham Cty	356	11.8 (20)	197	0.5 (38)	185	0.0 (1)
Olive Hill	301	7.0 (73)	175	0.0 (70)	166	0.0 (1)
Oneida Ele	30	* *	18	* *	17	* *
Owen Cty	162	13.6 (14)	79	* *	72	* *
Owsley Cty	135	2.2(139)	83	* *	77	* *
Park Duval	909	8.0 (55)	445	0.4 (51)	383	0.0 (1)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2000 - 2002 Pediatric Nutrition Surveillance⁽⁴⁾
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)						
Clinic	Breastfeeding (6)					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Pendleton	252	9.9 (26)	148	0.7 (27)	141	0.0 (1)
Perry Cty	622	4.0(124)	360	0.3 (61)	346	0.0 (1)
Phelps	74	* *	35	* *	27	* *
Pike Cty	892	6.5 (80)	481	0.2 (65)	401	0.0 (1)
Portland	786	6.6 (78)	397	0.5 (38)	328	0.0 (1)
Powell Cty	368	4.1(123)	194	0.5 (38)	185	0.0 (1)
Prenatal	103	0.0(148)	2	* *	0	* *
Prenatal	35	* *	0	* *	0	* *
Pulaski Ct	1,204	10.2 (25)	641	0.6 (34)	548	0.0 (1)
Radcliffe	1,215	8.8 (38)	668	0.1 (69)	556	0.0 (1)
Richmond	855	8.2 (52)	429	1.2 (13)	360	0.0 (1)
Robertson	60	* *	31	* *	30	* *
Rockcastle	382	7.3 (67)	187	0.5 (38)	167	0.0 (1)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2000 - 2002 Pediatric Nutrition Surveillance⁽⁴⁾
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)						
Clinic	Breastfeeding (6)					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Rowan Cty	419	3.6(133)	235	0.4 (51)	203	0.0 (1)
Russell Ct	387	8.0 (55)	221	0.0 (70)	210	0.0 (1)
Scott Cty	472	8.5 (45)	240	0.4 (51)	223	0.0 (1)
Shelby Cty	536	5.4(102)	281	0.0 (70)	255	0.0 (1)
Siler	24	* *	19	* *	13	* *
Simpson Ct	293	8.5 (45)	152	0.0 (70)	137	0.0 (1)
South Shor	89	* *	50	* *	43	* *
Spencer Ct	163	6.1 (86)	88	* *	83	* *
TAPP-South	61	* *	30	* *	25	* *
TAPP-West	59	* *	33	* *	28	* *
Taylor Cty	454	7.7 (60)	255	0.0 (70)	242	0.0 (1)
Todd Cty H	263	7.2 (69)	160	0.6 (34)	152	0.0 (1)
Trigg Cty	181	13.8 (13)	102	0.0 (70)	94	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2000 - 2002 Pediatric Nutrition Surveillance⁽⁴⁾
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5)						
Clinic	Breastfeeding (6)					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Trimble Ct	150	8.0 (55)	86	* *	84	* *
Union Cty	294	7.1 (71)	170	0.0 (70)	147	0.0 (1)
Wallins	69	* *	43	* *	42	* *
Washington	219	4.6(115)	130	0.0 (70)	124	0.0 (1)
Wayne Cty	498	5.2(105)	264	0.8 (22)	238	0.0 (1)
Webster Ct	256	5.9 (91)	141	1.4 (6)	105	0.0 (1)
Wheelbrigh	142	4.2(122)	78	* *	69	* *
Williamsbu	518	5.0(107)	292	0.0 (70)	256	0.0 (1)
Wolfe Cty	239	3.3(136)	140	0.0 (70)	132	0.0 (1)
Woodford C	344	14.2 (9)	207	0.5 (38)	189	0.0 (1)
Unknown Clinics	1,242	1.6(146)	62	* *	57	* *
Kentucky	73,122	8.1	38,745	0.5	34,374	0.0

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.