

**Table 4B**  
**2002 Pediatric Nutrition Surveillance** <sup>(1)</sup>  
**Kentucky**  
**Comparison of Racial and Ethnic Distribution by County**  
**Children Aged < 5 Years** <sup>(2)</sup>

<b>Comparison of Racial and Ethnic Distribution by County</b>							
<b>County</b>	<b>Number Included in Analysis (3)</b>	<b>White, Not Hispanic %</b>	<b>Black, Not Hispanic %</b>	<b>Hispanic %</b>	<b>American Indian/Alaskan %</b>	<b>Asian/Pacific Islander %</b>	<b>All Other / Unknown %</b>
Adair	515	93.2	4.9	1.9	0.0	0.0	0.0
Allen	600	96.8	1.0	1.0	0.0	1.2	0.0
Anderson	345	91.3	5.2	3.5	0.0	0.0	0.0
Ballard	268	89.2	10.1	0.4	0.0	0.4	0.0
Barren	995	90.8	6.6	1.8	0.2	0.6	0.0
Bath	427	97.9	1.2	0.5	0.5	0.0	0.0
Bell	1,632	95.6	2.3	1.8	0.2	0.1	0.0
Boone	1,980	88.5	3.3	7.6	0.0	0.5	0.0
Bourbon	474	78.1	9.1	11.6	0.2	1.1	0.0
Boyd	1,425	96.8	1.6	1.3	0.1	0.1	0.0
Boyle	567	94.2	5.5	0.0	0.0	0.4	0.0
Bracken	230	99.6	0.0	0.4	0.0	0.0	0.0
Breathitt	687	99.3	0.6	0.0	0.1	0.0	0.0
Breckinrid	492	95.7	3.9	0.4	0.0	0.0	0.0
Bullitt	1,122	99.1	0.5	0.1	0.0	0.3	0.0
Butler	456	96.9	1.1	0.0	0.0	2.0	0.0
Caldwell	421	92.2	7.4	0.2	0.0	0.2	0.0
Calloway	919	86.6	7.9	3.4	0.1	2.0	0.0
Campbell	1,425	92.7	6.0	0.9	0.1	0.3	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

\* Percentages are not calculated if < 100 records are available for analysis after exclusions.

**Table 4B**  
**2002 Pediatric Nutrition Surveillance** <sup>(1)</sup>  
**Kentucky**  
**Comparison of Racial and Ethnic Distribution by County**  
**Children Aged < 5 Years** <sup>(2)</sup>

<b>Comparison of Racial and Ethnic Distribution by County</b>							
<b>County</b>	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan %	Asian/Pacific Islander %	All Other / Unknown %
Carlisle	190	93.7	5.8	0.5	0.0	0.0	0.0
Carroll	405	84.9	1.2	13.8	0.0	0.0	0.0
Carter	1,256	99.2	0.4	0.0	0.3	0.1	0.0
Casey	552	97.3	0.0	2.7	0.0	0.0	0.0
Christian	4,981	64.1	30.2	4.9	0.1	0.8	0.0
Clark	823	89.6	6.6	3.3	0.0	0.6	0.0
Clay	1,139	98.0	1.1	0.9	0.1	0.0	0.0
Clinton	506	95.3	0.2	4.2	0.0	0.4	0.0
Crittenden	256	99.6	0.4	0.0	0.0	0.0	0.0
Cumberland	296	93.6	4.1	2.4	0.0	0.0	0.0
Daviess	2,646	85.4	11.0	3.1	0.1	0.5	0.0
Edmonson	362	100.0	0.0	0.0	0.0	0.0	0.0
Elliott	298	99.3	0.0	0.3	0.3	0.0	0.0
Estill	595	98.8	0.7	0.0	0.0	0.5	0.0
Fayette	5,022	47.9	31.0	14.9	0.4	5.8	0.0
Fleming	551	97.5	1.6	0.9	0.0	0.0	0.0
Floyd	1,854	99.5	0.3	0.2	0.0	0.0	0.0
Franklin	1,011	77.0	16.2	4.5	0.3	2.1	0.0
Fulton	373	59.5	40.5	0.0	0.0	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

\* Percentages are not calculated if < 100 records are available for analysis after exclusions.

**Table 4B**  
**2002 Pediatric Nutrition Surveillance** <sup>(1)</sup>  
**Kentucky**  
**Comparison of Racial and Ethnic Distribution by County**  
**Children Aged < 5 Years** <sup>(2)</sup>

<b>Comparison of Racial and Ethnic Distribution by County</b>							
<b>County</b>	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan %	Asian/Pacific Islander %	All Other / Unknown %
Gallatin	264	84.1	2.3	13.3	0.0	0.4	0.0
Garrard	551	94.7	2.4	2.7	0.2	0.0	0.0
Grant	1,061	97.8	0.9	0.4	0.0	0.8	0.0
Graves	1,199	75.6	10.3	11.8	0.1	2.3	0.0
Grayson	901	98.6	0.9	0.4	0.1	0.0	0.0
Green	332	95.2	3.6	0.9	0.0	0.3	0.0
Greenup	976	99.1	0.8	0.0	0.0	0.1	0.0
Hancock	258	97.7	0.8	1.2	0.4	0.0	0.0
Hardin	3,476	77.7	19.5	2.0	0.1	0.6	0.0
Harlan	1,658	97.3	2.4	0.2	0.0	0.1	0.0
Harrison	536	92.4	2.8	4.9	0.0	0.0	0.0
Hart	627	95.4	4.3	0.3	0.0	0.0	0.0
Henderson	1,152	84.5	11.9	0.9	0.4	2.3	0.0
Henry	370	88.4	3.5	8.1	0.0	0.0	0.0
Hickman	127	80.3	17.3	1.6	0.0	0.8	0.0
Hopkins	1,513	84.6	13.0	2.0	0.1	0.3	0.0
Jackson	640	100.0	0.0	0.0	0.0	0.0	0.0
Jefferson	14,738	48.2	44.7	4.4	0.3	2.3	0.0
Jessamine	1,053	90.8	5.2	2.6	0.1	1.3	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

\* Percentages are not calculated if < 100 records are available for analysis after exclusions.

**Table 4B**  
**2002 Pediatric Nutrition Surveillance** <sup>(1)</sup>  
**Kentucky**  
**Comparison of Racial and Ethnic Distribution by County**  
**Children Aged < 5 Years** <sup>(2)</sup>

<b>Comparison of Racial and Ethnic Distribution by County</b>							
<b>County</b>	<b>Number Included in Analysis (3)</b>	<b>White, Not Hispanic %</b>	<b>Black, Not Hispanic %</b>	<b>Hispanic %</b>	<b>American Indian/Alaskan %</b>	<b>Asian/Pacific Islander %</b>	<b>All Other / Unknown %</b>
Johnson	1,038	100.0	0.0	0.0	0.0	0.0	0.0
Kenton	2,166	83.7	14.9	1.0	0.2	0.3	0.0
Knott	720	99.4	0.6	0.0	0.0	0.0	0.0
Knox	1,520	98.8	1.1	0.1	0.0	0.0	0.0
Larue	434	89.2	5.5	4.6	0.7	0.0	0.0
Laurel	1,640	98.7	0.7	0.3	0.0	0.3	0.0
Lawrence	600	99.8	0.2	0.0	0.0	0.0	0.0
Lee	347	99.7	0.3	0.0	0.0	0.0	0.0
Leslie	650	99.5	0.5	0.0	0.0	0.0	0.0
Letcher	866	99.4	0.5	0.0	0.0	0.1	0.0
Lewis	500	99.6	0.4	0.0	0.0	0.0	0.0
Lincoln	874	93.6	3.0	3.3	0.0	0.1	0.0
Livingston	198	97.5	0.0	1.0	0.0	1.5	0.0
Logan	716	80.7	16.2	2.4	0.7	0.0	0.0
Lyon	172	98.8	1.2	0.0	0.0	0.0	0.0
Madison	1,785	89.9	7.7	1.1	0.1	1.2	0.0
Magoffin	727	98.3	0.0	1.7	0.0	0.0	0.0
Marion	677	83.3	11.5	4.7	0.0	0.4	0.0
Marshall	859	99.5	0.5	0.0	0.0	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

\* Percentages are not calculated if < 100 records are available for analysis after exclusions.

**Table 4B**  
**2002 Pediatric Nutrition Surveillance** <sup>(1)</sup>  
**Kentucky**  
**Comparison of Racial and Ethnic Distribution by County**  
**Children Aged < 5 Years** <sup>(2)</sup>

<b>Comparison of Racial and Ethnic Distribution by County</b>							
<b>County</b>	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan %	Asian/Pacific Islander %	All Other / Unknown %
Martin	711	99.9	0.0	0.1	0.0	0.0	0.0
Mason	592	90.9	7.3	0.7	0.5	0.7	0.0
McCracken	1,988	63.5	33.0	1.6	0.5	1.4	0.0
McCreary	884	99.8	0.2	0.0	0.0	0.0	0.0
McLean	452	97.3	0.7	1.8	0.0	0.2	0.0
Meade	697	97.1	2.9	0.0	0.0	0.0	0.0
Menifee	286	99.0	0.0	1.0	0.0	0.0	0.0
Mercer	592	89.2	4.7	5.7	0.0	0.3	0.0
Metcalfe	350	97.7	1.1	0.9	0.3	0.0	0.0
Monroe	408	88.7	5.6	5.4	0.2	0.0	0.0
Montgomery	872	95.5	1.3	2.6	0.0	0.6	0.0
Morgan	564	99.8	0.0	0.2	0.0	0.0	0.0
Muhlenberg	1,033	96.6	3.0	0.3	0.0	0.1	0.0
Nelson	1,025	89.7	8.5	1.4	0.2	0.3	0.0
Nicholas	285	98.9	0.0	1.1	0.0	0.0	0.0
Ohio	830	95.3	0.4	3.7	0.0	0.6	0.0
Oldham	588	82.8	5.1	11.7	0.0	0.3	0.0
Owen	280	92.9	0.0	6.4	0.4	0.4	0.0
Owsley	272	99.6	0.4	0.0	0.0	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

\* Percentages are not calculated if < 100 records are available for analysis after exclusions.

**Table 4B**  
**2002 Pediatric Nutrition Surveillance** <sup>(1)</sup>  
**Kentucky**  
**Comparison of Racial and Ethnic Distribution by County**  
**Children Aged < 5 Years** <sup>(2)</sup>

<b>Comparison of Racial and Ethnic Distribution by County</b>							
<b>County</b>	<b>Number Included in Analysis (3)</b>	<b>White, Not Hispanic %</b>	<b>Black, Not Hispanic %</b>	<b>Hispanic %</b>	<b>American Indian/Alaskan %</b>	<b>Asian/Pacific Islander %</b>	<b>All Other / Unknown %</b>
Pendleton	404	96.0	2.0	1.0	0.7	0.2	0.0
Perry	1,209	97.8	1.9	0.2	0.0	0.1	0.0
Pike	2,071	99.5	0.4	0.0	0.0	0.1	0.0
Powell	681	97.8	0.6	0.9	0.0	0.7	0.0
Pulaski	1,901	97.4	0.9	1.5	0.1	0.1	0.0
Robertson	111	94.6	0.9	4.5	0.0	0.0	0.0
Rockcastle	632	99.7	0.2	0.2	0.0	0.0	0.0
Rowan	690	97.1	1.0	1.2	0.4	0.3	0.0
Russell	607	95.6	0.2	3.1	0.2	1.0	0.0
Scott	723	86.7	7.6	4.8	0.4	0.4	0.0
Shelby	791	52.2	13.5	34.3	0.0	0.0	0.0
Simpson	446	83.6	14.6	0.9	0.2	0.7	0.0
Spencer	275	92.7	1.5	5.1	0.4	0.4	0.0
Taylor	783	92.3	7.4	0.1	0.1	0.0	0.0
Todd	460	81.5	11.5	5.4	0.2	1.3	0.0
Trigg	301	82.7	15.6	0.0	0.7	1.0	0.0
Trimble	247	96.4	0.4	3.2	0.0	0.0	0.0
Union	494	85.4	13.6	0.2	0.0	0.8	0.0
Warren	2,281	68.7	15.3	13.3	0.3	2.5	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

\* Percentages are not calculated if < 100 records are available for analysis after exclusions.

**Table 4B**  
**2002 Pediatric Nutrition Surveillance (1)**  
**Kentucky**  
**Comparison of Racial and Ethnic Distribution by County**  
**Children Aged < 5 Years (2)**

<b>Comparison of Racial and Ethnic Distribution by County</b>							
<b>County</b>	<b>Number Included in Analysis (3)</b>	<b>White, Not Hispanic %</b>	<b>Black, Not Hispanic %</b>	<b>Hispanic %</b>	<b>American Indian/Alaskan %</b>	<b>Asian/Pacific Islander %</b>	<b>All Other / Unknown %</b>
Washington	374	75.7	12.6	11.5	0.0	0.3	0.0
Wayne	879	92.3	1.6	6.1	0.0	0.0	0.0
Webster	428	85.7	4.2	7.2	0.0	2.8	0.0
Whitley	2,211	99.7	0.1	0.0	0.0	0.1	0.0
Wolfe	449	99.3	0.7	0.0	0.0	0.0	0.0
Woodford	533	78.2	4.7	16.5	0.2	0.4	0.0
Unknown Counties	1,324	83.8	12.8	2.7	0.0	0.6	0.0
<b>Kentucky</b>	<b>120,031</b>	<b>82.9</b>	<b>12.6</b>	<b>3.4</b>	<b>0.1</b>	<b>0.9</b>	<b>0.0</b>
<b>Nation (Prior Year)</b>	<b>5,174,210</b>	<b>40.2</b>	<b>21.4</b>	<b>31.4</b>	<b>1.4</b>	<b>2.8</b>	<b>2.9</b>

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

\* Percentages are not calculated if < 100 records are available for analysis after exclusions.

**Table 5B**  
**2002 Pediatric Nutrition Surveillance (1)**  
**Kentucky**  
**Comparison of Age Distribution by County**  
**Children Aged < 5 Years (2)**

<b>Comparison of Age Distribution by County</b>				
<b>County</b>	Number Included in Analysis (3,4)	0 - 11 Months  %	12 - 23 Months  %	24 - 59 Months  %
Adair	515	29.1	25.6	45.2
Allen	600	24.8	24.3	50.8
Anderson	345	31.0	25.2	43.8
Ballard	268	25.7	28.7	45.5
Barren	995	32.3	26.3	41.4
Bath	427	25.5	27.6	46.8
Bell	1,632	23.3	25.4	51.3
Boone	1,980	31.7	25.5	42.8
Bourbon	474	27.0	32.3	40.7
Boyd	1,425	33.9	25.7	40.4
Boyle	567	35.1	26.3	38.6
Bracken	230	33.9	23.5	42.6
Breathitt	687	24.6	21.1	54.3
Breckinrid	492	27.2	26.4	46.3
Bullitt	1,122	36.5	25.6	37.9
Butler	456	28.9	25.2	45.8
Caldwell	421	29.9	27.3	42.8
Calloway	919	30.5	26.7	42.9

(1) Reporting period is January 1 through December 31.  
(2) Analyses based on one record per child.  
(3) Excludes records with unknown data or errors.  
(4) Records rejected if date of birth and date of visit unknown.  
\* Percentages are not calculated if < 100 records are available for analysis after exclusions.



**Table 5B**  
**2002 Pediatric Nutrition Surveillance** (1)  
**Kentucky**  
**Comparison of Age Distribution by County**  
**Children Aged < 5 Years** (2)

<b>Comparison of Age Distribution by County</b>				
<b>County</b>	Number Included in Analysis (3,4)	0 - 11 Months  %	12 - 23 Months  %	24 - 59 Months  %
Campbell	1,425	35.2	23.4	41.3
Carlisle	190	20.0	25.8	54.2
Carroll	405	28.9	28.4	42.7
Carter	1,256	26.2	25.6	48.2
Casey	552	29.0	23.2	47.8
Christian	4,981	34.8	22.4	42.8
Clark	823	36.6	23.6	39.9
Clay	1,139	23.5	26.3	50.2
Clinton	506	23.3	27.7	49.0
Crittenden	256	26.2	28.1	45.7
Cumberland	296	23.6	28.4	48.0
Daviess	2,646	33.1	24.3	42.6
Edmonson	362	39.0	19.6	41.4
Elliott	298	21.8	26.5	51.7
Estill	595	30.3	24.2	45.5
Fayette	5,022	37.2	25.9	36.9
Fleming	551	24.0	25.4	50.6
Floyd	1,854	26.1	25.7	48.3

(1) Reporting period is January 1 through December 31.  
(2) Analyses based on one record per child.  
(3) Excludes records with unknown data or errors.  
(4) Records rejected if date of birth and date of visit unknown.  
\* Percentages are not calculated if < 100 records are available for analysis after exclusions.

**Table 5B**  
**2002 Pediatric Nutrition Surveillance (1)**  
**Kentucky**  
**Comparison of Age Distribution by County**  
**Children Aged < 5 Years (2)**

<b>Comparison of Age Distribution by County</b>				
<b>County</b>	Number Included in Analysis (3,4)	0 - 11 Months  %	12 - 23 Months  %	24 - 59 Months  %
Franklin	1,011	40.7	22.8	36.5
Fulton	373	31.6	24.1	44.2
Gallatin	264	32.2	23.5	44.3
Garrard	551	25.0	27.8	47.2
Grant	1,061	22.2	27.0	50.7
Graves	1,199	27.9	26.4	45.6
Grayson	901	25.0	26.1	48.9
Green	332	30.1	28.0	41.9
Greenup	976	32.9	23.5	43.6
Hancock	258	27.5	23.3	49.2
Hardin	3,476	33.7	21.9	44.4
Harlan	1,658	26.0	23.5	50.5
Harrison	536	27.1	27.4	45.5
Hart	627	33.2	23.6	43.2
Henderson	1,152	38.6	22.3	39.1
Henry	370	30.0	30.0	40.0
Hickman	127	30.7	26.8	42.5
Hopkins	1,513	31.0	23.5	45.5

(1) Reporting period is January 1 through December 31.  
(2) Analyses based on one record per child.  
(3) Excludes records with unknown data or errors.  
(4) Records rejected if date of birth and date of visit unknown.  
\* Percentages are not calculated if < 100 records are available for analysis after exclusions.

**Table 5B**  
**2002 Pediatric Nutrition Surveillance (1)**  
**Kentucky**  
**Comparison of Age Distribution by County**  
**Children Aged < 5 Years (2)**

<b>Comparison of Age Distribution by County</b>				
<b>County</b>	Number Included in Analysis (3,4)	0 - 11 Months  %	12 - 23 Months  %	24 - 59 Months  %
Jackson	640	21.9	29.5	48.6
Jefferson	14,738	38.0	23.9	38.1
Jessamine	1,053	32.0	24.3	43.7
Johnson	1,038	24.6	26.0	49.4
Kenton	2,166	31.8	26.7	41.5
Knott	720	25.3	21.0	53.8
Knox	1,520	25.6	24.5	49.9
Larue	434	31.3	24.0	44.7
Laurel	1,640	36.3	23.3	40.4
Lawrence	600	27.2	28.3	44.5
Lee	347	23.3	27.1	49.6
Leslie	650	24.2	22.0	53.8
Letcher	866	31.8	23.0	45.3
Lewis	500	27.0	26.8	46.2
Lincoln	874	33.9	22.2	43.9
Livingston	198	25.3	31.3	43.4
Logan	716	33.7	28.6	37.7
Lyon	172	29.1	27.3	43.6

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

\* Percentages are not calculated if < 100 records are available for analysis after exclusions.

**Table 5B**  
**2002 Pediatric Nutrition Surveillance** <sup>(1)</sup>  
**Kentucky**  
**Comparison of Age Distribution by County**  
**Children Aged < 5 Years** <sup>(2)</sup>

<b>Comparison of Age Distribution by County</b>				
<b>County</b>	Number Included in Analysis (3,4)	0 - 11 Months  %	12 - 23 Months  %	24 - 59 Months  %
Madison	1,785	35.9	23.6	40.5
Magoffin	727	22.7	24.6	52.7
Marion	677	28.8	25.1	46.1
Marshall	859	29.6	26.9	43.5
Martin	711	23.5	23.1	53.4
Mason	592	30.2	26.4	43.4
McCracken	1,988	29.3	28.0	42.7
McCreary	884	24.3	26.5	49.2
McLean	452	24.3	27.7	48.0
Meade	697	31.7	23.4	44.9
Menifee	286	22.7	26.2	51.0
Mercer	592	33.8	22.6	43.6
Metcalfe	350	31.1	23.7	45.1
Monroe	408	31.4	22.5	46.1
Montgomery	872	32.7	22.9	44.4
Morgan	564	23.2	25.5	51.2
Muhlenberg	1,033	27.0	28.1	44.9
Nelson	1,025	32.4	23.6	44.0

(1) Reporting period is January 1 through December 31.  
(2) Analyses based on one record per child.  
(3) Excludes records with unknown data or errors.  
(4) Records rejected if date of birth and date of visit unknown.  
\* Percentages are not calculated if < 100 records are available for analysis after exclusions.

**Table 5B**  
**2002 Pediatric Nutrition Surveillance** (1)  
**Kentucky**  
**Comparison of Age Distribution by County**  
**Children Aged < 5 Years** (2)

<b>Comparison of Age Distribution by County</b>				
<b>County</b>	Number Included in Analysis (3,4)	0 - 11 Months  %	12 - 23 Months  %	24 - 59 Months  %
Nicholas	285	32.3	23.9	43.9
Ohio	830	30.1	22.5	47.3
Oldham	588	29.6	27.0	43.4
Owen	280	32.9	22.9	44.3
Owsley	272	21.7	26.1	52.2
Pendleton	404	27.5	30.0	42.6
Perry	1,209	31.8	24.5	43.8
Pike	2,071	32.5	22.6	44.9
Powell	681	27.6	24.1	48.3
Pulaski	1,901	35.7	24.1	40.1
Robertson	111	25.2	26.1	48.6
Rockcastle	632	33.7	22.8	43.5
Rowan	690	31.9	24.1	44.1
Russell	607	29.7	28.7	41.7
Scott	723	35.8	27.2	36.9
Shelby	791	38.2	26.9	34.9
Simpson	446	34.3	27.4	38.3
Spencer	275	28.7	28.0	43.3

(1) Reporting period is January 1 through December 31.  
(2) Analyses based on one record per child.  
(3) Excludes records with unknown data or errors.  
(4) Records rejected if date of birth and date of visit unknown.  
\* Percentages are not calculated if < 100 records are available for analysis after exclusions.

**Table 5B**  
**2002 Pediatric Nutrition Surveillance (1)**  
**Kentucky**  
**Comparison of Age Distribution by County**  
**Children Aged < 5 Years (2)**

<b>Comparison of Age Distribution by County</b>				
<b>County</b>	Number Included in Analysis (3,4)	0 - 11 Months  %	12 - 23 Months  %	24 - 59 Months  %
Taylor	783	27.2	26.6	46.2
Todd	460	23.7	30.2	46.1
Trigg	301	28.9	26.9	44.2
Trimble	247	24.7	29.6	45.7
Union	494	31.2	25.9	42.9
Warren	2,281	39.5	23.3	37.2
Washington	374	25.9	29.1	44.9
Wayne	879	29.8	22.1	48.1
Webster	428	37.4	20.3	42.3
Whitley	2,211	29.0	23.0	47.9
Wolfe	449	24.1	26.3	49.7
Woodford	533	30.0	30.8	39.2
Unknown Counties	1,324	89.7	3.2	7.1
<b>Kentucky</b>	<b>120,031</b>	<b>32.4</b>	<b>24.5</b>	<b>43.1</b>
<b>Nation (Prior Year)</b>	<b>5,174,210</b>	<b>37.3</b>	<b>21.6</b>	<b>41.1</b>

(1) Reporting period is January 1 through December 31.  
(2) Analyses based on one record per child.  
(3) Excludes records with unknown data or errors.  
(4) Records rejected if date of birth and date of visit unknown.  
\* Percentages are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**2002 Pediatric Nutrition Surveillance** (1)  
**Kentucky**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years** (2)

<b>Comparison of Growth and Anemia Indicators by County</b> (3,4)												
<b>County</b>	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low		High	Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Adair	135	6.7 (11)	6.7 (30)	501	6.8(50)	5.2(88)	12.4(12)	223	13.9(14)	11.7 (9)	305	5.2 (14)
Allen	138	10.9 (67)	10.1 (79)	585	7.4(58)	3.4(38)	17.1(63)	290	22.8(10)	19.0(78)	368	8.4 (49)
Anderson	81	* *	* *	327	9.2(92)	5.8(95)	20.8(10)	147	19.0(75)	23.8(10)	253	5.9 (20)
Ballard	62	* *	* *	250	7.6(63)	6.0(10)	16.4(53)	119	9.2 (2)	10.9 (8)	198	4.5 (10)
Barren	250	9.6 (47)	7.2 (42)	949	9.1(89)	3.6(47)	18.2(77)	397	17.6(52)	17.4(62)	671	9.1 (53)
Bath	100	6.0 (7)	16.0 (92)	416	12.3(11)	2.4(15)	20.4(96)	200	19.5(84)	20.5(88)	272	9.2 (54)
Bell	334	9.3 (44)	6.6 (25)	1,563	7.4(58)	3.6(47)	19.9(90)	819	17.2(44)	17.1(59)	1,081	7.6 (39)
Boone	487	10.3 (59)	9.2 (74)	1,895	8.6(79)	4.9(82)	18.3(79)	826	18.9(73)	18.3(71)	1,384	14.7 (94)
Bourbon	117	11.1 (72)	9.4 (75)	450	8.9(84)	2.0 (9)	21.1(10)	178	22.5(10)	21.9(98)	342	29.8(118)
Boyd	425	11.3 (74)	7.5 (48)	1,357	6.8(50)	10.2(11)	14.2(22)	561	18.0(59)	16.8(53)	880	10.9 (68)
Boyle	181	12.7 (87)	6.6 (25)	527	9.1(89)	5.3(91)	16.3(51)	199	19.1(77)	18.1(68)	325	8.3 (48)

- (1) Reporting period is January 1 through December 31.  
(2) Analyses based on one record per child.  
(3) Excludes records with unknown data or errors.  
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.  
(5) Infants born during the reporting period included in the analysis.  
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.  
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.  
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.  
\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**2002 Pediatric Nutrition Surveillance** (1)  
**Kentucky**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years** (2)

<b>Comparison of Growth and Anemia Indicators by County</b> (3,4)												
<b>County</b>	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Bracken	69	* *	* *	220	8.2(71)	6.8(10)	22.7(10)	95	* *	* *	146	4.1 (6)
Breathitt	154	8.4 (27)	9.1 (72)	639	6.4(39)	3.3(33)	20.0(91)	361	14.4(17)	17.2(60)	506	5.3 (15)
Breckinrid	113	8.0 (21)	6.2 (20)	481	7.5(62)	3.3(33)	15.0(33)	226	21.2(92)	13.3(25)	367	3.3 (4)
Bullitt	368	8.2 (24)	9.0 (70)	1,067	8.2(71)	3.6(47)	16.8(59)	410	15.4(28)	17.6(64)	625	19.2(110)
Butler	115	6.1 (8)	7.0 (38)	446	5.4(23)	4.7(76)	19.5(88)	204	19.1(77)	24.5(11)	306	7.2 (33)
Caldwell	108	2.8 (1)	7.4 (45)	405	6.7(46)	7.2(10)	16.3(51)	178	15.2(26)	16.3(47)	298	15.8(101)
Calloway	243	8.6 (31)	13.2 (90)	892	7.0(54)	3.5(44)	14.9(32)	388	17.3(46)	15.2(38)	582	16.2(103)
Campbell	385	11.2 (73)	8.1 (56)	1,340	8.4(75)	3.3(33)	18.1(76)	563	20.2(87)	23.3(10)	873	14.7 (94)
Carlisle	35	* *	* *	183	6.0(33)	2.7(18)	22.4(10)	102	18.6(65)	24.5(11)	138	24.6(116)
Carroll	106	8.5 (29)	4.7 (8)	396	11.9(11)	3.8(56)	20.7(10)	171	25.1(11)	24.0(10)	295	11.5 (72)
Carter	300	6.3 (9)	11.3 (85)	1,205	3.9 (8)	7.2(10)	12.3(10)	581	15.0(23)	12.6(19)	793	9.6 (58)

- (1) Reporting period is January 1 through December 31.  
(2) Analyses based on one record per child.  
(3) Excludes records with unknown data or errors.  
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.  
(5) Infants born during the reporting period included in the analysis.  
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.  
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.  
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.  
\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



**Table 6B**  
**2002 Pediatric Nutrition Surveillance** (1)  
**Kentucky**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years** (2)

<b>Comparison of Growth and Anemia Indicators by County</b> (3,4)												
<b>County</b>	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low		High	Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Casey	136	10.3 (59)	3.7 (5)	532	9.4(96)	4.7(76)	13.2(16)	254	15.0(23)	13.0(21)	337	11.6 (75)
Christian	1,231	9.0 (39)	8.6 (62)	4,788	5.6(26)	6.9(10)	10.3 (3)	2,092	12.2 (9)	8.1 (5)	3,334	26.7(117)
Clark	255	14.1 (90)	8.6 (62)	791	7.7(65)	3.7(53)	15.7(40)	315	21.9(99)	15.2(38)	495	12.7 (84)
Clay	257	11.3 (74)	6.2 (20)	1,098	6.5(41)	5.9(97)	20.3(94)	551	19.1(77)	19.4(83)	846	7.8 (43)
Clinton	111	9.0 (39)	12.6 (89)	486	6.0(33)	4.3(68)	16.7(57)	239	16.3(31)	14.2(34)	308	5.8 (19)
Crittenden	62	* *	* *	248	5.6(26)	6.5(10)	12.1 (7)	110	16.4(32)	6.4 (1)	181	11.0 (69)
Cumberland	66	* *	* *	290	3.4 (4)	3.8(56)	14.8(31)	140	17.1(41)	13.6(27)	160	13.8 (90)
Daviess	722	9.3 (44)	5.4 (14)	2,551	10.5(10)	2.1(11)	21.2(10)	1,112	17.0(39)	19.3(82)	1,831	13.7 (88)
Edmonson	102	9.8 (51)	5.9 (17)	350	4.9(20)	4.9(82)	13.1(14)	146	18.5(63)	15.1(37)	219	4.6 (11)
Elliott	55	* *	* *	287	3.1 (2)	4.9(82)	14.3(24)	151	17.2(44)	13.2(24)	213	9.9 (61)
Estill	158	6.3 (9)	7.6 (51)	578	3.8 (7)	4.8(80)	15.9(42)	267	18.0(59)	18.4(72)	421	7.1 (31)

- (1) Reporting period is January 1 through December 31.  
(2) Analyses based on one record per child.  
(3) Excludes records with unknown data or errors.  
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.  
(5) Infants born during the reporting period included in the analysis.  
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.  
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.  
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.  
\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**2002 Pediatric Nutrition Surveillance** (1)  
**Kentucky**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years** (2)

<b>Comparison of Growth and Anemia Indicators by County</b> (3,4)												
<b>County</b>	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low		High	Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Fayette	1,554	12.5 (83)	7.5 (48)	4,791	7.2(56)	4.0(63)	14.3(24)	1,810	19.4(83)	15.6(43)	3,073	15.6 (99)
Fleming	127	7.1 (12)	11.8 (88)	539	6.7(46)	5.2(88)	14.5(27)	275	16.4(32)	16.0(45)	418	16.7(105)
Floyd	440	7.7 (17)	8.0 (55)	1,779	4.5(16)	9.4(11)	13.3(18)	856	12.3(10)	14.1(32)	1,320	7.1 (31)
Franklin	308	8.1 (23)	6.2 (20)	968	8.9(84)	7.0(10)	18.6(84)	363	18.7(68)	21.5(96)	683	13.2 (85)
Fulton	95	* *	* *	353	6.5(41)	2.8(19)	20.1(93)	165	23.0(11)	20.6(89)	248	14.9 (96)
Gallatin	74	* *	* *	255	5.5(25)	5.1(87)	17.3(66)	115	10.4 (4)	20.9(91)	173	11.6 (75)
Garrard	119	8.4 (27)	6.7 (30)	527	4.9(20)	3.8(56)	16.1(45)	255	16.9(37)	18.4(72)	340	11.5 (72)
Grant	207	9.7 (50)	7.2 (42)	1,026	5.6(26)	5.6(92)	11.2 (5)	521	8.1 (1)	7.7 (4)	836	5.5 (16)
Graves	313	8.6 (31)	8.9 (68)	1,145	7.9(67)	5.9(97)	17.9(73)	532	20.5(88)	18.2(69)	771	15.6 (99)
Grayson	201	9.0 (39)	6.5 (24)	879	13.2(11)	2.2(13)	20.4(96)	432	19.0(75)	17.6(64)	680	2.4 (2)
Green	93	* *	* *	318	6.3(38)	1.9 (7)	23.3(11)	138	22.5(10)	24.6(11)	207	7.2 (33)

- (1) Reporting period is January 1 through December 31.  
(2) Analyses based on one record per child.  
(3) Excludes records with unknown data or errors.  
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.  
(5) Infants born during the reporting period included in the analysis.  
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.  
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.  
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.  
\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**2002 Pediatric Nutrition Surveillance** (1)  
**Kentucky**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years** (2)

<b>Comparison of Growth and Anemia Indicators by County</b> (3,4)												
<b>County</b>	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low		High	Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Greenup	248	11.7 (78)	7.7 (52)	942	3.3 (3)	8.4(11)	12.3(10)	418	17.9(57)	12.2(16)	662	6.8 (28)
Hancock	58	* *	* *	249	6.8(50)	4.8(80)	16.1(45)	127	16.5(34)	14.2(34)	196	4.1 (6)
Hardin	921	10.2 (58)	7.4 (45)	3,338	7.4(58)	2.4(15)	15.2(35)	1,517	22.2(10)	15.5(41)	2,469	7.3 (35)
Harlan	371	8.9 (37)	10.8 (82)	1,591	4.4(15)	7.4(11)	19.0(86)	812	14.9(21)	20.1(86)	1,129	6.1 (21)
Harrison	123	13.8 (89)	8.1 (56)	509	4.1(11)	6.1(10)	9.6 (2)	245	11.4 (7)	6.9 (2)	384	13.8 (90)
Hart	184	7.6 (16)	8.7 (64)	613	6.4(39)	3.6(47)	15.5(37)	270	14.8(20)	17.0(58)	401	5.7 (18)
Henderson	362	10.5 (63)	5.2 (11)	1,120	8.9(84)	4.5(73)	15.9(42)	438	22.1(10)	22.1(10)	271	1.8 (1)
Henry	86	* *	* *	350	16.9(11)	3.4(38)	25.1(11)	141	20.6(89)	22.7(10)	273	10.3 (64)
Hickman	36	* *	* *	121	4.1(11)	5.8(95)	10.7 (4)	54	* *	* *	88	* *
Hopkins	346	14.5 (91)	3.2 (2)	1,429	13.7(11)	1.5 (4)	21.6(10)	664	21.1(91)	18.7(76)	1,087	12.3 (81)
Jackson	131	11.5 (76)	10.7 (81)	627	6.7(46)	1.6 (6)	17.7(71)	301	14.0(15)	15.9(44)	502	3.0 (3)

- (1) Reporting period is January 1 through December 31.  
(2) Analyses based on one record per child.  
(3) Excludes records with unknown data or errors.  
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.  
(5) Infants born during the reporting period included in the analysis.  
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.  
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.  
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.  
\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**2002 Pediatric Nutrition Surveillance** (1)  
**Kentucky**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years** (2)

<b>Comparison of Growth and Anemia Indicators by County</b> (3,4)												
<b>County</b>	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Jefferson	4,703	11.0 (70)	6.6 (25)	14,020	8.4(75)	4.7(76)	15.6(39)	5,441	17.6(52)	16.3(47)	8,921	12.2 (80)
Jessamine	263	5.3 (3)	8.7 (64)	1,013	8.5(77)	2.3(14)	17.5(68)	449	18.9(73)	17.6(64)	742	9.6 (58)
Johnson	233	9.9 (55)	10.3 (80)	998	5.4(23)	5.9(97)	17.8(72)	505	14.9(21)	16.8(53)	769	16.6(104)
Kenton	507	12.6 (85)	8.3 (59)	2,067	8.3(74)	3.6(47)	20.0(91)	886	21.9(99)	23.1(10)	1,591	14.5 (92)
Knott	142	10.6 (64)	4.2 (6)	681	10.7(10)	4.3(68)	20.4(96)	370	11.9 (8)	13.8(30)	570	8.2 (47)
Knox	334	12.6 (85)	6.0 (19)	1,471	6.1(36)	3.2(31)	17.9(73)	740	17.7(56)	20.0(85)	1,116	11.6 (75)
Larue	110	5.5 (4)	7.3 (44)	413	9.0(88)	4.4(72)	20.6(99)	189	17.5(49)	18.0(67)	323	9.0 (52)
Laurel	481	9.8 (51)	9.1 (72)	1,591	6.9(53)	5.2(88)	14.5(27)	635	20.6(89)	19.2(80)	1,019	14.6 (93)
Lawrence	154	11.7 (78)	4.5 (7)	575	5.2(22)	4.9(82)	20.3(94)	262	18.7(68)	21.0(92)	402	7.5 (37)
Lee	63	* *	* *	338	9.2(92)	0.9 (2)	26.6(11)	174	24.1(11)	21.8(97)	280	15.4 (98)
Leslie	128	7.8 (19)	11.7 (87)	621	3.9 (8)	3.1(27)	22.7(10)	343	19.2(81)	23.0(10)	513	8.4 (49)

- (1) Reporting period is January 1 through December 31.  
(2) Analyses based on one record per child.  
(3) Excludes records with unknown data or errors.  
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.  
(5) Infants born during the reporting period included in the analysis.  
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.  
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.  
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.  
\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**2002 Pediatric Nutrition Surveillance** (1)  
**Kentucky**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years** (2)

<b>Comparison of Growth and Anemia Indicators by County</b> (3,4)												
<b>County</b>	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Letcher	204	7.8 (19)	6.9 (34)	829	11.9(11)	3.1(27)	25.8(11)	381	17.3(46)	21.3(93)	638	4.1 (6)
Lewis	125	10.4 (62)	6.4 (23)	480	7.1(55)	2.9(24)	16.7(57)	218	22.0(10)	20.2(87)	315	4.8 (13)
Lincoln	227	11.9 (80)	8.4 (61)	834	7.6(63)	5.0(86)	18.5(81)	369	17.1(41)	19.8(84)	626	7.7 (41)
Livingston	44	* *	* *	193	4.7(18)	7.8(11)	16.1(45)	83	* *	* *	153	6.5 (25)
Logan	199	7.5 (14)	3.5 (3)	691	6.5(41)	3.3(33)	16.2(49)	257	21.8(97)	16.3(47)	465	17.6(108)
Lyon	38	* *	* *	162	9.3(94)	4.3(68)	18.5(81)	74	* *	* *	123	6.5 (25)
Madison	524	9.0 (39)	9.5 (76)	1,736	8.5(77)	5.6(92)	13.7(21)	699	21.9(99)	19.2(80)	1,219	11.3 (70)
Magoffin	147	8.2 (24)	8.8 (67)	703	10.7(10)	4.0(63)	26.3(11)	363	16.0(30)	24.0(10)	511	20.5(112)
Marion	159	11.9 (80)	6.9 (34)	660	11.5(11)	3.8(56)	22.6(10)	308	20.1(86)	17.2(60)	478	10.5 (65)
Marshall	228	8.3 (26)	7.5 (48)	822	11.2(10)	6.7(10)	14.5(27)	365	12.6(12)	7.1 (3)	558	7.7 (41)
Martin	149	9.4 (46)	4.7 (8)	683	5.9(31)	3.1(27)	17.6(70)	363	17.6(52)	18.5(75)	544	5.5 (16)

- (1) Reporting period is January 1 through December 31.  
(2) Analyses based on one record per child.  
(3) Excludes records with unknown data or errors.  
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.  
(5) Infants born during the reporting period included in the analysis.  
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.  
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.  
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.  
\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**2002 Pediatric Nutrition Surveillance** (1)  
**Kentucky**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years** (2)

<b>Comparison of Growth and Anemia Indicators by County</b> (3,4)												
<b>County</b>	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low		High	Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Mason	169	5.9 (5)	8.3 (59)	573	7.9(67)	7.5(11)	16.2(49)	254	16.9(37)	13.4(26)	389	11.8 (78)
McCracken	537	12.5 (83)	7.8 (53)	1,840	8.2(71)	4.2(67)	21.5(10)	815	17.4(48)	16.8(53)	1,396	18.1 (109)
McCreary	184	10.9 (67)	10.9 (83)	855	9.1(89)	3.4(38)	14.5(27)	420	21.2(92)	13.8(30)	663	8.1 (46)
McLean	99	* *	* *	428	11.2(10)	2.8(19)	21.0(10)	202	21.8(97)	24.3(11)	329	4.0 (5)
Meade	172	8.7 (34)	3.5 (3)	677	10.8(10)	3.4(38)	19.1(87)	310	25.2(11)	18.4(72)	518	13.7 (88)
Menifee	62	* *	* *	280	4.6(17)	3.9(61)	12.9(13)	140	18.6(65)	13.6(27)	189	10.1 (62)
Mercer	165	11.5 (76)	5.5 (15)	575	8.7(81)	4.3(68)	16.0(44)	249	23.7(11)	18.9(77)	410	7.8 (43)
Metcalfe	92	* *	* *	333	5.7(30)	3.6(47)	13.2(16)	149	21.5(96)	16.8(53)	225	9.3 (56)
Monroe	99	* *	* *	398	8.0(69)	3.8(56)	13.6(19)	185	15.1(25)	11.9(13)	286	11.5 (72)
Montgomery	229	9.6 (47)	7.0 (38)	831	6.7(46)	3.2(31)	14.2(22)	379	13.2(13)	12.9(20)	622	9.2 (54)
Morgan	106	8.5 (29)	11.3 (85)	544	6.1(36)	3.7(53)	23.2(11)	281	19.2(81)	22.4(10)	444	20.3(111)

- (1) Reporting period is January 1 through December 31.  
(2) Analyses based on one record per child.  
(3) Excludes records with unknown data or errors.  
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.  
(5) Infants born during the reporting period included in the analysis.  
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.  
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.  
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.  
\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**2002 Pediatric Nutrition Surveillance** (1)  
**Kentucky**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years** (2)

<b>Comparison of Growth and Anemia Indicators by County</b> (3,4)												
<b>County</b>	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low		High	Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Muhlenberg	256	9.8 (51)	9.0 (70)	1,004	4.3(13)	7.2(10)	12.1 (7)	466	14.6(18)	13.7(29)	661	9.5 (57)
Nelson	271	10.0 (56)	8.1 (56)	996	8.7(81)	3.5(44)	18.8(85)	445	17.5(49)	14.6(36)	715	7.0 (30)
Nicholas	78	* *	* *	274	4.7(18)	10.6(12)	11.7 (6)	128	9.4 (3)	11.7 (9)	202	12.4 (82)
Ohio	209	9.6 (47)	11.0 (84)	800	6.0(33)	2.8(19)	15.5(37)	389	18.3(62)	16.5(51)	601	10.5 (65)
Oldham	150	10.7 (66)	6.7 (30)	558	12.7(11)	2.9(24)	18.3(79)	241	22.0(10)	16.6(52)	373	15.8(101)
Owen	86	* *	* *	271	3.0 (1)	8.1(11)	14.4(26)	123	17.1(41)	15.4(40)	191	6.3 (24)
Owsley	51	* *	* *	259	9.7(98)	1.5 (4)	29.3(11)	137	21.2(92)	24.1(11)	219	21.5(114)
Pendleton	102	9.8 (51)	6.9 (34)	397	9.3(94)	4.0(63)	13.6(19)	170	18.8(71)	14.1(32)	297	7.4 (36)
Perry	357	9.0 (39)	5.9 (17)	1,109	10.6(10)	2.0 (9)	33.4(11)	512	18.2(61)	21.9(98)	813	8.0 (45)
Pike	554	10.3 (59)	7.8 (53)	1,960	3.5 (5)	8.7(11)	17.2(65)	888	18.6(65)	21.3(93)	1,363	6.8 (28)
Powell	173	7.5 (14)	5.2 (11)	659	3.9 (8)	3.3(33)	15.3(36)	330	15.2(26)	15.5(41)	501	6.2 (22)

- (1) Reporting period is January 1 through December 31.  
(2) Analyses based on one record per child.  
(3) Excludes records with unknown data or errors.  
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.  
(5) Infants born during the reporting period included in the analysis.  
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.  
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.  
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.  
\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**2002 Pediatric Nutrition Surveillance** (1)  
**Kentucky**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years** (2)

<b>Comparison of Growth and Anemia Indicators by County</b> (3,4)												
<b>County</b>	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low		High	Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Pulaski	533	8.6 (31)	6.8 (33)	1,818	8.9(84)	4.7(76)	12.2 (9)	741	15.8(29)	13.0(21)	1,131	6.5 (25)
Robertson	25	* *	* *	108	13.9(11)	0.9 (2)	18.5(81)	53	* *	* *	85	* *
Rockcastle	183	8.7 (34)	6.6 (25)	611	6.5(41)	7.5(11)	17.3(66)	279	16.8(35)	12.5(18)	416	20.7(113)
Rowan	177	7.3 (13)	15.3 (91)	681	3.7 (6)	3.4(38)	15.0(33)	298	18.5(63)	13.1(23)	445	16.9(106)
Russell	157	8.9 (37)	8.9 (68)	580	8.6(79)	2.1(11)	17.1(63)	241	19.1(77)	20.7(90)	368	10.1 (62)
Scott	230	8.7 (34)	7.0 (38)	684	8.8(83)	5.7(94)	17.5(68)	256	11.3 (6)	12.1(14)	464	11.4 (71)
Shelby	231	10.0 (56)	5.6 (16)	738	22.1(12)	1.9 (7)	37.1(12)	265	21.9(99)	32.8(11)	515	13.6 (87)
Simpson	124	12.9 (88)	4.8 (10)	426	9.6(97)	2.8(19)	16.4(53)	165	29.1(11)	18.2(69)	275	22.2(115)
Spencer	69	* *	* *	258	10.9(10)	3.1(27)	18.2(77)	112	17.0(39)	16.1(46)	188	10.6 (67)
Taylor	195	3.6 (2)	8.7 (64)	759	6.6(45)	2.8(19)	16.9(61)	356	21.3(95)	17.4(62)	492	4.7 (12)
Todd	101	5.9 (5)	6.9 (34)	448	8.0(69)	4.0(63)	16.5(56)	205	14.1(16)	8.8 (6)	351	17.4(107)

- (1) Reporting period is January 1 through December 31.  
(2) Analyses based on one record per child.  
(3) Excludes records with unknown data or errors.  
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.  
(5) Infants born during the reporting period included in the analysis.  
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.  
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.  
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.  
\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



**Table 6B**  
**2002 Pediatric Nutrition Surveillance** (1)  
**Kentucky**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years** (2)

<b>Comparison of Growth and Anemia Indicators by County</b> (3,4)												
<b>County</b>	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low		High	Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Trigg	75	* *	* *	290	5.9(31)	4.5(73)	13.1(14)	131	16.8(35)	12.2(16)	199	7.5 (37)
Trimble	57	* *	* *	232	5.6(26)	8.6(11)	7.3 (1)	111	10.8 (5)	9.0 (7)	184	4.3 (9)
Union	119	10.9 (67)	2.5 (1)	474	12.0(11)	2.5(17)	16.9(61)	205	17.6(52)	11.7 (9)	369	7.6 (39)
Warren	728	11.0 (70)	6.6 (25)	2,202	7.2(56)	3.4(38)	15.7(40)	836	17.5(49)	16.3(47)	1,337	13.4 (86)
Washington	88	* *	* *	369	9.8(99)	3.5(44)	19.8(89)	165	18.8(71)	12.1(14)	283	12.4 (82)
Wayne	216	12.0 (82)	7.4 (45)	844	4.3(13)	3.0(26)	16.8(59)	402	17.9(57)	21.4(95)	575	9.6 (58)
Webster	113	18.6 (92)	5.3 (13)	414	10.6(10)	3.9(61)	16.4(53)	177	12.4(11)	16.9(57)	304	15.1 (97)
Whitley	493	7.7 (17)	9.9 (78)	2,133	7.4(58)	4.5(73)	17.9(73)	1,021	18.7(68)	19.1(79)	1,635	6.2 (22)
Wolfe	100	8.0 (21)	7.0 (38)	431	10.9(10)	0.5 (1)	25.3(11)	218	19.7(85)	23.4(10)	340	8.5 (51)
Woodford	132	10.6 (64)	9.8 (77)	515	7.8(66)	3.7(53)	16.1(45)	205	14.6(18)	11.7 (9)	349	12.0 (79)

- (1) Reporting period is January 1 through December 31.  
(2) Analyses based on one record per child.  
(3) Excludes records with unknown data or errors.  
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.  
(5) Infants born during the reporting period included in the analysis.  
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.  
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.  
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.  
\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**2002 Pediatric Nutrition Surveillance** (1)  
**Kentucky**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years** (2)

<b>Comparison of Growth and Anemia Indicators by County</b> (3,4)												
<b>County</b>	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% >95th (Rank)	Number
<b>Kentucky</b>	<b>32,542</b>	<b>10.0</b>	<b>7.5</b>	<b>114,943</b>	<b>7.6</b>	<b>4.6</b>	<b>16.8</b>	<b>50,285</b>	<b>17.7</b>	<b>16.8</b>	<b>79,270</b>	<b>11.6</b>
<b>Nation (Prior Year)</b>	<b>1,260,117</b>	<b>9.0</b>	<b>7.9</b>	<b>4,846,643</b>	<b>6.4</b>	<b>5.4</b>	<b>13.1</b>	<b>2,031,241</b>	<b>15.3</b>	<b>13.4</b>	<b>3,265,122</b>	<b>13.3</b>

- (1) Reporting period is January 1 through December 31.
- (2) Analyses based on one record per child.
- (3) Excludes records with unknown data or errors.
- (4) Rank compares this county's rate to other counties. Rank 1 = best rate.
- (5) Infants born during the reporting period included in the analysis.
- (6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
- (7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
- (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
- \* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 7B**  
**2002 Pediatric Nutrition Surveillance** (1)  
**Kentucky**

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County** (2)  
**Children Aged < 5 Years** (3)

<b>Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County</b> (4,5)						
<b>County</b>	Breastfeeding (6)					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Adair	135	11.1 (56)	49	* *	98	* *
Allen	138	24.6 (7)	51	* *	95	* *
Anderson	82	* *	43	* *	57	* *
Ballard	62	* *	26	* *	58	* *
Barren	252	15.1 (40)	104	1.9 (8)	188	0.0 (1)
Bath	100	8.0 (81)	35	* *	79	* *
Bell	343	10.8 (63)	152	0.0 (21)	302	0.0 (1)
Boone	487	12.7 (50)	235	3.4 (3)	331	0.0 (1)
Bourbon	119	17.6 (21)	63	* *	108	0.0 (1)
Boyd	425	14.1 (41)	127	0.8 (18)	238	0.0 (1)
Boyle	187	9.1 (73)	53	* *	104	0.0 (1)
Bracken	70	* *	20	* *	34	* *
Breathitt	154	9.7 (68)	52	* *	103	0.0 (1)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 7B**  
**2002 Pediatric Nutrition Surveillance** (1)  
**Kentucky**

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County** (2)  
**Children Aged < 5 Years** (3)

<b>Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County</b> (4,5)						
<b>County</b>	Breastfeeding (6)					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Breckinrid	114	16.7 (26)	41	* *	87	* *
Bullitt	374	16.6 (28)	102	0.0 (21)	213	0.0 (1)
Butler	115	9.6 (71)	41	* *	80	* *
Caldwell	108	8.3 (78)	55	* *	91	* *
Calloway	243	28.4 (4)	94	* *	173	0.0 (1)
Campbell	404	7.7 (84)	160	1.3 (12)	237	0.0 (1)
Carlisle	35	* *	21	* *	38	* *
Carroll	107	14.0 (43)	40	* *	82	* *
Carter	301	17.6 (21)	109	0.9 (16)	210	0.0 (1)
Casey	136	16.2 (34)	43	* *	85	* *
Christian	1,287	21.4 (12)	507	4.1 (2)	646	0.0 (1)
Clark	255	14.1 (41)	94	* *	140	0.0 (1)
Clay	258	8.9 (77)	102	0.0 (21)	217	0.0 (1)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 7B**  
**2002 Pediatric Nutrition Surveillance** (1)  
**Kentucky**

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County** (2)  
**Children Aged < 5 Years** (3)

<b>Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County</b> (4,5)						
<b>County</b>	Breastfeeding (6)					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Clinton	111	5.4 (88)	41	* *	87	* *
Crittenden	62	* *	22	* *	46	* *
Cumberland	66	* *	30	* *	55	* *
Daviess	722	13.3 (46)	251	2.0 (7)	433	0.0 (1)
Edmonson	102	19.6 (14)	38	* *	44	* *
Elliott	56	* *	30	* *	48	* *
Estill	158	17.1 (23)	45	* *	91	* *
Fayette	1,560	39.2 (1)	525	4.6 (1)	932	0.0 (1)
Fleming	127	11.0 (57)	58	* *	102	0.0 (1)
Floyd	440	4.5 (89)	177	0.0 (21)	322	0.0 (1)
Franklin	308	26.3 (5)	131	2.3 (5)	164	0.0 (1)
Fulton	95	* *	40	* *	64	* *
Gallatin	74	* *	20	* *	40	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 7B**  
**2002 Pediatric Nutrition Surveillance** (1)  
**Kentucky**

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County** (2)  
**Children Aged < 5 Years** (3)

<b>Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County</b> (4,5)						
<b>County</b>	Breastfeeding (6)					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Garrard	124	17.7 (20)	50	* *	114	0.0 (1)
Grant	210	9.0 (76)	116	0.0 (21)	196	0.0 (1)
Graves	314	22.3 (10)	95	* *	215	0.0 (1)
Grayson	203	29.6 (3)	67	* *	150	0.0 (1)
Green	93	* *	29	* *	61	* *
Greenup	250	18.0 (19)	103	1.0 (15)	143	0.0 (1)
Hancock	58	* *	30	* *	35	* *
Hardin	926	16.5 (29)	343	0.6 (19)	495	0.0 (1)
Harlan	371	12.9 (48)	161	0.0 (21)	288	0.0 (1)
Harrison	124	16.9 (24)	58	* *	99	* *
Hart	184	10.9 (60)	53	* *	91	* *
Henderson	362	10.2 (66)	115	0.0 (21)	156	0.0 (1)
Henry	86	* *	38	* *	73	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 7B  
2002 Pediatric Nutrition Surveillance (1)  
Kentucky**

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County (2)  
Children Aged < 5 Years (3)**

<b>Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County (4,5)</b>						
<b>County</b>	<b>Breastfeeding (6)</b>					
	<b>Number</b>	<b>% Ever Breastfed</b>	<b>Number</b>	<b>% Breastfed At Least 6 Months</b>	<b>Number</b>	<b>% Breastfed At Least 12 Months</b>
		<b>(Rank)</b>		<b>(Rank)</b>		<b>(Rank)</b>
Hickman	36	* *	13	* *	26	* *
Hopkins	353	11.0 (57)	125	0.0 (21)	222	0.0 (1)
Jackson	131	10.7 (64)	62	* *	140	0.0 (1)
Jefferson	4,729	18.9 (16)	1,535	1.4 (11)	2,404	0.0 (1)
Jessamine	264	16.7 (26)	110	1.8 (9)	166	0.0 (1)
Johnson	233	8.2 (79)	101	0.0 (21)	183	0.0 (1)
Kenton	511	7.8 (83)	247	1.2 (13)	364	0.0 (1)
Knott	142	4.2 (90)	65	* *	101	0.0 (1)
Knox	345	8.1 (80)	136	0.0 (21)	257	0.0 (1)
Larue	111	21.6 (11)	54	* *	73	* *
Laurel	481	10.4 (65)	163	1.2 (13)	262	0.0 (1)
Lawrence	154	9.7 (68)	59	* *	114	0.0 (1)
Lee	64	* *	32	* *	59	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 7B**  
**2002 Pediatric Nutrition Surveillance** (1)  
**Kentucky**

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County** (2)  
**Children Aged < 5 Years** (3)

<b>Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County</b> (4,5)						
<b>County</b>	Breastfeeding (6)					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Leslie	128	10.9 (60)	51	* *	97	* *
Letcher	209	12.0 (53)	97	* *	119	0.0 (1)
Lewis	125	16.8 (25)	53	* *	104	0.0 (1)
Lincoln	227	13.7 (45)	90	* *	126	0.0 (1)
Livingston	44	* *	19	* *	41	* *
Logan	202	3.5 (91)	80	* *	144	0.0 (1)
Lyon	38	* *	20	* *	28	* *
Madison	525	16.2 (34)	193	2.6 (4)	295	0.0 (1)
Magoffin	147	2.7 (92)	80	* *	133	0.0 (1)
Marion	159	16.4 (31)	57	* *	106	0.0 (1)
Marshall	230	20.0 (13)	88	* *	153	0.0 (1)
Martin	153	2.6 (93)	50	* *	121	0.0 (1)
Mason	169	13.0 (47)	57	* *	108	0.0 (1)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



**Table 7B**  
**2002 Pediatric Nutrition Surveillance** (1)  
**Kentucky**

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County** (2)  
**Children Aged < 5 Years** (3)

<b>Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County</b> (4,5)						
<b>County</b>	Breastfeeding (6)					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
McCracken	538	13.9 (44)	196	0.0 (21)	385	0.0 (1)
McCreary	187	9.1 (73)	92	* *	165	0.0 (1)
McLean	99	* *	42	* *	77	* *
Meade	172	25.6 (6)	59	* *	103	0.0 (1)
Menifee	62	* *	21	* *	46	* *
Mercer	165	10.9 (60)	54	* *	89	* *
Metcalfe	92	* *	43	* *	57	* *
Monroe	100	16.0 (36)	48	* *	59	* *
Montgomery	231	11.7 (55)	84	* *	126	0.0 (1)
Morgan	109	11.0 (57)	61	* *	104	0.0 (1)
Muhlenberg	256	9.4 (72)	99	* *	195	0.0 (1)
Nelson	271	15.9 (37)	104	0.0 (21)	148	0.0 (1)
Nicholas	78	* *	27	* *	39	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 7B**  
**2002 Pediatric Nutrition Surveillance** (1)  
**Kentucky**

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County** (2)  
**Children Aged < 5 Years** (3)

<b>Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County</b> (4,5)						
<b>County</b>	<b>Breastfeeding</b> (6)					
	<b>Number</b>	<b>% Ever Breastfed  (Rank)</b>	<b>Number</b>	<b>% Breastfed At Least 6 Months (Rank)</b>	<b>Number</b>	<b>% Breastfed At Least 12 Months (Rank)</b>
Ohio	210	18.1 (18)	78	* *	131	0.0 (1)
Oldham	150	24.0 (8)	61	* *	116	0.0 (1)
Owen	86	* *	26	* *	40	* *
Owsley	51	* *	36	* *	54	* *
Pendleton	102	22.5 (9)	48	* *	86	* *
Perry	361	5.8 (87)	114	0.9 (16)	201	0.0 (1)
Pike	556	12.9 (48)	214	0.5 (20)	330	0.0 (1)
Powell	173	6.9 (85)	61	* *	112	0.0 (1)
Pulaski	537	19.2 (15)	229	1.7 (10)	352	0.0 (1)
Robertson	25	* *	16	* *	24	* *
Rockcastle	184	12.0 (53)	50	* *	84	* *
Rowan	177	7.9 (82)	60	* *	105	0.0 (1)
Russell	158	15.8 (38)	62	* *	124	0.0 (1)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 7B**  
**2002 Pediatric Nutrition Surveillance** (1)  
**Kentucky**

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County** (2)  
**Children Aged < 5 Years** (3)

<b>Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County</b> (4,5)						
<b>County</b>	Breastfeeding (6)					
	Number	% Ever Breastfed  (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Scott	230	16.5 (29)	77	* *	131	0.0 (1)
Shelby	231	9.1 (73)	85	* *	155	0.0 (1)
Simpson	129	16.3 (33)	36	* *	81	* *
Spencer	69	* *	31	* *	57	* *
Taylor	195	16.4 (31)	73	* *	137	0.0 (1)
Todd	101	18.8 (17)	46	* *	97	* *
Trigg	75	* *	31	* *	50	* *
Trimble	57	* *	21	* *	48	* *
Union	120	12.5 (51)	55	* *	88	* *
Warren	752	15.8 (38)	231	2.2 (6)	389	0.0 (1)
Washington	88	* *	33	* *	72	* *
Wayne	223	9.9 (67)	61	* *	135	0.0 (1)
Webster	113	9.7 (68)	48	* *	58	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 7B**  
**2002 Pediatric Nutrition Surveillance** (1)  
**Kentucky**

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County** (2)  
**Children Aged < 5 Years** (3)

<b>Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County</b> (4,5)						
<b>County</b>	Breastfeeding (6)					
	Number	% Ever Breastfed  (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Whitley	493	12.4 (52)	234	0.0 (21)	355	0.0 (1)
Wolfe	100	6.0 (86)	53	* *	85	* *
Woodford	132	34.8 (2)	63	* *	112	0.0 (1)
Unknown Counties	1,175	1.9 (94)	4	* *	20	* *
<b>Kentucky</b>	<b>32,800</b>	<b>15.6</b>	<b>11,889</b>	<b>1.3</b>	<b>20,006</b>	<b>0.0</b>
<b>Nation (Prior Year)</b>	<b>874,598</b>	<b>51.1</b>	<b>493,552</b>	<b>20.3</b>	<b>484,228</b>	<b>13.3</b>

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**2000 - 2002 Pediatric Nutrition Surveillance<sup>(1)</sup>**  
**Kentucky**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years <sup>(2)</sup>**

<b>Comparison of Growth and Anemia Indicators by County</b> <span style="float: right;">(3,4)</span>												
<b>County</b>	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Adair	423	7.6 (21)	7.8 (49)	1,453	4.5(26)	7.0(10)	11.1 (6)	627	11.5 (8)	11.6(13)	871	6.0 (15)
Allen	426	12.2(111)	10.1 (99)	1,747	6.6(79)	4.2(38)	14.9(57)	809	19.9(10)	15.9(70)	1,125	10.0 (60)
Anderson	298	7.4 (18)	8.7 (73)	1,072	7.5(98)	5.0(59)	17.2(91)	476	21.0(11)	18.7(99)	812	7.1 (27)
Ballard	199	9.0 (48)	8.5 (67)	766	7.2(95)	6.8(99)	13.7(39)	354	13.0(15)	11.3(11)	587	6.3 (20)
Barren	856	8.8 (40)	8.8 (79)	2,956	5.6(53)	4.7(54)	15.1(61)	1,209	16.1(49)	15.9(70)	1,909	9.2 (52)
Bath	312	5.8 (6)	14.1(117)	1,346	9.1(11)	4.3(43)	19.2(10)	658	16.6(59)	17.5(89)	855	10.2 (62)
Bell	1,069	9.6 (65)	8.1 (59)	4,759	4.9(35)	3.8(23)	16.8(87)	2,433	15.3(36)	15.3(58)	3,389	7.0 (26)
Boone	1,458	8.9 (43)	9.4 (91)	5,209	6.8(90)	5.1(63)	14.4(50)	2,157	17.2(72)	13.6(35)	3,710	13.3 (83)
Bourbon	434	11.5(102)	9.0 (82)	1,451	5.7(55)	2.8 (9)	17.8(96)	604	21.7(11)	17.4(87)	1,024	29.7(120)
Boyd	1,212	10.3 (78)	8.5 (67)	4,132	6.7(85)	9.6(11)	12.4(20)	1,792	14.9(31)	13.8(40)	2,716	12.7 (79)
Boyle	535	11.6(105)	5.6 (8)	1,635	5.7(55)	5.5(74)	13.7(39)	636	15.7(43)	15.4(60)	1,001	9.5 (55)

- (1) Reporting period is January 1 through December 31.  
(2) Analyses based on one record per child.  
(3) Excludes records with unknown data or errors.  
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.  
(5) Infants born during the reporting period included in the analysis.  
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.  
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.  
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.  
\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**2000 - 2002 Pediatric Nutrition Surveillance<sup>(1)</sup>**  
**Kentucky**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years <sup>(2)</sup>**

<b>Comparison of Growth and Anemia Indicators by County</b> <span style="float: right;">(3,4)</span>												
<b>County</b>	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Bracken	197	12.2 (111)	7.1 (31)	646	7.3(97)	7.0(10)	19.2(10)	264	17.0(67)	22.0(11)	431	6.0 (15)
Breathitt	448	9.2 (55)	8.7 (73)	2,110	4.3(21)	4.0(30)	16.7(85)	1,188	13.0(15)	14.7(54)	1,712	6.9 (23)
Breckinrid	347	8.4 (31)	10.1 (99)	1,415	6.6(79)	3.7(20)	13.1(26)	642	17.3(74)	11.2(10)	1,053	3.9 (3)
Bullitt	1,071	8.1 (27)	7.3 (35)	3,130	6.4(73)	3.9(27)	14.5(52)	1,174	16.7(62)	14.0(43)	1,816	19.4(114)
Butler	348	8.6 (34)	8.0 (56)	1,332	4.1(19)	4.2(38)	17.9(97)	598	17.7(84)	21.4(11)	894	5.9 (14)
Caldwell	321	5.9 (7)	8.7 (73)	1,206	6.7(85)	8.1(11)	13.8(43)	508	15.2(34)	14.6(51)	896	15.6(102)
Calloway	725	8.1 (27)	10.9(109)	2,655	4.6(28)	4.3(43)	13.4(30)	1,134	15.3(36)	15.5(64)	1,706	16.9(107)
Campbell	1,082	10.4 (82)	7.6 (43)	3,601	6.2(68)	4.1(33)	15.7(69)	1,523	19.6(10)	19.6(10)	2,404	13.3 (83)
Carlisle	133	6.8 (11)	12.0(114)	581	5.0(38)	4.1(33)	20.7(11)	295	19.3(10)	26.1(11)	407	21.9(117)
Carroll	354	9.6 (65)	6.5 (20)	1,216	9.7(11)	6.7(97)	16.2(76)	505	17.6(81)	17.8(91)	882	14.6 (96)
Carter	913	7.8 (25)	12.0(114)	3,668	3.5(10)	8.7(11)	11.5(10)	1,741	15.0(32)	11.8(17)	2,511	8.4 (47)

- (1) Reporting period is January 1 through December 31.  
(2) Analyses based on one record per child.  
(3) Excludes records with unknown data or errors.  
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.  
(5) Infants born during the reporting period included in the analysis.  
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.  
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.  
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.  
\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**2000 - 2002 Pediatric Nutrition Surveillance<sup>(1)</sup>**  
**Kentucky**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years <sup>(2)</sup>**

<b>Comparison of Growth and Anemia Indicators by County</b> <span style="float: right;">(3,4)</span>												
<b>County</b>	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Casey	379	9.8 (69)	5.8 (12)	1,576	5.5(52)	5.6(76)	12.1(16)	737	16.4(53)	13.0(30)	988	7.5 (36)
Christian	4,037	8.9 (43)	9.1 (86)	15,040	5.1(44)	7.5(10)	10.0 (4)	6,398	12.9(12)	9.2 (6)	10,142	23.0(118)
Clark	744	11.8(106)	8.2 (62)	2,430	8.1(10)	3.4(17)	16.1(75)	964	19.9(10)	15.4(60)	1,658	12.0 (73)
Clay	804	11.8(106)	6.8 (26)	3,435	4.8(33)	7.9(11)	16.7(85)	1,751	15.6(41)	16.5(80)	2,569	9.3 (54)
Clinton	352	9.1 (51)	8.8 (79)	1,443	4.0(18)	6.2(90)	15.0(58)	666	12.3(11)	11.9(20)	972	6.2 (17)
Crittenden	197	5.6 (4)	10.7(106)	803	5.1(44)	6.4(92)	12.1(16)	362	14.4(26)	9.7 (7)	593	12.6 (76)
Cumberland	211	3.8 (1)	9.0 (82)	884	3.7(15)	6.3(91)	13.0(25)	411	15.3(36)	14.6(51)	444	14.9 (98)
Daviess	2,340	9.4 (61)	5.9 (13)	7,903	7.8(10)	2.2 (3)	17.6(93)	3,313	16.5(55)	15.6(65)	5,638	12.6 (76)
Edmonson	312	9.9 (71)	8.3 (66)	1,098	3.6(13)	6.1(85)	11.7(12)	449	14.7(28)	13.8(40)	707	5.2 (7)
Elliott	243	6.6 (10)	12.3(116)	959	2.8 (2)	6.0(82)	12.3(18)	458	14.4(26)	14.2(45)	671	5.7 (12)
Estill	441	9.3 (58)	10.0 (97)	1,820	2.8 (2)	5.7(77)	13.4(30)	873	17.3(74)	16.4(79)	1,380	7.3 (31)

- (1) Reporting period is January 1 through December 31.  
(2) Analyses based on one record per child.  
(3) Excludes records with unknown data or errors.  
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.  
(5) Infants born during the reporting period included in the analysis.  
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.  
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.  
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.  
\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**2000 - 2002 Pediatric Nutrition Surveillance<sup>(1)</sup>**  
**Kentucky**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years <sup>(2)</sup>**

<b>Comparison of Growth and Anemia Indicators by County</b> <span style="float: right;">(3,4)</span>												
<b>County</b>	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Fayette	4,858	10.7 (89)	7.3 (35)	13,820	5.0(38)	5.1(63)	12.9(22)	5,045	17.4(76)	14.6(51)	8,358	16.5(105)
Fleming	410	7.6 (21)	9.8 (96)	1,817	4.6(28)	6.5(95)	13.3(29)	862	15.7(43)	13.6(35)	1,415	16.5(105)
Floyd	1,321	8.7 (38)	8.5 (67)	5,264	3.4 (8)	10.4(12)	12.0(15)	2,512	10.1 (2)	11.9(20)	3,793	8.7 (49)
Franklin	969	7.6 (21)	6.9 (27)	3,034	8.0(10)	5.7(77)	17.7(94)	1,135	18.2(89)	18.4(95)	2,092	11.5 (69)
Fulton	291	11.3(100)	6.9 (27)	1,098	6.8(90)	4.1(33)	19.9(11)	521	17.1(69)	20.3(11)	762	14.4 (94)
Gallatin	216	10.6 (86)	7.4 (38)	796	5.7(55)	5.0(59)	12.9(22)	360	13.1(18)	14.4(48)	536	13.4 (87)
Garrard	343	10.5 (83)	9.0 (82)	1,536	5.7(55)	3.7(20)	14.8(54)	749	17.1(69)	17.9(93)	1,005	14.8 (97)
Grant	667	11.1 (95)	7.9 (51)	2,986	4.3(21)	7.3(10)	9.3 (1)	1,439	8.8 (1)	7.0 (1)	2,350	6.2 (17)
Graves	896	9.9 (71)	9.4 (91)	3,377	5.7(55)	6.1(85)	14.8(54)	1,553	18.2(89)	15.3(58)	2,261	16.4(104)
Grayson	680	7.4 (18)	8.2 (62)	2,632	10.0(11)	3.1(12)	18.8(10)	1,165	16.6(59)	16.5(80)	1,966	3.0 (1)
Green	270	8.5 (33)	9.6 (94)	956	4.6(28)	3.6(19)	17.7(94)	417	21.8(11)	17.7(90)	602	5.8 (13)

- (1) Reporting period is January 1 through December 31.  
(2) Analyses based on one record per child.  
(3) Excludes records with unknown data or errors.  
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.  
(5) Infants born during the reporting period included in the analysis.  
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.  
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.  
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.  
\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



**Table 6B**  
**2000 - 2002 Pediatric Nutrition Surveillance<sup>(1)</sup>**  
**Kentucky**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years <sup>(2)</sup>**

<b>Comparison of Growth and Anemia Indicators by County</b> <span style="float: right;">(3,4)</span>												
<b>County</b>	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Greenup	780	9.9 (71)	8.1 (59)	2,829	2.9 (4)	8.8(11)	11.7(12)	1,242	15.4(39)	12.2(23)	1,933	8.0 (44)
Hancock	194	10.3 (78)	4.1 (2)	766	5.0(38)	4.2(38)	16.2(76)	349	14.3(24)	16.6(82)	583	7.2 (29)
Hardin	2,794	8.9 (43)	7.4 (38)	10,567	5.8(60)	3.2(13)	13.7(39)	4,640	19.2(10)	15.1(56)	7,848	7.7 (40)
Harlan	1,132	9.0 (48)	10.8(108)	4,983	3.9(16)	8.4(11)	17.0(89)	2,598	16.8(63)	18.4(95)	3,584	7.6 (39)
Harrison	428	12.9(115)	7.2 (32)	1,472	3.5(10)	6.7(97)	9.9 (3)	635	10.2 (3)	7.9 (2)	1,038	12.0 (73)
Hart	500	7.0 (12)	9.2 (89)	1,832	4.6(28)	4.4(47)	14.3(49)	807	15.2(34)	15.7(66)	1,236	7.3 (31)
Henderson	1,032	11.1 (95)	5.1 (4)	3,357	6.4(73)	6.1(85)	13.9(44)	1,307	20.0(11)	19.0(10)	1,249	7.7 (40)
Henry	349	7.2 (15)	6.0 (16)	1,147	11.7(11)	4.7(54)	22.8(11)	445	19.3(10)	23.4(11)	830	15.3 (99)
Hickman	99	* *	* *	394	4.3(21)	5.8(79)	15.2(62)	179	15.6(41)	20.1(10)	305	7.5 (36)
Hopkins	1,156	10.8 (91)	5.4 (5)	4,279	9.9(11)	2.5 (5)	19.9(11)	1,882	19.3(10)	18.5(97)	3,086	13.8 (91)
Jackson	455	8.6 (34)	9.7 (95)	1,823	5.9(63)	4.1(33)	15.8(70)	848	13.0(15)	15.8(68)	1,405	3.3 (2)

- (1) Reporting period is January 1 through December 31.  
(2) Analyses based on one record per child.  
(3) Excludes records with unknown data or errors.  
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.  
(5) Infants born during the reporting period included in the analysis.  
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.  
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.  
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.  
\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**2000 - 2002 Pediatric Nutrition Surveillance<sup>(1)</sup>**  
**Kentucky**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years <sup>(2)</sup>**

<b>Comparison of Growth and Anemia Indicators by County</b> <span style="float: right;">(3,4)</span>												
<b>County</b>	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Jefferson	12,691	10.5 (83)	7.2 (32)	36,056	6.5(75)	6.0(82)	13.6(35)	13,413	15.5(40)	13.6(35)	22,572	13.3 (83)
Jessamine	890	7.0 (12)	9.0 (82)	3,026	6.6(79)	3.2(13)	15.3(66)	1,281	17.6(81)	15.4(60)	2,070	10.7 (65)
Johnson	754	7.0 (12)	10.3(102)	3,002	3.6(13)	8.1(11)	15.0(58)	1,403	13.5(20)	14.4(48)	2,208	17.0(108)
Kenton	2,046	11.5(102)	6.6 (23)	6,746	6.0(65)	5.2(67)	15.0(58)	2,673	19.7(10)	18.9(10)	4,753	12.9 (80)
Knott	459	11.5(102)	6.1 (18)	2,169	8.8(11)	6.0(82)	18.9(10)	1,134	11.7(10)	11.4(12)	1,789	8.8 (50)
Knox	1,031	9.3 (58)	8.2 (62)	4,554	5.2(49)	3.9(27)	16.0(74)	2,244	16.5(55)	16.8(84)	3,528	14.1 (93)
Larue	343	8.7 (38)	6.7 (25)	1,173	7.7(99)	5.3(69)	16.5(81)	510	17.5(78)	14.7(54)	896	10.9 (67)
Laurel	1,513	11.8(106)	9.2 (89)	4,850	5.0(38)	6.5(95)	13.6(35)	1,890	19.6(10)	15.9(70)	3,008	13.3 (83)
Lawrence	484	11.8(106)	7.0 (30)	1,732	4.2(20)	8.0(11)	16.6(82)	767	15.8(45)	16.2(77)	1,161	9.1 (51)
Lee	206	8.3 (30)	11.2(110)	1,026	6.1(67)	1.7 (1)	22.6(11)	528	19.1(10)	21.0(11)	841	18.2(110)
Leslie	405	7.2 (15)	11.9(113)	1,966	2.4 (1)	4.8(57)	18.3(10)	1,067	17.5(78)	19.3(10)	1,580	6.6 (22)

- (1) Reporting period is January 1 through December 31.  
(2) Analyses based on one record per child.  
(3) Excludes records with unknown data or errors.  
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.  
(5) Infants born during the reporting period included in the analysis.  
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.  
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.  
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.  
\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**2000 - 2002 Pediatric Nutrition Surveillance<sup>(1)</sup>**  
**Kentucky**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years <sup>(2)</sup>**

<b>Comparison of Growth and Anemia Indicators by County</b> <span style="float: right;">(3,4)</span>												
<b>County</b>	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Letcher	686	9.2 (55)	4.5 (3)	2,662	7.2(95)	4.1(33)	21.3(11)	1,202	16.8(63)	20.2(10)	1,998	4.0 (4)
Lewis	405	11.1 (95)	9.1 (86)	1,428	6.0(65)	5.3(69)	13.6(35)	615	21.0(11)	17.4(87)	968	5.3 (9)
Lincoln	671	10.7 (89)	7.6 (43)	2,433	5.6(53)	6.1(85)	15.5(68)	1,078	19.6(10)	17.3(86)	1,826	7.8 (42)
Livingston	174	10.3 (78)	7.5 (41)	712	5.1(44)	7.7(10)	16.6(82)	336	16.1(49)	15.8(68)	550	8.2 (45)
Logan	646	9.8 (69)	5.4 (5)	2,059	4.6(28)	3.9(27)	15.3(66)	794	20.2(11)	15.9(70)	1,317	19.3(113)
Lyon	123	5.7 (5)	10.6(104)	535	6.7(85)	5.4(72)	18.5(10)	258	16.3(51)	16.7(83)	418	14.4 (94)
Madison	1,626	9.6 (65)	10.6(104)	5,455	6.5(75)	7.6(10)	11.7(12)	2,195	18.7(95)	15.9(70)	3,846	11.9 (72)
Magoffin	475	8.6 (34)	10.3(102)	2,201	6.8(90)	5.2(67)	19.8(11)	1,103	15.9(46)	17.1(85)	1,545	19.0(111)
Marion	464	10.6 (86)	7.5 (41)	1,980	8.7(10)	4.7(54)	18.8(10)	899	16.9(66)	15.1(56)	1,413	9.8 (58)
Marshall	660	9.1 (51)	7.9 (51)	2,437	6.6(79)	6.9(10)	13.7(39)	1,055	10.4 (4)	8.0 (3)	1,766	7.2 (29)
Martin	475	11.8(106)	5.9 (13)	2,154	4.9(35)	5.3(69)	16.3(79)	1,118	18.2(89)	17.8(91)	1,729	6.5 (21)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**2000 - 2002 Pediatric Nutrition Surveillance<sup>(1)</sup>**  
**Kentucky**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years<sup>(2)</sup>**

<b>Comparison of Growth and Anemia Indicators by County</b> <span style="float: right;">(3,4)</span>												
<b>County</b>	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low		High	Short Stature (6)			Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Mason	498	7.2 (15)	8.0 (56)	1,682	8.3(10)	7.3(10)	14.7(53)	714	14.3(24)	11.8(17)	1,164	13.5 (88)
McCracken	1,594	11.1 (95)	8.1 (59)	5,560	6.6(79)	5.4(72)	18.1(98)	2,408	16.5(55)	13.8(40)	4,131	19.1 (112)
McCreary	607	11.0 (93)	10.7(106)	2,678	5.0(38)	5.8(79)	13.4(30)	1,314	12.9(12)	11.6(13)	2,077	7.1 (27)
McLean	316	11.4(101)	5.7 (11)	1,312	7.9(10)	3.3(15)	18.9(10)	597	18.8(97)	22.1(11)	1,011	5.2 (7)
Meade	557	8.6 (34)	7.9 (51)	2,035	8.1(10)	4.3(43)	15.9(73)	917	17.6(81)	15.7(66)	1,527	19.4(114)
Menifee	206	9.2 (55)	7.3 (35)	895	4.8(33)	5.5(74)	11.4 (9)	425	17.4(76)	13.2(32)	580	10.7 (65)
Mercer	486	9.5 (63)	6.6 (23)	1,770	6.7(85)	3.8(23)	13.6(35)	786	18.2(89)	15.4(60)	1,299	8.4 (47)
Metcalfe	291	12.4(113)	11.3(112)	1,037	3.9(16)	4.5(50)	9.8 (2)	436	16.5(55)	13.1(31)	668	9.6 (56)
Monroe	332	5.1 (2)	8.7 (73)	1,230	6.6(79)	2.8 (9)	12.3(18)	500	16.6(59)	12.2(23)	820	13.0 (82)
Montgomery	679	9.9 (71)	11.2(110)	2,448	5.3(51)	4.2(38)	13.4(30)	1,047	15.9(46)	12.3(26)	1,813	12.0 (73)
Morgan	343	5.2 (3)	10.2(101)	1,661	4.3(21)	4.5(50)	19.6(10)	838	18.1(87)	19.5(10)	1,332	19.4(114)

- (1) Reporting period is January 1 through December 31.  
(2) Analyses based on one record per child.  
(3) Excludes records with unknown data or errors.  
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.  
(5) Infants born during the reporting period included in the analysis.  
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.  
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.  
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.  
\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**2000 - 2002 Pediatric Nutrition Surveillance<sup>(1)</sup>**  
**Kentucky**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years <sup>(2)</sup>**

<b>Comparison of Growth and Anemia Indicators by County</b> <span style="float: right;">(3,4)</span>												
<b>County</b>	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Muhlenberg	854	9.7 (68)	7.8 (49)	3,007	2.9 (4)	7.5(10)	11.2 (8)	1,294	11.6 (9)	10.7 (8)	1,898	11.1 (68)
Nelson	834	11.0 (93)	7.2 (32)	3,102	6.9(93)	4.6(53)	14.2(48)	1,336	16.4(53)	12.1(22)	2,078	9.8 (58)
Nicholas	222	13.5(118)	5.4 (5)	843	4.5(26)	10.0(11)	10.1 (5)	379	10.6 (6)	8.7 (4)	645	13.8 (91)
Ohio	627	10.0 (75)	8.6 (70)	2,448	4.4(25)	3.0(11)	15.2(62)	1,156	16.8(63)	16.1(76)	1,876	10.0 (60)
Oldham	437	10.8 (91)	8.7 (73)	1,597	8.2(10)	3.3(15)	15.2(62)	691	18.5(94)	14.5(50)	1,029	12.6 (76)
Owen	231	7.8 (25)	5.6 (8)	792	5.9(63)	6.4(92)	14.0(45)	339	15.0(32)	13.6(35)	553	6.9 (23)
Owsley	163	7.4 (18)	8.0 (56)	814	5.8(60)	2.7 (8)	25.3(11)	436	21.3(11)	21.1(11)	669	15.8(103)
Pendleton	321	10.6 (86)	5.9 (13)	1,127	6.2(68)	4.4(47)	12.4(20)	479	18.2(89)	13.4(33)	812	8.3 (46)
Perry	1,098	10.3 (78)	7.7 (46)	3,523	7.7(99)	2.2 (3)	30.1(11)	1,596	17.5(78)	22.6(11)	2,614	5.3 (9)
Pike	1,737	9.0 (48)	8.2 (62)	6,464	2.9 (4)	9.1(11)	15.8(70)	2,906	18.8(97)	18.5(97)	4,394	7.3 (31)
Powell	471	10.0 (75)	7.9 (51)	1,899	3.3 (7)	3.8(23)	13.1(26)	939	14.8(30)	13.6(35)	1,477	5.6 (11)

- (1) Reporting period is January 1 through December 31.  
(2) Analyses based on one record per child.  
(3) Excludes records with unknown data or errors.  
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.  
(5) Infants born during the reporting period included in the analysis.  
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.  
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.  
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.  
\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**2000 - 2002 Pediatric Nutrition Surveillance<sup>(1)</sup>**  
**Kentucky**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years <sup>(2)</sup>**

<b>Comparison of Growth and Anemia Indicators by County</b> <span style="float: right;">(3,4)</span>												
<b>County</b>	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Pulaski	1,589	8.4 (31)	7.6 (43)	5,397	5.2(49)	5.8(79)	11.5(10)	2,241	12.9(12)	12.2(23)	3,445	7.3 (31)
Robertson	69	* *	* *	298	8.7(10)	4.0(30)	17.4(92)	137	10.9 (7)	11.7(16)	237	4.2 (6)
Rockcastle	490	6.3 (8)	8.8 (79)	1,787	5.0(38)	6.4(92)	16.6(82)	797	16.3(51)	13.4(33)	1,278	15.3 (99)
Rowan	561	8.9 (43)	14.6(118)	1,987	3.4 (8)	5.0(59)	13.5(34)	836	16.0(48)	14.1(44)	1,269	17.2(109)
Russell	507	8.1 (27)	7.7 (46)	1,732	6.2(68)	4.3(43)	16.2(76)	687	18.9(99)	17.9(93)	1,029	7.4 (35)
Scott	658	9.3 (58)	6.5 (20)	1,826	6.5(75)	6.1(85)	17.1(90)	696	14.7(28)	12.6(27)	1,236	11.8 (71)
Shelby	696	9.1 (51)	8.6 (70)	2,138	14.6(12)	2.6 (6)	32.2(12)	783	22.0(12)	27.8(12)	1,486	11.6 (70)
Simpson	419	9.5 (63)	7.4 (38)	1,261	7.1(94)	4.4(47)	14.4(50)	450	21.8(11)	16.0(75)	758	25.6(119)
Spencer	216	13.4(117)	5.6 (8)	746	10.2(11)	3.8(23)	18.2(99)	296	17.2(72)	16.2(77)	531	9.2 (52)
Taylor	582	7.6 (21)	7.9 (51)	2,173	4.9(35)	5.0(59)	14.1(46)	978	17.1(69)	14.3(47)	1,366	4.1 (5)
Todd	333	10.2 (77)	6.0 (16)	1,323	6.7(85)	4.2(38)	14.1(46)	592	13.9(22)	8.8 (5)	1,005	15.3 (99)

- (1) Reporting period is January 1 through December 31.  
(2) Analyses based on one record per child.  
(3) Excludes records with unknown data or errors.  
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.  
(5) Infants born during the reporting period included in the analysis.  
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.  
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.  
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.  
\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**2000 - 2002 Pediatric Nutrition Surveillance<sup>(1)</sup>**  
**Kentucky**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years<sup>(2)</sup>**

<b>Comparison of Growth and Anemia Indicators by County</b> (3,4)												
<b>County</b>	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Trigg	231	12.6 (114)	6.5 (20)	953	5.1(44)	4.8(57)	12.9(22)	429	13.8(21)	10.7 (8)	717	10.2 (62)
Trimble	202	8.9 (43)	6.9 (27)	667	6.3(71)	7.2(10)	11.1 (6)	286	10.5 (5)	12.9(28)	481	6.9 (23)
Union	398	11.1 (95)	2.3 (1)	1,468	10.4(11)	3.7(20)	14.8(54)	662	18.7(95)	11.8(17)	1,092	7.5 (36)
Warren	2,130	10.5 (83)	7.7 (46)	6,382	5.8(60)	5.1(63)	13.2(28)	2,327	17.0(67)	12.9(28)	3,821	13.6 (89)
Washington	263	6.5 (9)	8.7 (73)	1,080	6.5(75)	3.4(17)	15.2(62)	481	17.9(85)	11.6(13)	813	9.7 (57)
Wayne	626	8.8 (40)	8.6 (70)	2,537	3.5(10)	4.5(50)	16.4(80)	1,201	18.1(87)	20.3(11)	1,690	10.2 (62)
Webster	347	13.3(116)	6.1 (18)	1,170	10.0(11)	2.6 (6)	19.4(10)	470	13.4(19)	19.4(10)	845	12.9 (80)
Whitley	1,665	8.8 (40)	10.0 (97)	6,309	6.3(71)	4.0(30)	16.9(88)	2,882	18.0(86)	18.8(10)	4,663	6.2 (17)
Wolfe	287	9.1 (51)	9.4 (91)	1,308	8.0(10)	1.8 (2)	21.9(11)	670	21.2(11)	20.1(10)	1,051	7.8 (42)
Woodford	417	9.4 (61)	9.1 (86)	1,442	5.1(44)	5.1(63)	15.8(70)	593	14.2(23)	14.2(45)	941	13.6 (89)

- (1) Reporting period is January 1 through December 31.  
(2) Analyses based on one record per child.  
(3) Excludes records with unknown data or errors.  
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.  
(5) Infants born during the reporting period included in the analysis.  
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.  
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.  
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.  
\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**2000 - 2002 Pediatric Nutrition Surveillance<sup>(1)</sup>**  
**Kentucky**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years<sup>(2)</sup>**

<b>Comparison of Growth and Anemia Indicators by County</b> <span style="float: right;">(3,4)</span>												
<b>County</b>	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number
<b>Kentucky</b>	<b>100,497</b>	<b>9.7</b>	<b>8.0</b>	<b>344,976</b>	<b>5.8</b>	<b>5.5</b>	<b>14.8</b>	<b>147,248</b>	<b>16.3</b>	<b>15.0</b>	<b>234,409</b>	<b>11.7</b>

- (1) Reporting period is January 1 through December 31.
- (2) Analyses based on one record per child.
- (3) Excludes records with unknown data or errors.
- (4) Rank compares this county's rate to other counties. Rank 1 = best rate.
- (5) Infants born during the reporting period included in the analysis.
- (6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
- (7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
- (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
- \* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



**Table 7B**  
**2000 - 2002 Pediatric Nutrition Surveillance<sup>(4)</sup>**  
**Kentucky**

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County** (2)  
**Children Aged < 5 Years** (3)

<b>Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County</b> (4,5)						
<b>County</b>	Breastfeeding (6)					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Adair	313	5.8 (72)	162	0.0 (54)	153	0.0 (1)
Allen	322	12.4 (10)	178	0.0 (54)	172	0.0 (1)
Anderson	208	12.5 (9)	121	0.8 (19)	102	0.0 (1)
Ballard	161	6.2 (63)	93	* *	88	* *
Barren	609	7.6 (48)	350	0.9 (15)	298	0.0 (1)
Bath	229	3.9 (99)	127	0.0 (54)	123	0.0 (1)
Bell	870	4.7 (90)	520	0.0 (54)	501	0.0 (1)
Boone	1,233	6.8 (56)	689	1.3 (9)	574	0.0 (1)
Bourbon	310	7.1 (53)	190	0.0 (54)	179	0.0 (1)
Boyd	898	7.0 (55)	460	0.2 (50)	416	0.0 (1)
Boyle	377	5.0 (83)	181	0.0 (54)	175	0.0 (1)
Bracken	142	9.9 (17)	69	* *	64	* *
Breathitt	346	5.2 (81)	180	0.0 (54)	167	0.0 (1)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 7B**  
**2000 - 2002 Pediatric Nutrition Surveillance<sup>(4)</sup>**  
**Kentucky**

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County** (2)  
**Children Aged < 5 Years** (3)

<b>Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County</b> (4,5)						
<b>County</b>	Breastfeeding (6)					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Breckinrid	285	6.7 (58)	164	0.0 (54)	150	0.0 (1)
Bullitt	729	9.7 (21)	338	0.0 (54)	319	0.0 (1)
Butler	270	4.8 (88)	145	0.0 (54)	138	0.0 (1)
Caldwell	259	3.5(107)	150	0.0 (54)	139	0.0 (1)
Calloway	557	13.6 (7)	311	1.0 (12)	285	0.0 (1)
Campbell	923	4.7 (90)	484	0.6 (24)	404	0.0 (1)
Carlisle	99	* *	58	* *	56	* *
Carroll	253	6.3 (62)	145	0.0 (54)	137	0.0 (1)
Carter	699	8.0 (40)	389	0.3 (45)	367	0.0 (1)
Casey	305	7.5 (49)	163	0.6 (24)	150	0.0 (1)
Christian	3,070	10.9 (14)	1,713	1.5 (4)	1,355	0.0 (1)
Clark	533	7.9 (43)	268	0.4 (37)	236	0.0 (1)
Clay	616	3.9 (99)	353	0.0 (54)	345	0.0 (1)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 7B**  
**2000 - 2002 Pediatric Nutrition Surveillance<sup>(4)</sup>**  
**Kentucky**

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County** (2)  
**Children Aged < 5 Years** (3)

<b>Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County</b> (4,5)						
<b>County</b>	Breastfeeding (6)					
	Number	% Ever Breastfed  (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Clinton	278	2.2(112)	165	0.0 (54)	159	0.0 (1)
Crittenden	149	6.0 (67)	84	* *	81	* *
Cumberland	169	5.3 (79)	98	* *	96	* *
Daviess	1,636	6.8 (56)	881	1.0 (12)	747	0.0 (1)
Edmonson	232	9.5 (23)	126	0.0 (54)	94	* *
Elliott	160	3.8(105)	99	* *	91	* *
Estill	344	9.0 (28)	178	0.0 (54)	163	0.0 (1)
Fayette	3,402	21.0 (1)	1,730	1.8 (1)	1,507	0.0 (1)
Fleming	283	4.9 (85)	152	0.0 (54)	150	0.0 (1)
Floyd	1,026	2.3(111)	564	0.0 (54)	537	0.0 (1)
Franklin	671	13.9 (5)	352	1.7 (2)	275	0.0 (1)
Fulton	225	4.9 (85)	123	1.6 (3)	108	0.0 (1)
Gallatin	157	4.5 (93)	78	* *	70	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 7B**  
**2000 - 2002 Pediatric Nutrition Surveillance<sup>(4)</sup>**  
**Kentucky**

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County** (2)  
**Children Aged < 5 Years** (3)

<b>Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County</b> (4,5)						
<b>County</b>	Breastfeeding (6)					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Garrard	311	7.7 (44)	183	0.5 (28)	173	0.0 (1)
Grant	599	5.0 (83)	355	0.3 (45)	339	0.0 (1)
Graves	694	11.4 (13)	370	0.5 (28)	351	0.0 (1)
Grayson	495	14.1 (4)	272	0.0 (54)	265	0.0 (1)
Green	205	11.7 (12)	114	0.0 (54)	107	0.0 (1)
Greenup	580	8.3 (37)	318	0.3 (45)	265	0.0 (1)
Hancock	141	9.2 (24)	81	* *	70	* *
Hardin	2,029	8.6 (31)	1,067	0.3 (45)	883	0.0 (1)
Harlan	897	5.8 (72)	518	0.0 (54)	480	0.0 (1)
Harrison	317	8.2 (39)	183	1.1 (11)	165	0.0 (1)
Hart	379	6.1 (64)	187	0.0 (54)	171	0.0 (1)
Henderson	749	5.6 (76)	368	0.0 (54)	306	0.0 (1)
Henry	238	5.9 (69)	148	1.4 (5)	127	0.0 (1)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 7B**  
**2000 - 2002 Pediatric Nutrition Surveillance<sup>(4)</sup>**  
**Kentucky**

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County** (2)  
**Children Aged < 5 Years** (3)

<b>Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County</b> (4,5)						
<b>County</b>	<b>Breastfeeding</b> (6)					
	<b>Number</b>	<b>% Ever Breastfed  (Rank)</b>	<b>Number</b>	<b>% Breastfed At Least 6 Months (Rank)</b>	<b>Number</b>	<b>% Breastfed At Least 12 Months (Rank)</b>
Hickman	77	* *	42	* *	38	* *
Hopkins	874	5.3 (79)	491	0.4 (37)	424	0.0 (1)
Jackson	361	4.4 (94)	222	0.0 (54)	215	0.0 (1)
Jefferson	9,751	10.4 (15)	4,743	0.5 (28)	4,091	0.0 (1)
Jessamine	626	8.9 (29)	343	0.9 (15)	290	0.0 (1)
Johnson	568	3.9 (99)	326	0.0 (54)	312	0.0 (1)
Kenton	1,369	4.0 (98)	792	0.4 (37)	653	0.0 (1)
Knott	362	1.7(115)	218	0.0 (54)	190	0.0 (1)
Knox	812	3.9 (99)	458	0.2 (50)	426	0.0 (1)
Larue	256	9.8 (19)	144	0.0 (54)	125	0.0 (1)
Laurel	1,039	5.9 (69)	533	0.8 (19)	454	0.0 (1)
Lawrence	350	4.9 (85)	190	0.5 (28)	184	0.0 (1)
Lee	181	3.9 (99)	115	0.9 (15)	104	0.0 (1)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 7B**  
**2000 - 2002 Pediatric Nutrition Surveillance<sup>(4)</sup>**  
**Kentucky**

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County** (2)  
**Children Aged < 5 Years** (3)

<b>Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County</b> (4,5)						
<b>County</b>	Breastfeeding (6)					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Leslie	335	4.8 (88)	201	0.0 (54)	182	0.0 (1)
Letcher	510	6.1 (64)	290	1.0 (12)	240	0.0 (1)
Lewis	291	7.2 (51)	164	0.0 (54)	158	0.0 (1)
Lincoln	517	7.7 (44)	285	0.4 (37)	232	0.0 (1)
Livingston	118	8.5 (33)	72	* *	68	* *
Logan	481	1.7(115)	269	0.0 (54)	246	0.0 (1)
Lyon	103	9.7 (21)	64	* *	56	* *
Madison	1,128	9.2 (24)	580	1.4 (5)	498	0.0 (1)
Magoffin	382	1.6(117)	228	0.0 (54)	221	0.0 (1)
Marion	377	7.7 (44)	204	0.0 (54)	188	0.0 (1)
Marshall	520	9.8 (19)	280	0.4 (37)	262	0.0 (1)
Martin	353	2.0(114)	191	0.0 (54)	185	0.0 (1)
Mason	356	6.5 (59)	188	0.0 (54)	179	0.0 (1)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 7B**  
**2000 - 2002 Pediatric Nutrition Surveillance<sup>(4)</sup>**  
**Kentucky**

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County** (2)  
**Children Aged < 5 Years** (3)

<b>Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County</b> (4,5)						
<b>County</b>	Breastfeeding (6)					
	Number	% Ever Breastfed  (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
McCracken	1,237	6.5 (59)	678	0.0 (54)	644	0.0 (1)
McCreary	486	3.9 (99)	287	0.0 (54)	273	0.0 (1)
McLean	251	9.2 (24)	149	1.3 (9)	144	0.0 (1)
Meade	423	14.2 (2)	235	0.9 (15)	200	0.0 (1)
Menifee	157	8.3 (37)	96	* *	95	* *
Mercer	349	5.7 (75)	185	0.0 (54)	162	0.0 (1)
Metcalfe	208	2.9(110)	117	0.0 (54)	101	0.0 (1)
Monroe	242	8.7 (30)	142	1.4 (5)	112	0.0 (1)
Montgomery	517	6.0 (67)	285	0.4 (37)	242	0.0 (1)
Morgan	299	4.3 (95)	183	0.5 (28)	173	0.0 (1)
Muhlenberg	606	4.3 (95)	344	0.0 (54)	329	0.0 (1)
Nelson	609	9.2 (24)	328	0.0 (54)	285	0.0 (1)
Nicholas	171	5.8 (72)	88	* *	75	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 7B**  
**2000 - 2002 Pediatric Nutrition Surveillance<sup>(4)</sup>**  
**Kentucky**

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County** (2)  
**Children Aged < 5 Years** (3)

<b>Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County</b> (4,5)						
<b>County</b>	Breastfeeding (6)					
	Number	% Ever Breastfed  (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Ohio	464	8.4 (36)	249	0.0 (54)	222	0.0 (1)
Oldham	356	11.8 (11)	197	0.5 (28)	185	0.0 (1)
Owen	162	13.6 (7)	79	* *	72	* *
Owsley	135	2.2(112)	83	* *	77	* *
Pendleton	252	9.9 (17)	148	0.7 (22)	141	0.0 (1)
Perry	725	3.4(108)	362	0.3 (45)	346	0.0 (1)
Pike	1,228	6.4 (61)	653	0.2 (50)	560	0.0 (1)
Powell	368	4.1 (97)	194	0.5 (28)	185	0.0 (1)
Pulaski	1,204	10.2 (16)	641	0.6 (24)	548	0.0 (1)
Robertson	60	* *	31	* *	30	* *
Rockcastle	382	7.3 (50)	187	0.5 (28)	167	0.0 (1)
Rowan	419	3.6(106)	235	0.4 (37)	203	0.0 (1)
Russell	387	8.0 (40)	221	0.0 (54)	210	0.0 (1)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



**Table 7B**  
**2000 - 2002 Pediatric Nutrition Surveillance<sup>(4)</sup>**  
**Kentucky**

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County** (2)  
**Children Aged < 5 Years** (3)

<b>Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County</b> (4,5)						
<b>County</b>	Breastfeeding (6)					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Scott	472	8.5 (33)	240	0.4 (37)	223	0.0 (1)
Shelby	536	5.4 (78)	281	0.0 (54)	255	0.0 (1)
Simpson	293	8.5 (33)	152	0.0 (54)	137	0.0 (1)
Spencer	163	6.1 (64)	88	* *	83	* *
Taylor	454	7.7 (44)	255	0.0 (54)	242	0.0 (1)
Todd	263	7.2 (51)	160	0.6 (24)	152	0.0 (1)
Trigg	181	13.8 (6)	102	0.0 (54)	94	* *
Trimble	150	8.0 (40)	86	* *	84	* *
Union	294	7.1 (53)	170	0.0 (54)	147	0.0 (1)
Warren	1,504	8.6 (31)	727	0.7 (22)	625	0.0 (1)
Washington	219	4.6 (92)	130	0.0 (54)	124	0.0 (1)
Wayne	498	5.2 (81)	264	0.8 (19)	238	0.0 (1)
Webster	256	5.9 (69)	141	1.4 (5)	105	0.0 (1)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 7B**  
**2000 - 2002 Pediatric Nutrition Surveillance<sup>(4)</sup>**  
**Kentucky**

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County** (2)  
**Children Aged < 5 Years** (3)

<b>Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County</b> (4,5)						
<b>County</b>	Breastfeeding (6)					
	Number	% Ever Breastfed  (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Whitley	1,238	5.5 (77)	715	0.1 (53)	612	0.0 (1)
Wolfe	239	3.3(109)	140	0.0 (54)	132	0.0 (1)
Woodford	344	14.2 (2)	207	0.5 (28)	189	0.0 (1)
Unknown Counties	1,242	1.6(117)	62	* *	57	* *
<b>Kentucky</b>	<b>73,122</b>	<b>8.1</b>	<b>38,745</b>	<b>0.5</b>	<b>34,374</b>	<b>0.0</b>

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.