

Table 4B
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky
Comparison of Racial and Ethnic Distribution by County
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution by County							
County	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other / Unknown %
Adair	514	93.8	4.3	1.9	0.0	0.0	0.0
Allen	707	98.2	0.6	0.7	0.0	0.6	0.0
Anderson	362	90.1	6.6	3.3	0.0	0.0	0.0
Ballard	304	91.1	8.2	0.0	0.0	0.7	0.0
Barren	1,299	90.2	6.6	2.2	0.0	0.9	0.0
Bath	874	98.9	0.2	0.5	0.5	0.0	0.0
Bell	1,877	96.2	2.1	1.5	0.1	0.1	0.0
Boone	2,158	89.1	2.6	8.0	0.0	0.4	0.0
Bourbon	534	77.3	9.0	12.9	0.4	0.4	0.0
Boyd	1,373	96.8	1.9	0.9	0.1	0.2	0.0
Boyle	634	93.5	6.5	0.0	0.0	0.0	0.0
Bracken	269	99.6	0.0	0.4	0.0	0.0	0.0
Breathitt	708	99.9	0.1	0.0	0.0	0.0	0.0
Breckinrid	639	95.0	3.9	1.1	0.0	0.0	0.0
Bullitt	1,183	98.0	1.4	0.4	0.0	0.2	0.0
Butler	554	95.7	1.1	0.0	0.0	3.2	0.0
Caldwell	432	92.4	7.6	0.0	0.0	0.0	0.0
Calloway	993	87.5	7.4	3.4	0.1	1.6	0.0
Campbell	1,475	90.6	7.2	1.8	0.1	0.3	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4B
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky
Comparison of Racial and Ethnic Distribution by County
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution by County							
County	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other / Unknown %
Carlisle	234	94.9	5.1	0.0	0.0	0.0	0.0
Carroll	423	80.6	2.1	17.0	0.0	0.2	0.0
Carter	1,325	99.2	0.5	0.2	0.1	0.0	0.0
Casey	550	94.5	0.0	5.5	0.0	0.0	0.0
Christian	4,806	66.0	28.3	4.6	0.3	0.9	0.0
Clark	817	91.6	5.8	1.6	0.1	1.0	0.0
Clay	1,280	98.9	0.8	0.2	0.1	0.0	0.0
Clinton	546	93.0	0.0	6.6	0.0	0.4	0.0
Crittenden	305	99.7	0.3	0.0	0.0	0.0	0.0
Cumberland	336	94.6	4.5	0.6	0.0	0.3	0.0
Daviess	2,756	84.1	11.7	3.7	0.0	0.4	0.0
Edmonson	452	100.0	0.0	0.0	0.0	0.0	0.0
Elliott	369	99.2	0.0	0.3	0.5	0.0	0.0
Estill	645	98.6	0.8	0.0	0.0	0.6	0.0
Fayette	5,371	46.1	29.4	17.9	0.6	6.0	0.0
Fleming	633	96.5	1.4	1.6	0.2	0.3	0.0
Floyd	2,051	99.6	0.2	0.1	0.0	0.0	0.0
Franklin	971	75.8	14.8	7.7	0.3	1.3	0.0
Fulton	419	58.9	41.1	0.0	0.0	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4B
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky
Comparison of Racial and Ethnic Distribution by County
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution by County							
County	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other / Unknown %
Gallatin	284	82.0	2.5	15.5	0.0	0.0	0.0
Garrard	602	94.5	1.8	3.7	0.0	0.0	0.0
Grant	1,069	97.9	1.4	0.1	0.2	0.4	0.0
Graves	1,276	72.3	10.5	15.4	0.2	1.5	0.0
Grayson	975	98.2	0.6	0.7	0.2	0.3	0.0
Green	395	93.9	4.6	1.3	0.0	0.3	0.0
Greenup	1,048	99.5	0.5	0.0	0.0	0.0	0.0
Hancock	284	96.1	2.1	1.8	0.0	0.0	0.0
Hardin	3,428	78.7	18.5	2.0	0.2	0.6	0.0
Harlan	1,794	97.1	2.6	0.2	0.0	0.1	0.0
Harrison	597	92.6	2.8	4.5	0.0	0.0	0.0
Hart	621	96.5	3.4	0.2	0.0	0.0	0.0
Henderson	1,221	82.5	12.3	1.8	0.5	2.9	0.0
Henry	444	85.4	3.4	11.3	0.0	0.0	0.0
Hickman	142	82.4	16.9	0.0	0.0	0.7	0.0
Hopkins	1,480	84.5	12.3	2.7	0.2	0.3	0.0
Jackson	726	99.9	0.0	0.1	0.0	0.0	0.0
Jefferson	15,135	47.5	44.0	5.4	0.3	2.9	0.0
Jessamine	1,106	88.2	6.3	3.1	0.1	2.3	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4B
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky
Comparison of Racial and Ethnic Distribution by County
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution by County							
County	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other / Unknown %
Johnson	1,186	99.9	0.1	0.0	0.0	0.0	0.0
Kenton	2,175	85.4	13.6	0.4	0.3	0.3	0.0
Knott	794	99.5	0.4	0.1	0.0	0.0	0.0
Knox	1,606	98.9	1.1	0.0	0.0	0.0	0.0
Larue	461	88.1	5.0	6.1	0.4	0.4	0.0
Laurel	1,631	98.7	0.8	0.1	0.1	0.3	0.0
Lawrence	649	99.8	0.2	0.0	0.0	0.0	0.0
Lee	399	99.0	1.0	0.0	0.0	0.0	0.0
Leslie	680	99.7	0.3	0.0	0.0	0.0	0.0
Letcher	920	99.3	0.5	0.0	0.0	0.1	0.0
Lewis	598	99.7	0.0	0.0	0.0	0.3	0.0
Lincoln	906	93.3	2.9	3.8	0.0	0.1	0.0
Livingston	242	98.3	0.0	0.4	0.0	1.2	0.0
Logan	775	82.5	13.9	3.0	0.6	0.0	0.0
Lyon	177	98.3	1.7	0.0	0.0	0.0	0.0
Madison	1,959	88.9	7.9	1.7	0.2	1.3	0.0
Magoffin	872	99.1	0.0	0.9	0.0	0.0	0.0
Marion	703	81.9	11.0	5.8	0.0	1.3	0.0
Marshall	878	99.1	0.8	0.1	0.0	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4B
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky
Comparison of Racial and Ethnic Distribution by County
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution by County							
County	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other / Unknown %
Martin	746	99.2	0.7	0.1	0.0	0.0	0.0
Mason	547	91.0	7.3	0.9	0.4	0.4	0.0
McCracken	2,031	65.5	30.8	1.9	0.5	1.3	0.0
McCreary	1,065	99.7	0.3	0.0	0.0	0.0	0.0
McLean	441	97.5	0.7	1.4	0.2	0.2	0.0
Meade	743	97.6	2.4	0.0	0.0	0.0	0.0
Menifee	371	99.2	0.0	0.8	0.0	0.0	0.0
Mercer	656	87.2	4.7	8.1	0.0	0.0	0.0
Metcalfe	394	97.5	1.5	0.5	0.5	0.0	0.0
Monroe	411	91.7	4.6	3.6	0.0	0.0	0.0
Montgomery	990	96.1	0.7	2.8	0.0	0.4	0.0
Morgan	660	99.8	0.2	0.0	0.0	0.0	0.0
Muhlenberg	978	94.9	4.5	0.5	0.0	0.1	0.0
Nelson	1,178	90.2	8.7	0.8	0.0	0.3	0.0
Nicholas	307	99.3	0.0	0.7	0.0	0.0	0.0
Ohio	896	92.3	0.9	6.8	0.0	0.0	0.0
Oldham	747	79.8	4.6	15.3	0.0	0.4	0.0
Owen	316	90.2	0.0	8.9	0.9	0.0	0.0
Owsley	299	99.0	1.0	0.0	0.0	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4B
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky
Comparison of Racial and Ethnic Distribution by County
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution by County							
County	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other / Unknown %
Pendleton	381	96.6	2.6	0.8	0.0	0.0	0.0
Perry	1,356	98.0	1.8	0.1	0.0	0.1	0.0
Pike	2,133	98.9	0.5	0.1	0.2	0.2	0.0
Powell	745	97.0	0.9	1.3	0.0	0.7	0.0
Pulaski	2,092	96.8	0.8	2.3	0.0	0.1	0.0
Robertson	118	93.2	0.0	6.8	0.0	0.0	0.0
Rockcastle	663	99.5	0.2	0.2	0.0	0.2	0.0
Rowan	695	97.7	1.3	0.3	0.1	0.6	0.0
Russell	627	91.9	1.1	6.2	0.0	0.8	0.0
Scott	724	85.9	6.1	6.4	1.1	0.6	0.0
Shelby	841	49.6	11.4	39.0	0.0	0.0	0.0
Simpson	474	79.3	18.4	1.5	0.0	0.8	0.0
Spencer	292	91.8	0.3	6.5	0.7	0.7	0.0
Taylor	842	91.9	7.6	0.2	0.2	0.0	0.0
Todd	489	72.2	16.0	4.9	0.2	6.7	0.0
Trigg	333	82.6	15.6	0.6	0.9	0.3	0.0
Trimble	252	95.6	0.0	4.4	0.0	0.0	0.0
Union	483	84.9	14.1	0.2	0.0	0.8	0.0
Warren	2,186	66.4	15.2	13.4	0.5	4.5	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4B
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky
Comparison of Racial and Ethnic Distribution by County
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution by County							
County	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other / Unknown %
Washington	432	76.9	8.8	13.2	0.0	1.2	0.0
Wayne	959	91.3	1.6	7.1	0.0	0.0	0.0
Webster	475	79.6	3.4	14.1	0.0	2.9	0.0
Whitley	2,474	99.5	0.2	0.0	0.1	0.2	0.0
Wolfe	457	99.3	0.7	0.0	0.0	0.0	0.0
Unknown Counties	1,168	93.1	5.8	0.8	0.1	0.3	0.0
Kentucky	126,883	83.0	11.9	3.9	0.2	1.0	0.0
Nation (Prior Year)	5,333,226	39.1	20.8	32.2	1.2	3.3	3.5

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5B
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky
Comparison of Age Distribution by County
Children Aged < 5 Years ⁽²⁾

Comparison of Age Distribution by County				
County	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Adair	514	22.6	28.2	49.2
Allen	707	18.1	24.0	57.9
Anderson	362	26.0	22.1	51.9
Ballard	304	20.1	25.3	54.6
Barren	1,299	26.5	24.6	49.0
Bath	874	20.4	25.4	54.2
Bell	1,877	17.3	24.6	58.1
Boone	2,158	29.3	25.9	44.9
Bourbon	534	22.1	27.3	50.6
Boyd	1,373	28.1	25.1	46.8
Boyle	634	26.0	29.8	44.2
Bracken	269	23.8	21.9	54.3
Breathitt	708	19.6	26.4	54.0
Breckinrid	639	24.7	22.1	53.2
Bullitt	1,183	28.7	27.9	43.4
Butler	554	22.7	21.5	55.8
Caldwell	432	19.7	29.2	51.2
Calloway	993	22.6	28.4	49.0

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5B
2003 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Age Distribution by County
Children Aged < 5 Years (2)

Comparison of Age Distribution by County				
County	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Campbell	1,475	31.5	23.9	44.6
Carlisle	234	16.7	24.4	59.0
Carroll	423	25.8	28.6	45.6
Carter	1,325	16.9	26.6	56.5
Casey	550	21.1	27.6	51.3
Christian	4,806	28.7	23.1	48.1
Clark	817	28.0	27.8	44.2
Clay	1,280	18.4	25.4	56.3
Clinton	546	16.8	26.7	56.4
Crittenden	305	21.6	30.8	47.5
Cumberland	336	17.0	23.5	59.5
Daviess	2,756	28.4	24.1	47.5
Edmonson	452	29.9	25.0	45.1
Elliott	369	20.9	21.7	57.5
Estill	645	20.3	24.7	55.0
Fayette	5,371	30.6	25.4	44.0
Fleming	633	16.1	25.6	58.3
Floyd	2,051	18.6	25.2	56.2

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5B
2003 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Age Distribution by County
Children Aged < 5 Years (2)

Comparison of Age Distribution by County				
County	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Franklin	971	34.0	26.0	40.1
Fulton	419	17.2	27.4	55.4
Gallatin	284	26.8	27.8	45.4
Garrard	602	19.4	28.4	52.2
Grant	1,069	21.0	24.3	54.7
Graves	1,276	21.5	25.9	52.7
Grayson	975	19.4	26.4	54.3
Green	395	22.5	24.8	52.7
Greenup	1,048	26.2	22.8	51.0
Hancock	284	20.8	23.2	56.0
Hardin	3,428	26.6	24.1	49.3
Harlan	1,794	18.3	24.6	57.1
Harrison	597	23.8	25.5	50.8
Hart	621	21.4	28.8	49.8
Henderson	1,221	28.3	23.3	48.5
Henry	444	31.8	22.1	46.2
Hickman	142	28.2	25.4	46.5
Hopkins	1,480	26.6	23.8	49.7

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5B
2003 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Age Distribution by County
Children Aged < 5 Years (2)

Comparison of Age Distribution by County				
County	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Jackson	726	18.0	22.5	59.5
Jefferson	15,135	31.0	24.4	44.6
Jessamine	1,106	26.3	26.8	46.9
Johnson	1,186	18.0	24.4	57.7
Kenton	2,175	31.3	25.4	43.3
Knott	794	20.0	22.5	57.4
Knox	1,606	20.4	22.5	57.0
Larue	461	29.1	23.4	47.5
Laurel	1,631	28.9	24.6	46.5
Lawrence	649	18.5	25.3	56.2
Lee	399	18.0	24.6	57.4
Leslie	680	16.5	23.8	59.7
Letcher	920	23.6	25.1	51.3
Lewis	598	15.6	25.6	58.9
Lincoln	906	24.9	23.7	51.3
Livingston	242	21.1	26.0	52.9
Logan	775	24.3	25.5	50.2
Lyon	177	19.8	26.0	54.2

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5B
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky
Comparison of Age Distribution by County
Children Aged < 5 Years ⁽²⁾

Comparison of Age Distribution by County				
County	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Madison	1,959	28.4	26.2	45.4
Magoffin	872	16.2	23.1	60.8
Marion	703	22.2	24.8	53.1
Marshall	878	20.4	29.8	49.8
Martin	746	16.9	27.2	55.9
Mason	547	18.5	28.5	53.0
McCracken	2,031	20.8	31.1	48.1
McCreary	1,065	21.2	24.3	54.5
McLean	441	16.3	24.9	58.7
Meade	743	25.4	23.0	51.5
Menifee	371	21.6	21.8	56.6
Mercer	656	25.6	23.9	50.5
Metcalfe	394	19.5	29.7	50.8
Monroe	411	21.7	25.3	53.0
Montgomery	990	23.5	25.1	51.4
Morgan	660	15.8	25.9	58.3
Muhlenberg	978	21.8	22.4	55.8
Nelson	1,178	26.5	24.3	49.2

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5B
2003 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Age Distribution by County
Children Aged < 5 Years (2)

Comparison of Age Distribution by County				
County	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Nicholas	307	16.9	28.3	54.7
Ohio	896	23.5	24.3	52.1
Oldham	747	23.2	26.1	50.7
Owen	316	19.3	27.5	53.2
Owsley	299	14.7	28.4	56.9
Pendleton	381	17.6	31.0	51.4
Perry	1,356	25.2	25.9	48.9
Pike	2,133	25.8	25.3	48.9
Powell	745	16.5	29.0	54.5
Pulaski	2,092	26.3	26.8	46.8
Robertson	118	11.9	28.0	60.2
Rockcastle	663	23.5	25.0	51.4
Rowan	695	24.9	25.2	49.9
Russell	627	18.8	30.1	51.0
Scott	724	28.2	29.8	42.0
Shelby	841	29.5	27.3	43.2
Simpson	474	25.7	24.7	49.6
Spencer	292	24.7	26.4	49.0

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5B
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky
Comparison of Age Distribution by County
Children Aged < 5 Years ⁽²⁾

Comparison of Age Distribution by County				
County	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Taylor	842	22.1	24.8	53.1
Todd	489	23.1	23.5	53.4
Trigg	333	24.3	31.2	44.4
Trimble	252	23.4	28.6	48.0
Union	483	24.2	24.4	51.3
Warren	2,186	33.9	24.0	42.1
Washington	432	20.8	22.2	56.9
Wayne	959	20.1	24.8	55.1
Webster	475	27.8	26.9	45.3
Whitley	2,474	23.7	22.9	53.4
Wolfe	457	19.0	25.2	55.8
Unknown Counties	1,168	100.0	0.0	0.0
Kentucky	126,883	25.7	24.9	49.4
Nation (Prior Year)	5,333,226	36.6	21.7	41.7

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2003 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Adair	101	7.9 (18)	8.9 (53)	491	4.7(18)	4.5(82)	14.7(22)	244	16.0(37)	13.5(18)	236	7.6 (32)
Allen	122	8.2 (20)	11.5 (73)	688	6.3(44)	2.2(17)	17.2(46)	398	20.4(93)	17.3(61)	372	10.5 (62)
Anderson	74	* *	* *	347	7.5(65)	2.9(38)	22.2(99)	182	20.3(91)	20.3(83)	277	10.1 (57)
Ballard	57	* *	* *	289	6.6(49)	9.3(11)	13.1(11)	163	11.0 (5)	11.7 (8)	234	6.4 (17)
Barren	287	10.5 (51)	7.7 (37)	1,249	7.8(68)	4.7(90)	18.2(65)	611	15.9(34)	16.9(54)	726	12.4 (81)
Bath	148	12.2 (69)	12.2 (75)	842	9.3(92)	2.6(29)	21.9(98)	466	18.5(66)	20.6(88)	386	8.8 (43)
Bell	273	6.6 (10)	8.4 (44)	1,807	7.1(57)	4.3(74)	18.1(62)	1,057	17.3(53)	16.0(41)	1,065	7.2 (26)
Boone	511	6.3 (7)	10.4 (65)	2,085	8.1(73)	3.8(60)	19.5(77)	945	20.4(93)	15.6(37)	1,374	14.3 (90)
Bourbon	108	11.1 (59)	7.4 (32)	516	9.5(94)	2.3(20)	20.5(87)	263	16.3(44)	22.8(10)	394	26.1 (116)
Boyd	343	13.7 (73)	7.9 (39)	1,318	6.7(51)	6.2(11)	14.6(21)	626	19.3(77)	15.5(35)	757	10.0 (54)
Boyle	147	8.8 (28)	8.8 (49)	595	4.9(23)	4.0(65)	18.5(68)	265	15.1(23)	18.5(76)	361	8.0 (36)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2003 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Bracken	52	* *	* *	256	12.9(11)	5.9(11)	27.0(11)	139	15.1(23)	34.5(11)	181	7.2 (26)
Breathitt	126	11.1 (59)	8.7 (48)	668	6.0(39)	4.6(84)	20.8(90)	374	19.5(81)	21.4(95)	426	10.3 (58)
Breckinrid	113	8.8 (28)	4.4 (7)	625	6.6(49)	2.6(29)	16.3(34)	334	18.9(71)	14.4(23)	492	2.8 (2)
Bullitt	307	6.2 (5)	11.1 (70)	1,141	7.4(62)	4.2(71)	17.7(55)	503	15.7(30)	15.5(35)	572	20.8(109)
Butler	118	6.8 (14)	8.5 (47)	539	4.5(13)	2.6(29)	21.0(93)	304	20.7(99)	24.7(10)	280	8.9 (45)
Caldwell	65	* *	* *	421	5.9(36)	4.3(74)	20.9(91)	219	14.6(20)	17.4(63)	349	10.0 (54)
Calloway	183	6.6 (10)	11.5 (73)	962	7.3(60)	1.8(14)	16.3(34)	480	20.6(98)	17.1(56)	603	25.5(114)
Campbell	366	8.7 (27)	9.6 (60)	1,392	8.6(81)	3.5(53)	17.0(44)	630	19.5(81)	14.6(26)	893	15.5 (94)
Carlisle	33	* *	* *	231	6.5(47)	2.2(17)	28.6(11)	138	20.3(91)	28.3(11)	123	20.3(107)
Carroll	94	* *	* *	411	9.7(97)	3.9(62)	20.9(91)	186	26.3(11)	21.0(90)	308	11.7 (74)
Carter	200	10.0 (43)	9.0 (54)	1,280	4.5(13)	6.1(11)	12.9(10)	729	16.6(48)	13.0(14)	651	10.9 (66)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2003 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Casey	90	* *	* *	535	10.3(10)	2.2(17)	16.4(37)	279	16.1(39)	19.4(79)	259	16.2 (96)
Christian	916	8.3 (21)	7.9 (39)	4,616	5.6(35)	5.7(10)	11.3 (2)	2,253	13.2(13)	9.9 (3)	2,617	27.0(118)
Clark	192	11.5 (65)	3.1 (3)	784	4.8(21)	4.2(71)	14.0(16)	357	15.1(23)	18.5(76)	460	12.8 (86)
Clay	219	11.0 (57)	7.3 (30)	1,231	6.1(41)	5.0(99)	19.7(80)	700	18.3(63)	20.7(89)	955	8.5 (41)
Clinton	84	* *	* *	534	6.0(39)	4.5(82)	20.0(82)	299	20.1(90)	16.7(49)	270	11.5 (71)
Crittenden	53	* *	* *	297	2.4 (1)	4.0(65)	13.8(14)	143	14.0(18)	14.7(28)	196	13.8 (89)
Cumberland	54	* *	* *	329	4.6(17)	4.3(74)	14.0(16)	196	13.3(14)	16.3(45)	140	19.3(103)
Daviess	615	8.5 (25)	6.2 (21)	2,656	10.8(10)	2.0(16)	20.3(86)	1,280	17.7(58)	17.1(56)	1,807	9.2 (50)
Edmonson	94	* *	* *	443	4.5(13)	6.1(11)	11.3 (2)	201	13.9(17)	11.4 (7)	249	6.4 (17)
Elliott	66	* *	* *	360	4.4(12)	8.6(11)	15.3(26)	209	12.9(12)	15.3(33)	187	25.7(115)
Estill	114	5.3 (2)	8.8 (49)	626	6.1(41)	1.6(10)	18.2(65)	345	19.1(75)	21.2(92)	520	11.5 (71)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2003 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Fayette	1,319	10.2 (46)	6.5 (25)	5,149	7.3(60)	3.5(53)	16.8(41)	2,307	19.4(80)	18.2(74)	2,812	13.6 (88)
Fleming	89	* *	* *	611	5.9(36)	3.4(48)	17.8(57)	355	12.1(10)	16.3(45)	518	16.2 (96)
Floyd	352	11.4 (63)	7.4 (32)	1,938	5.0(25)	7.3(11)	15.9(32)	1,084	11.8 (9)	14.6(26)	1,319	10.4 (60)
Franklin	257	9.7 (37)	7.8 (38)	938	8.0(71)	6.0(11)	17.9(58)	379	19.8(85)	20.3(83)	694	19.9(106)
Fulton	50	* *	* *	399	7.8(68)	3.0(41)	18.0(61)	224	21.9(10)	14.7(28)	258	15.1 (91)
Gallatin	61	* *	* *	278	11.2(10)	4.0(65)	14.0(16)	127	12.6(11)	16.5(47)	184	12.0 (78)
Garrard	105	6.7 (12)	5.7 (15)	574	8.7(83)	2.3(20)	12.5 (8)	298	23.8(11)	9.7 (2)	282	18.1(101)
Grant	203	10.3 (47)	10.8 (68)	1,024	5.0(25)	4.4(79)	11.4 (4)	571	11.7 (8)	9.1 (1)	827	5.2 (10)
Graves	238	11.3 (61)	8.8 (49)	1,222	8.7(83)	5.3(10)	17.9(58)	650	19.1(75)	18.8(78)	726	19.4(104)
Grayson	173	4.0 (1)	9.8 (63)	949	12.2(11)	3.3(44)	22.6(10)	513	17.7(58)	17.7(69)	734	3.1 (5)
Green	77	* *	* *	383	12.0(11)	1.6(10)	26.1(11)	198	17.7(58)	25.3(11)	205	8.8 (43)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2003 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Greenup	219	10.5 (51)	6.8 (27)	1,019	3.4 (3)	7.7(11)	14.9(24)	526	18.3(63)	15.2(31)	584	7.9 (35)
Hancock	45	* *	* *	275	5.5(33)	1.1 (2)	17.5(51)	157	17.2(51)	15.3(33)	230	8.3 (38)
Hardin	662	9.5 (36)	6.8 (27)	3,324	6.7(51)	2.5(27)	13.7(13)	1,651	17.9(62)	10.5 (4)	2,643	7.2 (26)
Harlan	276	9.8 (38)	9.4 (58)	1,715	5.0(25)	4.7(90)	21.4(95)	985	17.8(61)	23.7(10)	1,102	6.4 (17)
Harrison	122	10.7 (54)	0.8 (1)	576	6.3(44)	5.0(99)	12.0 (6)	293	9.9 (3)	10.6 (5)	354	15.8 (95)
Hart	111	6.3 (7)	9.0 (54)	611	5.2(31)	3.4(48)	16.9(43)	303	15.5(28)	22.4(10)	391	6.6 (21)
Henderson	249	13.3 (71)	5.2 (10)	1,187	9.9(99)	3.5(53)	17.9(58)	579	25.6(11)	23.3(10)	303	5.6 (12)
Henry	111	10.8 (56)	6.3 (23)	421	10.9(10)	3.3(44)	22.8(10)	200	16.5(45)	17.5(66)	219	11.4 (68)
Hickman	30	* *	* *	141	2.8 (2)	5.0(99)	10.6 (1)	66	* *	* *	89	* *
Hopkins	289	10.0 (43)	7.3 (30)	1,424	13.3(11)	1.3 (3)	19.1(75)	713	21.3(10)	18.0(71)	1,097	10.5 (62)
Jackson	114	8.8 (28)	8.8 (49)	708	3.8 (6)	4.0(65)	18.2(65)	421	18.3(63)	15.9(40)	594	3.5 (7)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2003 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Jefferson	3,874	10.5 (51)	6.4 (24)	14,441	8.4(79)	4.6(84)	16.4(37)	6,584	17.5(55)	15.7(38)	9,392	9.9 (53)
Jessamine	224	10.7 (54)	11.2 (71)	1,071	10.4(10)	3.7(57)	17.6(52)	513	17.5(55)	14.0(21)	695	9.1 (48)
Johnson	183	9.3 (34)	5.5 (13)	1,143	3.9 (8)	5.5(10)	16.8(41)	663	16.6(48)	16.1(42)	805	24.2(113)
Kenton	535	9.9 (40)	5.6 (14)	2,094	9.5(94)	3.4(48)	22.2(99)	925	25.6(11)	26.4(11)	1,549	12.1 (79)
Knott	108	12.0 (68)	3.7 (5)	761	12.2(11)	3.7(57)	22.5(10)	436	16.5(45)	18.3(75)	649	7.2 (26)
Knox	270	11.9 (67)	10.4 (65)	1,544	4.2 (9)	3.2(42)	18.8(73)	891	21.4(10)	21.3(93)	1,034	12.4 (81)
Larue	104	5.8 (4)	6.7 (26)	428	7.9(70)	2.6(29)	22.7(10)	210	16.2(41)	16.7(49)	340	10.0 (54)
Laurel	379	11.3 (61)	6.1 (19)	1,561	7.4(62)	5.0(99)	18.1(62)	730	21.0(10)	19.6(80)	752	15.4 (93)
Lawrence	107	15.9 (76)	7.5 (35)	628	6.5(47)	4.6(84)	18.5(68)	358	23.7(11)	17.3(61)	393	5.6 (12)
Lee	52	* *	* *	386	8.5(80)	2.6(29)	24.4(11)	220	20.9(10)	23.6(10)	321	15.3 (92)
Leslie	92	* *	* *	657	3.8 (6)	2.3(20)	24.7(11)	401	19.7(84)	24.9(10)	581	9.1 (48)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2003 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Letcher	169	8.9 (31)	5.9 (18)	887	8.3(76)	2.9(38)	20.6(88)	460	15.7(30)	16.7(49)	718	2.4 (1)
Lewis	85	* *	* *	581	4.5(13)	4.0(65)	14.5(19)	344	21.2(10)	13.4(16)	284	4.9 (9)
Lincoln	155	11.0 (57)	9.0 (54)	873	10.0(10)	3.4(48)	22.9(10)	453	19.9(87)	20.5(87)	684	5.3 (11)
Livingston	42	* *	* *	233	4.7(18)	3.9(62)	17.6(52)	127	11.0 (5)	13.4(16)	197	3.0 (3)
Logan	149	7.4 (16)	8.1 (43)	744	7.5(65)	2.3(20)	17.6(52)	379	22.4(10)	16.6(48)	405	21.7(112)
Lyon	22	* *	* *	172	6.4(46)	5.2(10)	22.7(10)	94	* *	* *	102	7.8 (34)
Madison	452	10.4 (50)	10.2 (64)	1,904	8.9(87)	5.6(10)	15.4(28)	874	19.9(87)	20.3(83)	1,397	8.4 (39)
Magoffin	129	9.3 (34)	10.9 (69)	837	6.8(53)	2.6(29)	25.2(11)	511	20.5(97)	21.3(93)	445	26.3(117)
Marion	119	6.7 (12)	8.4 (44)	687	12.8(11)	1.6(10)	23.3(10)	366	18.6(68)	19.7(81)	496	11.7 (74)
Marshall	153	6.5 (9)	10.5 (67)	842	7.7(67)	5.7(10)	16.6(40)	426	14.6(20)	10.8 (6)	628	7.3 (31)
Martin	112	9.8 (38)	8.0 (41)	724	8.0(71)	3.3(44)	21.5(96)	407	23.6(11)	17.2(58)	618	6.8 (22)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2003 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low		High	Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Mason	90	* *	* *	526	10.1(10)	4.9(95)	20.2(85)	285	19.3(77)	15.8(39)	349	12.3 (80)
McCracken	367	8.4 (23)	5.4 (11)	1,927	8.1(73)	4.9(95)	20.0(82)	947	17.6(57)	17.2(58)	1,412	21.1 (110)
McCreary	195	10.3 (47)	4.6 (8)	1,034	8.3(76)	3.3(44)	15.2(25)	570	19.8(85)	18.1(73)	791	7.2 (26)
McLean	60	* *	* *	420	16.7(11)	2.4(26)	22.6(10)	249	13.7(16)	26.1(11)	302	3.3 (6)
Meade	154	8.4 (23)	8.4 (44)	717	9.5(94)	4.6(84)	17.4(49)	378	15.9(34)	14.8(30)	551	11.4 (68)
Menifee	72	* *	* *	362	5.0(25)	4.1(70)	12.4 (7)	205	10.2 (4)	16.1(42)	188	9.0 (47)
Mercer	133	15.0 (75)	7.5 (35)	633	9.0(90)	2.5(27)	16.3(34)	322	16.8(50)	17.4(63)	478	9.6 (51)
Metcalfe	67	* *	* *	379	5.3(32)	4.7(90)	15.6(29)	197	14.7(22)	16.2(44)	228	11.0 (67)
Monroe	65	* *	* *	399	5.0(25)	4.3(74)	15.8(30)	211	16.1(39)	18.0(71)	244	20.5(108)
Montgomery	176	11.4 (63)	5.7 (15)	943	6.9(54)	2.3(20)	13.6(12)	493	15.6(29)	12.6(13)	767	7.0 (25)
Morgan	93	* *	* *	643	4.8(21)	3.7(57)	21.2(94)	378	20.4(93)	20.4(86)	555	11.7 (74)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2003 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low		High	Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Muhlenberg	193	5.7 (3)	6.2 (21)	953	4.7(18)	5.7(10)	15.3(26)	538	15.2(26)	17.5(66)	481	12.7 (84)
Nelson	233	9.9 (40)	2.6 (2)	1,146	9.2(91)	4.4(79)	18.1(62)	565	15.9(34)	11.7 (8)	729	10.4 (60)
Nicholas	46	* *	* *	295	7.1(57)	3.4(48)	18.6(70)	161	8.7 (1)	13.7(19)	246	19.1 (102)
Ohio	175	9.1 (32)	5.7 (15)	871	7.2(59)	4.2(71)	14.7(22)	458	16.2(41)	16.8(52)	654	9.8 (52)
Oldham	143	14.7 (74)	3.5 (4)	718	11.7(11)	1.3 (3)	20.1(84)	368	16.0(37)	17.9(70)	412	11.4 (68)
Owen	55	* *	* *	304	4.9(23)	3.9(62)	13.8(14)	164	19.5(81)	15.2(31)	248	6.9 (23)
Owsley	35	* *	* *	286	7.0(56)	1.4 (6)	27.3(11)	161	21.7(10)	25.5(11)	261	12.6 (83)
Pendleton	62	* *	* *	378	8.2(75)	1.6(10)	17.2(46)	195	19.0(72)	13.8(20)	316	8.9 (45)
Perry	314	9.2 (33)	5.4 (11)	1,265	8.9(87)	1.3 (3)	32.5(11)	642	19.0(72)	22.4(10)	858	12.7 (84)
Pike	433	9.9 (40)	6.9 (29)	2,041	3.4 (3)	4.9(95)	19.0(74)	998	14.2(19)	21.4(95)	1,163	8.2 (37)
Powell	115	7.8 (17)	6.1 (19)	725	5.9(36)	1.5 (8)	19.2(76)	402	19.9(87)	22.1(10)	621	6.9 (23)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2003 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Pulaski	413	13.3 (71)	4.8 (9)	2,006	5.1(30)	4.6(84)	12.7 (9)	951	17.4(54)	12.5(12)	1,002	8.5 (41)
Robertson	10	* *	* *	115	3.5 (5)	0.9 (1)	17.4(49)	69	* *	* *	108	6.5 (20)
Rockcastle	118	8.5 (25)	4.2 (6)	650	9.8(98)	4.9(95)	19.5(77)	337	15.7(30)	13.1(15)	490	18.0(100)
Rowan	146	6.2 (5)	9.6 (60)	669	4.2 (9)	4.8(93)	17.2(46)	340	16.2(41)	17.4(63)	475	10.7 (65)
Russell	104	11.5 (65)	9.6 (60)	605	6.9(54)	2.3(20)	18.7(71)	309	18.8(69)	21.0(90)	315	11.7 (74)
Scott	180	8.3 (21)	9.4 (58)	700	9.3(92)	2.7(35)	19.6(79)	298	15.8(33)	16.8(52)	488	13.1 (87)
Shelby	184	10.3 (47)	9.2 (57)	801	14.6(11)	1.5 (8)	32.3(11)	350	22.6(10)	30.0(11)	599	7.7 (33)
Simpson	97	* *	* *	452	8.6(81)	2.9(38)	19.9(81)	228	19.3(77)	21.9(98)	259	21.6(111)
Spencer	62	* *	* *	274	10.6(10)	4.4(79)	21.5(96)	132	27.3(11)	14.4(23)	156	10.3 (58)
Taylor	151	7.9 (18)	15.2 (76)	812	7.4(62)	4.6(84)	15.8(30)	437	20.4(93)	19.9(82)	366	3.0 (3)
Todd	98	* *	* *	468	8.8(86)	4.3(74)	14.5(19)	253	11.5 (7)	12.3(11)	358	17.9 (99)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2003 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)													
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)			
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% >95th (Rank)	Number	% (Rank)
Trigg	68	* *	* *	321	8.7(83)	2.8(36)	15.9(32)	145	17.2(51)	14.5(25)		214	8.4 (39)
Trimble	54	* *	* *	240	4.2 (9)	9.6(11)	11.7 (5)	114	8.8 (2)	14.0(21)		185	5.9 (15)
Union	83	* *	* *	474	10.3(10)	1.9(15)	16.5(39)	245	22.0(10)	12.2(10)		381	4.2 (8)
Warren	565	7.3 (15)	7.4 (32)	2,123	5.5(33)	3.2(42)	17.1(45)	901	18.8(69)	17.5(66)		1,139	11.5 (71)
Washington	80	* *	* *	423	10.4(10)	2.8(36)	24.1(10)	243	16.5(45)	24.3(10)		266	10.5 (62)
Wayne	162	12.3 (70)	8.0 (41)	928	6.1(41)	3.8(60)	17.7(55)	518	18.5(66)	17.0(55)		490	16.9 (98)
Webster	98	* *	* *	454	11.5(11)	4.8(93)	18.7(71)	209	13.4(15)	17.2(58)		346	19.7(105)
Whitley	464	10.1 (45)	11.4 (72)	2,394	8.3(76)	3.6(56)	20.6(88)	1,281	19.0(72)	21.9(98)		1,846	5.9 (15)
Wolfe	76	* *	* *	438	8.9(87)	1.4 (6)	24.7(11)	247	15.4(27)	21.5(97)		371	5.7 (14)
Kentucky	26,648	9.6	7.4	121,974	7.4	4.0	17.7	60,984	17.8	17.2		79,078	11.5

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2003 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number
Nation (Prior Year)	1,208,400	8.9	7.6	4,913,316	6.3	5.4	13.5	2,069,605	15.4	14.3	3,295,974	13.1

- (1) Reporting period is January 1 through December 31.
- (2) Analyses based on one record per child.
- (3) Excludes records with unknown data or errors.
- (4) Rank compares this county's rate to other counties. Rank 1 = best rate.
- (5) Infants born during the reporting period included in the analysis.
- (6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
- (7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
- (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7B
2003 Pediatric Nutrition Surveillance (1)
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County (4,5)						
County	Breastfeeding (6)					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Adair	102	17.6 (48)	48	* *	95	* *
Allen	122	27.9 (11)	49	* *	126	4.0 (60)
Anderson	74	* *	23	* *	49	* *
Ballard	57	* *	30	* *	53	* *
Barren	289	29.8 (10)	114	6.1 (27)	203	6.9 (24)
Bath	150	24.0 (21)	70	* *	156	3.8 (64)
Bell	288	11.1 (72)	153	7.8 (23)	318	5.7 (36)
Boone	514	17.5 (50)	225	16.0 (4)	361	11.6 (3)
Bourbon	109	18.3 (46)	40	* *	96	* *
Boyd	343	21.0 (34)	140	10.7 (13)	251	5.2 (42)
Boyle	151	17.2 (52)	63	* *	135	1.5 (75)
Bracken	52	* *	21	* *	41	* *
Breathitt	127	3.9 (78)	71	* *	128	7.8 (13)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7B
2003 Pediatric Nutrition Surveillance (1)
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County (4,5)						
County	Breastfeeding (6)					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Breckinrid	114	16.7 (54)	55	* *	85	* *
Bullitt	307	26.4 (14)	128	10.2 (15)	238	5.0 (46)
Butler	119	17.6 (48)	39	* *	73	* *
Caldwell	65	* *	49	* *	88	* *
Calloway	183	37.2 (2)	104	23.1 (1)	190	13.2 (2)
Campbell	385	16.4 (57)	151	9.3 (19)	227	6.2 (33)
Carlisle	33	* *	15	* *	36	* *
Carroll	95	* *	43	* *	76	* *
Carter	200	22.0 (29)	131	6.1 (27)	254	4.3 (54)
Casey	90	* *	50	* *	107	6.5 (29)
Christian	971	27.6 (12)	462	13.6 (9)	609	5.7 (36)
Clark	192	21.4 (31)	87	* *	151	4.6 (52)
Clay	222	18.5 (44)	104	5.8 (29)	237	2.5 (69)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7B
2003 Pediatric Nutrition Surveillance (1)
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County (4,5)						
County	Breastfeeding (6)					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Clinton	84	* *	52	* *	105	1.9 (74)
Crittenden	53	* *	28	* *	66	* *
Cumberland	54	* *	28	* *	52	* *
Daviess	615	13.8 (66)	261	5.7 (30)	411	5.1 (43)
Edmonson	94	* *	41	* *	60	* *
Elliott	66	* *	27	* *	45	* *
Estill	114	16.7 (54)	57	* *	112	8.0 (12)
Fayette	1,322	42.2 (1)	556	19.2 (3)	897	11.6 (3)
Fleming	89	* *	52	* *	118	6.8 (26)
Floyd	352	10.2 (74)	168	10.1 (17)	353	5.7 (36)
Franklin	260	23.1 (24)	98	* *	153	11.1 (5)
Fulton	50	* *	39	* *	78	* *
Gallatin	61	* *	27	* *	59	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7B
2003 Pediatric Nutrition Surveillance (1)
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County (4,5)						
County	Breastfeeding (6)					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Garrard	107	19.6 (40)	57	* *	113	5.3 (41)
Grant	207	24.6 (18)	79	* *	177	5.1 (43)
Graves	241	34.4 (4)	124	13.7 (8)	226	7.1 (23)
Grayson	173	26.0 (16)	82	* *	192	3.6 (67)
Green	77	* *	34	* *	73	* *
Greenup	222	18.9 (42)	118	15.3 (5)	158	8.9 (9)
Hancock	45	* *	26	* *	41	* *
Hardin	663	14.8 (62)	345	13.9 (6)	533	10.5 (7)
Harlan	276	14.1 (65)	159	6.3 (26)	304	3.9 (63)
Harrison	122	36.1 (3)	46	* *	97	* *
Hart	111	19.8 (39)	72	* *	138	5.1 (43)
Henderson	249	23.7 (23)	137	8.0 (22)	186	3.8 (64)
Henry	111	22.5 (27)	42	* *	64	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7B
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County ^(4,5)						
County	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Hickman	30	* *	19	* *	27	* *
Hopkins	296	18.6 (43)	127	10.2 (15)	192	4.2 (57)
Jackson	114	14.9 (61)	61	* *	116	4.3 (54)
Jefferson	3,909	25.6 (17)	1,476	13.8 (7)	2,436	7.7 (16)
Jessamine	225	20.4 (38)	117	22.2 (2)	173	11.0 (6)
Johnson	185	16.2 (58)	109	10.1 (17)	203	5.4 (40)
Kenton	547	19.6 (40)	232	5.6 (32)	340	3.5 (68)
Knott	108	12.0 (69)	83	* *	122	4.1 (59)
Knox	280	10.4 (73)	129	9.3 (19)	233	4.7 (48)
Larue	105	21.0 (34)	37	* *	67	* *
Laurel	379	17.7 (47)	175	5.7 (30)	254	4.7 (48)
Lawrence	107	11.2 (71)	56	* *	109	7.3 (19)
Lee	62	* *	29	* *	57	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7B
2003 Pediatric Nutrition Surveillance (1)
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County (4,5)						
County	Breastfeeding (6)					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Leslie	92	* *	67	* *	112	6.3 (32)
Letcher	171	12.3 (68)	80	* *	138	14.5 (1)
Lewis	85	* *	51	* *	115	4.3 (54)
Lincoln	156	24.4 (20)	81	* *	118	4.2 (57)
Livingston	43	* *	22	* *	36	* *
Logan	157	14.6 (63)	71	* *	140	2.1 (72)
Lyon	22	* *	20	* *	30	* *
Madison	453	23.0 (25)	220	10.9 (12)	340	7.6 (17)
Magoffin	129	7.0 (77)	57	* *	135	2.2 (70)
Marion	122	24.6 (18)	60	* *	103	7.8 (13)
Marshall	153	32.7 (6)	70	* *	183	10.4 (8)
Martin	117	9.4 (75)	66	* *	128	4.7 (48)
Mason	90	* *	56	* *	126	4.0 (60)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7B
2003 Pediatric Nutrition Surveillance (1)
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County (4,5)						
County	Breastfeeding (6)					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
McCracken	367	33.5 (5)	232	10.3 (14)	439	7.3 (19)
McCreary	196	13.8 (66)	75	* *	164	6.1 (34)
McLean	60	* *	33	* *	77	* *
Meade	154	22.1 (28)	69	* *	98	* *
Menifee	72	* *	30	* *	58	* *
Mercer	134	17.2 (52)	70	* *	110	4.5 (53)
Metcalfe	67	* *	48	* *	73	* *
Monroe	65	* *	36	* *	60	* *
Montgomery	178	18.5 (44)	99	* *	186	8.6 (10)
Morgan	93	* *	48	* *	104	3.8 (64)
Muhlenberg	193	11.4 (70)	77	* *	152	2.0 (73)
Nelson	234	20.9 (36)	122	7.4 (24)	201	4.0 (60)
Nicholas	46	* *	29	* *	60	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7B
2003 Pediatric Nutrition Surveillance (1)
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County (4,5)						
County	Breastfeeding (6)					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Ohio	175	26.9 (13)	77	* *	152	7.2 (21)
Oldham	144	32.6 (7)	74	* *	130	6.9 (24)
Owen	56	* *	22	* *	66	* *
Owsley	35	* *	24	* *	46	* *
Pendleton	62	* *	35	* *	75	* *
Perry	319	8.2 (76)	122	7.4 (24)	263	5.7 (36)
Pike	434	16.1 (59)	213	4.7 (33)	339	4.7 (48)
Powell	115	26.1 (15)	70	* *	160	7.5 (18)
Pulaski	414	23.9 (22)	215	11.6 (11)	344	6.7 (27)
Robertson	10	* *	16	* *	22	* *
Rockcastle	121	30.6 (8)	75	* *	109	6.4 (30)
Rowan	146	14.4 (64)	73	* *	118	8.5 (11)
Russell	104	17.3 (51)	68	* *	125	6.4 (30)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7B
2003 Pediatric Nutrition Surveillance (1)
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County (4,5)						
County	Breastfeeding (6)					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Scott	181	30.4 (9)	75	* *	139	2.2 (70)
Shelby	184	20.7 (37)	91	* *	152	7.2 (21)
Simpson	101	22.8 (26)	43	* *	67	* *
Spencer	62	* *	32	* *	49	* *
Taylor	151	21.9 (30)	78	* *	138	5.8 (35)
Todd	98	* *	30	* *	80	* *
Trigg	68	* *	33	* *	64	* *
Trimble	54	* *	16	* *	46	* *
Union	84	* *	50	* *	71	* *
Warren	584	21.2 (32)	226	12.8 (10)	359	7.8 (13)
Washington	80	* *	42	* *	68	* *
Wayne	163	16.0 (60)	80	* *	164	4.9 (47)
Webster	98	* *	56	* *	82	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7B
2003 Pediatric Nutrition Surveillance (1)
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County (4,5)						
County	Breastfeeding (6)					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Whitley	465	16.6 (56)	224	8.9 (21)	333	6.6 (28)
Wolfe	77	* *	29	* *	72	* *
Unknown Counties	1,168	21.1 (33)	0	* *	0	* *
Kentucky	26,921	22.2	11,948	10.5	20,842	6.4
Nation (Prior Year)	969,228	52.5	527,765	20.8	555,990	12.3

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2001 - 2003 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Adair	384	7.6 (22)	9.1 (91)	1,476	5.0(27)	5.6(93)	12.9(11)	664	13.3(12)	13.0(18)	827	7.0 (21)
Allen	396	9.3 (52)	10.6(109)	1,871	6.0(44)	3.1(19)	16.3(57)	962	20.2(10)	17.3(72)	1,135	9.7 (57)
Anderson	265	8.3 (34)	7.5 (54)	1,042	8.4(98)	4.4(68)	20.2(10)	493	20.1(10)	21.7(10)	806	8.6 (46)
Ballard	196	12.8(114)	9.7 (98)	807	6.4(57)	8.1(11)	13.6(18)	406	10.1 (5)	10.8 (6)	631	6.3 (15)
Barren	848	9.1 (47)	7.9 (65)	3,213	7.2(66)	4.5(70)	17.0(70)	1,403	16.0(38)	16.9(69)	2,029	10.8 (65)
Bath	353	8.2 (32)	12.2(117)	1,728	9.7(10)	2.8(13)	21.2(10)	895	17.8(70)	19.8(98)	945	9.6 (54)
Bell	975	8.7 (38)	7.6 (56)	4,963	6.1(48)	3.9(45)	17.9(80)	2,654	16.2(41)	16.0(58)	3,296	7.9 (40)
Boone	1,542	9.1 (47)	9.4 (96)	5,777	7.5(78)	4.6(72)	16.7(65)	2,477	18.7(83)	15.0(45)	4,010	13.9 (89)
Bourbon	396	11.4 (99)	9.1 (91)	1,502	7.5(78)	2.4 (8)	19.8(99)	665	19.4(99)	20.6(10)	1,094	28.9(120)
Boyd	1,155	11.5(102)	7.7 (59)	4,090	6.9(63)	8.6(11)	13.6(18)	1,807	17.5(64)	14.8(41)	2,577	12.1 (72)
Boyle	505	10.1 (72)	6.9 (36)	1,681	6.0(44)	4.9(80)	16.2(56)	672	15.6(31)	17.6(78)	1,020	7.6 (35)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2001 - 2003 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Bracken	176	13.1 (115)	6.8 (32)	686	9.0(10)	6.4(10)	22.3(11)	323	16.1(39)	27.9(11)	475	7.2 (25)
Breathitt	441	9.5 (57)	8.4 (80)	2,045	5.2(31)	4.3(62)	18.4(84)	1,153	14.7(21)	17.7(79)	1,515	7.6 (35)
Breckinrid	339	8.0 (29)	8.0 (68)	1,604	6.7(62)	3.2(23)	14.7(34)	785	18.3(77)	13.1(20)	1,235	3.1 (2)
Bullitt	1,034	7.4 (17)	8.4 (80)	3,271	7.0(65)	3.9(45)	16.0(52)	1,317	16.5(45)	14.7(40)	1,814	19.6(113)
Butler	350	7.4 (17)	8.0 (68)	1,433	4.6(18)	3.5(33)	18.9(90)	701	19.0(91)	22.7(11)	887	7.1 (23)
Caldwell	283	6.7 (5)	7.8 (63)	1,237	6.1(48)	6.8(10)	16.9(68)	564	14.7(21)	15.8(55)	949	13.2 (81)
Calloway	681	7.8 (26)	11.5(114)	2,743	5.8(41)	3.0(17)	14.8(36)	1,248	17.2(59)	15.8(55)	1,752	19.3(112)
Campbell	1,139	10.0 (69)	7.7 (59)	3,918	7.4(74)	3.8(43)	16.7(65)	1,698	19.3(96)	18.5(85)	2,539	14.5 (96)
Carlisle	120	4.2 (1)	10.0(104)	618	5.7(40)	3.4(28)	23.8(11)	336	19.3(96)	27.1(11)	396	19.9(114)
Carroll	323	10.5 (80)	4.6 (2)	1,243	10.5(11)	4.7(74)	19.1(92)	537	22.0(11)	19.9(99)	922	13.4 (83)
Carter	810	7.2 (14)	11.7(115)	3,745	4.0 (8)	7.2(11)	12.5(10)	1,910	15.9(36)	12.5(13)	2,276	9.9 (61)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2001 - 2003 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low		High	Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Casey	338	10.1 (72)	5.3 (6)	1,583	7.9(88)	4.4(68)	13.8(21)	780	15.8(35)	15.3(49)	926	10.5 (64)
Christian	3,519	8.6 (37)	8.8 (86)	14,678	5.3(32)	6.8(10)	10.4 (3)	6,605	12.5 (9)	9.3 (4)	9,529	25.3(119)
Clark	693	11.8(107)	6.9 (36)	2,392	7.3(69)	3.6(36)	15.9(51)	1,012	18.6(81)	17.2(71)	1,523	11.4 (71)
Clay	767	10.7 (86)	7.2 (47)	3,503	5.9(42)	6.5(10)	18.5(85)	1,853	17.6(68)	18.9(91)	2,636	7.6 (35)
Clinton	307	9.8 (63)	8.8 (86)	1,502	4.8(21)	4.9(80)	17.4(74)	763	16.4(43)	13.9(27)	903	7.2 (25)
Crittenden	174	8.0 (29)	5.7 (14)	799	4.3(12)	5.5(90)	13.1(14)	363	13.2(11)	11.0 (7)	563	13.3 (82)
Cumberland	194	6.7 (5)	6.7 (30)	912	3.9 (7)	4.5(70)	13.8(21)	462	14.7(21)	14.9(42)	445	17.8(110)
Daviess	2,180	8.9 (45)	6.1 (24)	7,955	9.3(10)	2.1 (7)	19.5(96)	3,527	17.1(56)	17.5(77)	5,532	12.6 (76)
Edmonson	297	11.8(107)	7.7 (59)	1,148	4.3(12)	5.4(89)	11.5 (5)	494	15.6(31)	12.8(15)	701	5.4 (9)
Elliott	217	8.8 (42)	12.0(116)	993	3.3 (4)	7.0(11)	13.5(16)	519	14.5(19)	13.9(27)	624	13.1 (80)
Estill	413	7.0 (12)	9.4 (96)	1,799	4.2(11)	4.3(62)	15.5(45)	907	18.4(79)	18.7(88)	1,396	8.5 (45)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2001 - 2003 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Fayette	4,618	10.7 (86)	7.2 (47)	14,774	6.2(52)	4.3(62)	14.7(34)	5,858	18.6(81)	16.8(65)	8,767	15.8 (99)
Fleming	358	7.3 (16)	10.3(106)	1,795	5.4(35)	4.7(74)	15.0(39)	928	13.9(17)	15.2(48)	1,431	16.2(101)
Floyd	1,230	9.1 (47)	7.9 (65)	5,471	4.3(12)	9.3(12)	13.8(21)	2,758	11.3 (6)	13.4(22)	3,905	8.7 (48)
Franklin	903	8.9 (45)	7.0 (38)	2,942	8.3(97)	6.2(99)	18.0(82)	1,119	19.4(99)	19.9(99)	2,068	14.5 (96)
Fulton	237	11.0 (91)	6.8 (32)	1,110	7.3(69)	4.0(50)	19.0(91)	560	21.3(11)	18.4(84)	766	14.2 (95)
Gallatin	208	10.6 (81)	7.2 (47)	810	8.0(92)	4.2(61)	14.1(29)	373	12.6(10)	16.4(62)	537	13.6 (88)
Garrard	355	9.3 (52)	7.6 (56)	1,625	6.6(60)	3.3(25)	14.1(29)	806	18.9(88)	14.0(32)	953	13.9 (89)
Grant	654	9.8 (63)	9.8 (99)	3,092	4.7(19)	6.2(99)	10.2 (2)	1,573	9.7 (3)	8.0 (1)	2,465	5.7 (11)
Graves	841	10.6 (81)	8.0 (68)	3,507	7.2(66)	5.8(96)	16.9(68)	1,693	19.1(93)	17.4(73)	2,266	16.8(105)
Grayson	594	7.2 (14)	8.4 (80)	2,729	11.4(11)	2.9(14)	20.4(10)	1,337	16.8(51)	17.4(73)	2,092	2.9 (1)
Green	260	6.9 (8)	10.0(104)	1,016	7.9(88)	2.5(10)	22.1(11)	468	20.3(10)	21.8(10)	598	7.5 (31)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2001 - 2003 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Greenup	719	11.4 (99)	7.6 (56)	2,928	3.1 (1)	8.3(11)	12.9(11)	1,370	16.9(53)	12.9(17)	1,905	7.4 (30)
Hancock	149	5.4 (2)	4.7 (3)	777	5.5(36)	3.2(23)	16.1(53)	392	16.6(47)	15.1(47)	631	7.3 (27)
Hardin	2,498	9.3 (52)	7.1 (42)	10,358	6.4(57)	2.7(12)	14.0(26)	4,780	19.5(10)	13.4(22)	7,886	7.5 (31)
Harlan	1,027	8.1 (31)	10.7(111)	4,972	4.3(12)	7.1(11)	18.5(85)	2,671	16.4(43)	20.8(10)	3,425	7.1 (23)
Harrison	395	12.4(113)	5.8 (18)	1,579	4.7(19)	5.7(94)	10.5 (4)	744	9.8 (4)	8.5 (2)	1,091	14.1 (94)
Hart	434	7.1 (13)	8.3 (79)	1,832	5.1(29)	4.0(50)	15.8(50)	849	15.1(28)	18.5(85)	1,228	6.8 (19)
Henderson	931	11.2 (96)	5.4 (8)	3,447	8.1(95)	5.5(90)	16.1(53)	1,471	22.2(11)	22.1(11)	915	5.2 (7)
Henry	317	8.8 (42)	5.7 (14)	1,169	12.5(11)	3.6(36)	23.4(11)	489	18.2(76)	19.6(97)	777	15.2 (98)
Hickman	100	12.0(111)	6.0 (22)	390	3.8 (5)	4.9(80)	12.3 (8)	175	16.6(47)	14.9(42)	276	8.7 (48)
Hopkins	1,050	11.0 (91)	5.6 (13)	4,370	11.9(11)	1.8 (4)	20.2(10)	2,014	20.3(10)	18.7(88)	3,219	13.0 (78)
Jackson	416	10.1 (72)	9.9(103)	1,949	5.3(32)	3.4(28)	17.2(72)	1,000	14.9(25)	15.6(54)	1,553	3.5 (3)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2001 - 2003 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Jefferson	12,580	10.8 (89)	6.8 (32)	39,655	7.6(80)	5.2(85)	15.0(39)	16,067	16.6(47)	14.9(42)	25,097	12.1 (72)
Jessamine	797	7.4 (17)	9.8 (99)	3,106	8.6(99)	3.4(28)	16.3(57)	1,373	16.9(53)	15.4(52)	2,125	10.4 (63)
Johnson	669	7.5 (20)	8.8 (86)	3,158	4.0 (8)	6.6(10)	16.1(53)	1,628	15.3(30)	15.3(49)	2,321	19.2(111)
Kenton	1,827	11.3 (97)	6.2 (27)	6,604	7.3(69)	4.1(54)	18.6(87)	2,745	21.9(11)	23.2(11)	4,835	13.4 (83)
Knott	401	10.2 (75)	5.5 (11)	2,190	10.5(11)	4.7(74)	20.1(10)	1,196	13.7(14)	14.0(32)	1,842	8.3 (43)
Knox	964	10.6 (81)	8.5 (83)	4,509	4.9(25)	3.7(42)	17.5(77)	2,344	19.1(93)	19.0(92)	3,290	13.4 (83)
Larue	332	6.9 (8)	5.7 (14)	1,229	8.2(96)	4.3(62)	19.6(97)	565	17.7(69)	15.4(52)	957	9.3 (52)
Laurel	1,382	10.9 (90)	8.0 (68)	4,787	6.0(44)	5.8(96)	15.3(44)	1,982	19.7(10)	18.1(82)	2,763	13.9 (89)
Lawrence	419	13.1 (115)	5.3 (6)	1,789	5.1(29)	5.7(94)	18.3(83)	875	19.9(10)	17.7(79)	1,198	7.5 (31)
Lee	190	7.9 (28)	10.5(107)	1,057	7.6(80)	1.5 (1)	24.9(11)	560	20.2(10)	22.9(11)	863	16.7(103)
Leslie	363	7.7 (23)	10.5(107)	1,938	3.2 (3)	3.6(36)	21.4(10)	1,099	18.7(83)	21.9(11)	1,623	8.2 (42)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2001 - 2003 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Letcher	603	8.8 (42)	5.5 (11)	2,610	8.6(99)	3.3(25)	22.0(11)	1,244	16.1(39)	19.5(96)	2,020	3.5 (3)
Lewis	348	10.3 (77)	9.2 (93)	1,554	5.5(36)	4.7(74)	14.3(32)	774	21.2(11)	16.0(58)	932	5.6 (10)
Lincoln	607	9.9 (67)	7.7 (59)	2,527	7.7(84)	4.9(80)	19.1(92)	1,193	18.8(86)	19.4(94)	1,931	6.9 (20)
Livingston	142	11.3 (97)	4.9 (4)	658	4.9(25)	6.8(10)	15.5(45)	324	15.1(28)	13.9(27)	531	5.3 (8)
Logan	557	9.5 (57)	5.7 (14)	2,138	6.0(44)	3.3(25)	16.5(61)	919	21.7(11)	16.4(62)	1,306	20.4(115)
Lyon	99	* *	* *	511	7.6(80)	5.1(84)	19.6(97)	254	16.9(53)	18.1(82)	366	9.6 (54)
Madison	1,549	9.6 (60)	10.7(111)	5,541	7.7(84)	6.2(99)	13.6(18)	2,334	18.9(88)	18.8(90)	3,926	10.9 (66)
Magoffin	439	10.0 (69)	9.8 (99)	2,304	7.4(74)	4.1(54)	22.9(11)	1,253	17.8(70)	20.2(10)	1,481	22.7(117)
Marion	424	9.9 (67)	5.9 (19)	1,990	10.8(11)	3.6(36)	21.7(10)	971	17.9(73)	17.4(73)	1,424	10.2 (62)
Marshall	599	8.3 (34)	7.5 (54)	2,477	7.8(86)	6.5(10)	15.1(41)	1,133	12.0 (7)	8.7 (3)	1,797	7.3 (27)
Martin	429	11.4 (99)	6.1 (24)	2,144	6.3(55)	4.1(54)	17.5(77)	1,153	21.0(11)	16.8(65)	1,753	6.6 (18)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2001 - 2003 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years ⁽²⁾

Comparison of Growth and Anemia Indicators by County ^(3,4)												
County	Birthweight ⁽⁵⁾			Height and Weight						Anemia Low Hb/Hct ⁽⁸⁾		
	Low	High		Short Stature ⁽⁶⁾	Under- weight ⁽⁷⁾	Over- weight ⁽⁷⁾		≥ 2 Yrs Over- weight ⁽⁷⁾				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Mason	401	8.2 (32)	8.2 (74)	1,647	9.1(10)	6.4(10)	16.7(65)	772	17.1(56)	13.1(20)	1,133	13.0 (78)
McCracken	1,437	11.6(105)	7.1 (42)	5,618	7.3(69)	5.2(85)	19.4(95)	2,556	17.3(61)	15.8(55)	4,172	20.4(115)
McCreary	597	11.9(110)	8.7 (84)	2,795	6.6(60)	4.8(78)	14.0(26)	1,431	15.9(36)	13.8(25)	2,149	7.5 (31)
McLean	250	9.6 (60)	6.0 (22)	1,289	11.6(11)	2.6(11)	21.6(10)	652	17.5(64)	24.8(11)	989	3.6 (5)
Meade	523	7.5 (20)	7.3 (51)	2,061	9.4(10)	4.1(54)	17.1(71)	983	18.7(83)	16.2(61)	1,555	15.8 (99)
Menifee	197	9.1 (47)	8.1 (73)	942	4.8(21)	4.0(50)	11.7 (6)	483	14.7(21)	13.9(27)	572	9.8 (59)
Mercer	462	11.0 (91)	6.5 (29)	1,826	8.0(92)	3.5(33)	15.1(41)	851	17.9(73)	17.4(73)	1,349	8.8 (50)
Metcalfe	254	13.4(117)	10.6(109)	1,065	4.5(17)	4.1(54)	12.4 (9)	499	17.2(59)	14.6(39)	687	9.8 (59)
Monroe	272	7.7 (23)	8.8 (86)	1,222	6.3(55)	3.4(28)	14.2(31)	552	15.6(31)	14.1(36)	819	14.0 (92)
Montgomery	632	10.3 (77)	9.3 (94)	2,626	6.2(52)	3.1(19)	13.4(15)	1,226	15.7(34)	12.6(14)	2,021	9.6 (54)
Morgan	318	6.9 (8)	8.2 (74)	1,747	4.8(21)	4.3(62)	21.5(10)	947	19.9(10)	21.1(10)	1,437	17.1(106)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2001 - 2003 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Muhlenberg	740	8.4 (36)	8.0 (68)	2,954	3.8 (5)	6.8(10)	12.9(11)	1,413	13.8(16)	14.4(38)	1,768	10.9 (66)
Nelson	766	10.6 (81)	5.9 (19)	3,229	8.0(92)	4.3(62)	16.3(57)	1,482	16.3(42)	12.1(10)	2,090	9.7 (57)
Nicholas	191	11.5 (102)	5.2 (5)	843	5.5(36)	8.4(11)	13.5(16)	410	9.3 (1)	11.2 (8)	661	17.1 (106)
Ohio	587	9.2 (51)	8.2 (74)	2,493	5.3(32)	3.5(33)	14.4(33)	1,242	17.1(56)	16.1(60)	1,897	9.5 (53)
Oldham	457	12.3 (112)	6.1 (24)	1,825	10.3(10)	2.4 (8)	17.9(80)	846	17.5(64)	16.8(65)	1,130	13.5 (87)
Owen	205	10.7 (86)	5.4 (8)	843	5.0(27)	5.3(87)	13.8(21)	404	16.8(51)	14.1(36)	633	7.0 (21)
Owsley	153	6.5 (3)	7.8 (63)	822	7.4(74)	1.9 (6)	27.0(11)	445	21.6(11)	23.4(11)	693	16.5 (102)
Pendleton	285	9.5 (57)	7.0 (38)	1,148	7.4(74)	3.4(28)	14.0(26)	523	18.0(75)	13.8(25)	869	8.3 (43)
Perry	1,066	10.4 (79)	7.0 (38)	3,605	8.9(10)	1.6 (3)	32.3(11)	1,694	18.8(86)	22.4(11)	2,580	8.6 (46)
Pike	1,579	9.4 (56)	7.1 (42)	6,214	3.1 (1)	8.0(11)	17.3(73)	2,886	17.3(61)	20.2(10)	4,021	7.7 (38)
Powell	435	8.7 (38)	7.4 (53)	2,022	4.3(12)	2.9(14)	15.7(48)	1,051	17.3(61)	16.9(69)	1,623	6.5 (17)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2001 - 2003 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years ⁽²⁾

Comparison of Growth and Anemia Indicators by County ^(3,4)												
County	Birthweight ⁽⁵⁾			Height and Weight							Anemia Low Hb/Hct ⁽⁸⁾	
	Low	High		Short Stature ⁽⁶⁾	Under- weight ⁽⁷⁾	Over- weight ⁽⁷⁾		≥ 2 Yrs Over- weight ⁽⁷⁾				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Pulaski	1,516	10.0 (69)	6.7 (30)	5,653	5.6(39)	5.3(87)	11.9 (7)	2,452	14.9(25)	12.3(11)	3,219	8.0 (41)
Robertson	63	* *	* *	327	7.6(80)	3.1(19)	16.5(61)	172	12.2 (8)	14.0(32)	276	5.8 (12)
Rockcastle	448	6.7 (5)	7.1 (42)	1,841	7.3(69)	6.0(98)	17.4(74)	872	16.6(47)	12.8(15)	1,324	17.5(109)
Rowan	508	7.7 (23)	13.4(118)	1,998	4.0 (8)	4.6(72)	15.1(41)	902	16.5(45)	15.3(49)	1,333	14.0 (92)
Russell	428	9.3 (52)	8.2 (74)	1,779	6.9(63)	3.1(19)	17.4(74)	777	18.3(77)	19.4(94)	1,041	9.0 (51)
Scott	634	8.7 (38)	7.3 (51)	2,005	7.8(86)	5.5(90)	17.7(79)	789	13.3(12)	13.6(24)	1,371	12.6 (76)
Shelby	652	9.8 (63)	8.9 (90)	2,270	16.2(12)	1.8 (4)	33.8(12)	889	22.0(11)	30.3(12)	1,625	11.0 (68)
Simpson	368	9.8 (63)	5.4 (8)	1,329	7.9(88)	3.6(36)	16.6(64)	547	22.5(12)	19.2(93)	810	22.7(117)
Spencer	207	13.5(118)	7.2 (47)	777	11.6(11)	3.6(36)	19.3(94)	337	21.4(11)	13.9(27)	508	11.0 (68)
Taylor	539	6.9 (8)	9.8 (99)	2,292	6.1(48)	4.1(54)	15.5(45)	1,117	19.2(95)	16.6(64)	1,296	3.9 (6)
Todd	321	11.5(102)	5.9 (19)	1,380	7.9(88)	4.0(50)	14.8(36)	665	13.7(14)	10.1 (5)	1,056	17.2(108)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2001 - 2003 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Trigg	216	11.6 (105)	7.9 (65)	933	6.4(57)	3.9(45)	13.8(21)	424	14.9(25)	12.3(11)	666	7.7 (38)
Trimble	188	9.6 (60)	6.4 (28)	703	6.1(48)	8.4(11)	10.1 (1)	322	9.3 (1)	13.0(18)	530	5.8 (12)
Union	359	11.1 (94)	2.8 (1)	1,481	10.5(11)	3.0(17)	15.7(48)	689	19.3(96)	11.9 (9)	1,122	6.4 (16)
Warren	2,017	10.2 (75)	7.0 (38)	6,490	6.2(52)	4.1(54)	14.8(36)	2,508	17.8(70)	15.0(45)	3,768	12.5 (75)
Washington	260	6.5 (3)	10.8(113)	1,161	9.1(10)	2.9(14)	19.8(99)	565	17.5(64)	16.8(65)	817	11.0 (68)
Wayne	575	10.6 (81)	8.2 (74)	2,608	4.8(21)	3.9(45)	16.4(60)	1,321	18.5(80)	18.5(85)	1,616	12.2 (74)
Webster	323	11.8(107)	6.8 (32)	1,238	10.3(10)	3.9(45)	18.6(87)	528	14.0(18)	18.0(81)	910	16.7(103)
Whitley	1,585	8.7 (38)	9.3 (94)	6,722	7.2(66)	3.8(43)	18.6(87)	3,264	19.0(91)	20.3(10)	5,021	6.2 (14)
Wolfe	281	7.8 (26)	7.1 (42)	1,320	9.0(10)	1.5 (1)	23.3(11)	697	18.9(88)	20.9(10)	1,065	7.3 (27)
Woodford	287	11.1 (94)	8.7 (84)	1,021	5.9(42)	4.8(78)	16.5(61)	408	14.5(19)	14.0(32)	674	13.4 (83)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2001 - 2003 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number
Kentucky	93,533	9.7	7.7	354,077	6.7	4.8	16.2	160,602	17.1	16.2	236,889	11.8

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7B
2001 - 2003 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County ^(4,5)						
County	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Adair	522	9.8 (88)	304	2.0 (90)	298	1.7 (71)
Allen	543	18.8 (13)	333	2.4 (72)	342	1.5 (82)
Anderson	338	21.0 (4)	204	4.9 (9)	182	2.2 (47)
Ballard	268	12.3 (59)	162	1.9 (96)	167	0.6(109)
Barren	1,164	17.2 (21)	692	4.0 (23)	621	2.6 (30)
Bath	589	9.3 (96)	355	1.7(104)	348	1.7 (71)
Bell	1,459	8.9(100)	961	2.6 (65)	960	2.1 (48)
Boone	2,173	11.8 (64)	1,291	5.8 (6)	1,127	4.3 (5)
Bourbon	500	12.6 (54)	329	3.6 (33)	325	3.1 (15)
Boyd	1,447	12.1 (62)	800	2.6 (65)	761	1.8 (64)
Boyle	660	8.0(106)	359	0.8(114)	364	0.5(111)
Bracken	234	16.2 (23)	126	3.2 (45)	122	2.5 (35)
Breathitt	606	5.6(116)	361	2.8 (58)	353	2.8 (25)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7B
2001 - 2003 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County ^(4,5)						
County	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Breckinrid	519	10.4 (79)	312	2.6 (65)	292	2.4 (40)
Bullitt	1,292	15.9 (27)	663	2.9 (49)	648	1.9 (55)
Butler	459	9.6 (93)	259	1.2(111)	257	0.4(114)
Caldwell	423	9.9 (86)	283	2.8 (58)	265	1.5 (82)
Calloway	952	23.0 (3)	596	7.0 (2)	565	4.8 (1)
Campbell	1,573	9.3 (96)	866	3.1 (47)	752	2.1 (48)
Carlisle	176	19.9 (7)	119	2.5 (69)	114	1.8 (64)
Carroll	433	12.7 (52)	260	1.9 (96)	257	1.9 (55)
Carter	1,121	14.1 (43)	734	2.6 (65)	722	1.8 (64)
Casey	505	10.9 (72)	311	2.9 (49)	300	2.3 (43)
Christian	4,853	16.4 (22)	3,012	4.9 (9)	2,457	1.9 (55)
Clark	893	12.3 (59)	508	2.8 (58)	461	1.5 (82)
Clay	1,052	9.7 (90)	677	1.3(107)	668	0.9(103)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7B
2001 - 2003 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County ^(4,5)						
County	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Clinton	450	5.3(117)	312	1.3(107)	309	1.0(101)
Crittenden	274	10.2 (82)	177	2.8 (58)	175	2.3 (43)
Cumberland	263	10.6 (76)	173	2.3 (77)	173	1.7 (71)
Daviess	2,769	10.3 (80)	1,603	2.8 (58)	1,407	1.8 (64)
Edmonson	430	16.0 (26)	262	1.1(112)	205	0.5(111)
Elliott	272	9.9 (86)	183	3.3 (40)	173	2.3 (43)
Estill	563	13.1 (50)	336	3.3 (40)	324	3.1 (15)
Fayette	5,827	29.6 (1)	3,224	7.6 (1)	2,863	4.2 (6)
Fleming	502	8.2(104)	311	3.5 (37)	313	3.2 (13)
Floyd	1,709	4.8(119)	1,072	2.1 (84)	1,051	1.9 (55)
Franklin	1,139	19.1 (12)	647	5.3 (7)	526	4.0 (8)
Fulton	367	9.3 (96)	245	3.3 (40)	224	1.8 (64)
Gallatin	281	12.5 (56)	162	2.5 (69)	148	2.7 (28)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7B
2001 - 2003 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County ^(4,5)						
County	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Garrard	540	12.4 (57)	346	2.3 (77)	343	1.7 (71)
Grant	940	12.0 (63)	593	2.2 (80)	596	2.0 (53)
Graves	1,163	19.9 (7)	706	4.5 (14)	681	2.6 (30)
Grayson	825	20.6 (6)	515	2.1 (84)	519	1.3 (95)
Green	345	23.5 (2)	198	4.0 (23)	201	2.5 (35)
Greenup	973	13.2 (49)	584	4.1 (19)	509	3.1 (15)
Hancock	230	15.7 (29)	157	6.4 (4)	137	4.4 (4)
Hardin	3,366	12.4 (57)	1,985	4.7 (13)	1,707	3.6 (10)
Harlan	1,479	9.6 (93)	957	1.8 (98)	923	1.5 (82)
Harrison	546	17.6 (18)	341	2.9 (49)	322	0.9(103)
Hart	611	11.1 (69)	367	4.1 (19)	351	2.0 (53)
Henderson	1,225	11.6 (66)	694	2.0 (90)	590	1.4 (91)
Henry	419	15.8 (28)	257	3.9 (28)	225	1.3 (95)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7B
2001 - 2003 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County ^(4,5)						
County	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Hickman	141	19.9 (7)	79	* *	71	* *
Hopkins	1,416	10.0 (85)	846	2.1 (84)	773	1.0(101)
Jackson	581	9.3 (96)	385	2.1 (84)	380	1.3 (95)
Jefferson	16,471	16.2 (23)	8,720	4.2 (17)	7,780	2.6 (30)
Jessamine	1,092	14.0 (45)	662	6.6 (3)	585	4.6 (2)
Johnson	941	10.6 (76)	613	2.4 (72)	605	1.8 (64)
Kenton	2,326	9.7 (90)	1,412	2.0 (90)	1,213	1.2 (98)
Knott	611	5.1(118)	415	1.7(104)	369	1.4 (91)
Knox	1,334	6.7(111)	825	2.2 (80)	792	1.4 (91)
Larue	447	15.7 (29)	268	3.7 (30)	234	3.4 (11)
Laurel	1,717	10.3 (80)	975	2.4 (72)	854	1.5 (82)
Lawrence	563	8.0(106)	358	3.4 (39)	347	2.9 (22)
Lee	306	7.2(109)	201	4.0 (23)	202	3.0 (21)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7B
2001 - 2003 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County ^(4,5)						
County	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Leslie	521	8.8(101)	355	3.1 (47)	343	2.3 (43)
Letcher	838	10.9 (72)	527	5.9 (5)	469	4.5 (3)
Lewis	481	11.2 (67)	314	2.2 (80)	309	1.6 (77)
Lincoln	849	12.7 (52)	538	2.0 (90)	448	1.6 (77)
Livingston	206	12.6 (54)	137	2.9 (49)	131	1.5 (82)
Logan	764	5.9(115)	470	1.1(112)	443	0.7(105)
Lyon	157	15.3 (33)	110	3.6 (33)	102	2.9 (22)
Madison	1,998	15.1 (35)	1,141	4.5 (14)	1,012	2.9 (22)
Magoffin	638	3.0(121)	420	0.7(116)	421	0.7(105)
Marion	624	13.9 (47)	383	3.7 (30)	361	2.8 (25)
Marshall	850	18.5 (14)	528	4.9 (9)	524	3.8 (9)
Martin	619	4.8(119)	385	2.1 (84)	387	1.6 (77)
Mason	542	12.2 (61)	331	1.8 (98)	333	1.5 (82)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7B
2001 - 2003 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County ^(4,5)						
County	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
McCracken	2,036	15.0 (36)	1,270	3.5 (37)	1,274	3.1 (15)
McCreary	863	8.0(106)	551	2.9 (49)	530	2.1 (48)
McLean	369	14.9 (37)	254	5.1 (8)	255	3.1 (15)
Meade	684	18.0 (16)	407	3.2 (45)	369	1.9 (55)
Menifee	271	17.3 (20)	170	1.8 (98)	176	1.7 (71)
Mercer	600	11.2 (67)	358	3.9 (28)	319	1.9 (55)
Metcalfe	359	8.4(103)	230	2.2 (80)	215	1.9 (55)
Monroe	378	14.3 (41)	252	4.0 (23)	215	2.8 (25)
Montgomery	882	11.7 (65)	547	4.9 (9)	487	3.3 (12)
Morgan	503	10.5 (78)	352	1.7(104)	344	1.5 (82)
Muhlenberg	905	7.2(109)	549	1.3(107)	547	0.5(111)
Nelson	1,075	14.8 (38)	651	2.0 (90)	571	1.4 (91)
Nicholas	271	11.1 (69)	172	2.3 (77)	161	1.2 (98)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7B
2001 - 2003 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County ^(4,5)						
County	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Ohio	803	15.6 (31)	478	2.9 (49)	441	2.5 (35)
Oldham	651	21.0 (4)	388	4.4 (16)	378	2.4 (40)
Owen	277	18.4 (15)	163	1.8 (98)	160	0.6(109)
Owsley	239	6.3(112)	171	1.8 (98)	162	1.9 (55)
Pendleton	382	17.5 (19)	252	2.0 (90)	258	1.6 (77)
Perry	1,277	6.2(114)	702	2.4 (72)	698	2.1 (48)
Pike	2,090	10.2 (82)	1,232	2.1 (84)	1,096	1.6 (77)
Powell	633	9.8 (88)	401	3.7 (30)	401	3.2 (13)
Pulaski	2,119	16.2 (23)	1,256	4.1 (19)	1,106	2.6 (30)
Robertson	102	19.6 (10)	68	* *	65	* *
Rockcastle	621	15.5 (32)	364	4.1 (19)	328	2.1 (48)
Rowan	670	8.2(104)	417	3.6 (33)	379	2.6 (30)
Russell	608	14.0 (45)	394	3.3 (40)	397	2.5 (35)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7B
2001 - 2003 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County ^(4,5)						
County	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Scott	808	15.2 (34)	448	1.3(107)	439	0.7(105)
Shelby	919	10.1 (84)	526	2.9 (49)	484	2.5 (35)
Simpson	475	12.8 (51)	278	1.8 (98)	254	1.2 (98)
Spencer	286	10.8 (74)	166	3.6 (33)	160	1.9 (55)
Taylor	747	14.1 (43)	460	2.8 (58)	451	1.8 (64)
Todd	440	13.6 (48)	274	2.9 (49)	270	1.5 (82)
Trigg	324	19.4 (11)	209	3.3 (40)	196	3.1 (15)
Trimble	248	14.5 (40)	150	2.7 (64)	153	0.7(105)
Union	473	14.2 (42)	302	0.3(118)	266	0.4(114)
Warren	2,535	14.8 (38)	1,338	4.2 (17)	1,153	2.7 (28)
Washington	355	10.7 (75)	226	4.0 (23)	221	4.1 (7)
Wayne	824	8.7(102)	495	2.4 (72)	479	1.7 (71)
Webster	471	9.6 (93)	285	2.5 (69)	231	0.4(114)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7B
2001 - 2003 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County ^(4,5)						
County	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Whitley	2,132	9.7 (90)	1,325	2.9 (49)	1,179	2.4 (40)
Wolfe	395	6.3(112)	252	0.8(114)	246	0.4(114)
Woodford	267	18.0 (16)	193	0.5(117)	188	0.0(118)
Unknown Counties	2,366	11.1 (69)	53	* *	55	* *
Kentucky	123,028	13.6	71,711	3.5	65,968	2.3

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.