




Kentucky Public Health
Prevent. Promote. Protect.

54 Kentucky Counties with Increased Vulnerability to Rapid Dissemination of HIV/HCV Infections Among People who Inject Drugs and Preventive Syringe Services Programs (SSPs)

220 U.S. Counties at Risk



National Ranking by County*

1	Wolfe	34	Martin	108	Gallatin
3	Breathitt	35	Boyle	125	Bath
4	Perry	39	Lawrence	126	Grayson
5	Clay	40	Rockcastle	129	Greenup
6	Bell	45	Harlan	132	Green
8	Leslie	48	McCreary	153	Casey
9	Knox	50	Letcher	154	Carter
10	Floyd	53	Johnson	163	Monroe
11	Clinton	54	Russell	167	Garrard
12	Owsley	56	Elliott	175	Robertson
14	Whitley	65	Laurel	178	Lewis
15	Powell	67	Carroll	179	Edmonson
17	Knott	75	Taylor	180	Allen
21	Pike	77	Grant	187	Boyd
23	Magoffin	93	Adair	191	Hickman
25	Estill	97	Lincoln	202	Breckinridge
30	Lee	99	Wayne	212	Campbell
31	Menifee	101	Cumberland	214	Mercer

* Vulnerable Counties in RED have Operating SSPs

Specific concerns regarding Kentucky Counties:

1. Dense drug user networks similar to Scott County, Indiana



54 Vulnerable Counties



73 Operating SSPs
(82 Counties) as of 7/27/2020

A large, stylized red ribbon graphic that loops around the top left and bottom of the main text.

**KNOW YOUR
STATUS**


Eastern Kentucky

get HIV tested





**Even if a mother has HIV, her baby
doesn't have to follow in her footsteps.**



CDC Resource Kit: HIV Screening of Pregnant Women





KENTUCKY


Since the beginning of the HIV epidemic 1982 there has been

- 95 pediatric HIV case reported to the Kentucky HIV/AIDS surveillance program.
- 77% were due to perinatal transmission through an HIV-infected mother.
- Eleven (11) cases were reported with a primary exposure route of pediatric hemophilia or coagulation disorders.
- Two (2) ca

it



Spring of 2021, I will be having a One Test Two Lives Campaign. Including billboards and posters throughout Ky to raise awareness.





ONE TEST TWO LIVES

Prevent HIV Kentucky

Contact your local Health Department



KentuckyPublicHealth
Prevent. Promote. Protect.

FAIRWAY

Should I get tested for HIV?

CDC recommends everyone between the ages of 13 and 64 get tested for HIV at least once.

bioLytical® Laboratories develops and **manufactures** a rapid **HIV** diagnostic product, the **INSTI**. At 60 seconds for a reading, **INSTI** is the fastest **HIV** diagnostic test in the world today and is 99.6% accurate



HIV HOT LINE: 1-844-294-2448



JAMI MCLEMORE

PROGRAM COORDINATOR

KENTUCKY DEPARTMENT FOR PUBLIC
HEALTH

HIV/AIDS SECTION

JAMI.MCLEMORE@KY.GOV

(502) 564-6539 EXT.4284



Kentucky Public Health
Prevent. Promote. Protect.
