


**Table 2-G**  
**Resident Deaths from Influenza and Pneumonia**  
**Number and Crude Rate\*: Kentucky, 1960 - 2004**

Year	Number	Rate	Year	Number	Rate	Year	Number	Rate
1960	1,407	46.3	1975	1,031	30.7	1990	1,405	37.7
1961	946	31.0	1976	1,016	30.0	1991	1,439	39.0
1962	1,005	32.6	1977	935	27.3	1992	1,355	36.5
1963	1,630	53.2	1978	1,076	31.1	1993	1,536	40.9
1964	1,019	33.5	1979	802	22.9	1994	1,510	39.5
1965	1,135	37.0	1980	988	28.0	1995	1,476	38.2
1966	1,091	34.7	1981	1,004	27.4	1996	1,350	34.8
1967	1,080	33.9	1982	838	22.6	1997	1,346	34.4
1968	1,254	39.2	1983	964	26.1	1998	1,496	38.0
1969	1,222	38.0	1984	1,025	27.6	1999**	604	15.3
1970	1,130	35.1	1985	1,268	34.1	2000	834	20.6
1971	958	29.5	1986	1,262	33.9	2001	997	24.5
1972	1,099	33.5	1987	1,117	30.0	2002	1,223	29.9
1973	1,086	32.8	1988	1,340	36.0	2003	1,030	25.0
1974	938	28.2	1989	1,309	35.1	2004	970	23.4

\*Crude Rate per 100,000 Total Population



\*\* The decrease after 1998 is partially the result of coding differences between ICD-9 and ICD-10.